

# Microbial diversity analyses in a changing landscape

## Lactic acid bacteria in food fermentations as a test case

MSc. Isabel Snauwaert




Dissertation submitted in fulfillment of the requirements for the degree of  
*Doctor (Ph.D.) in Sciences, Biochemistry and Biotechnology (Ghent University)*  
*Doctor (Ph.D.) in Bioengineering Sciences (Vrije Universiteit Brussel)*

Promotors: Prof. dr. Peter Vandamme (Ghent University)  
Prof. dr. ir. Luc De Vuyst (Vrije Universiteit Brussel)  
Prof. dr. Anita Van Landschoot (Ghent University)



I feel more microbe than man



Snauwaert, I. | Microbial diversity analyses in a changing landscape: Lactic acid bacteria in food fermentations as a test case

©2014, Isabel Snauwaert

ISBN-number: 978-9-4619724-0-8

All rights reserved. No part of this thesis protected by this copyright notice may be reproduced or utilised in any form or by any means, electronic or mechanical, including photocopying, recording or by any information storage or retrieval system without written permission of the author.

Printed by University Press | <http://www.universitypress.be>

Joint Ph.D. thesis, Faculty of Sciences, Ghent University, Ghent, Belgium  
Faculty of Sciences and Bioengineering Sciences, Vrije Universiteit Brussel, Brussels, Belgium

Publicly defended in Ghent, Belgium, November 25<sup>th</sup>, 2014

This Ph.D. work was supported by FWO-Flanders, BOF project, and the Vrije Universiteit Brussel (SRP, IRP, and IOF projects).

**Author's email address:** [isabelsnauwaert@hotmail.com](mailto:isabelsnauwaert@hotmail.com)



# Examination Committee

**Prof. dr. Savvas SAVVIDES** (Chairman)

L-Probe: Laboratory for Protein Biochemistry and Biomolecular Engineering  
Faculty of Sciences, Ghent University, Ghent, Belgium

**Prof. dr. Peter VANDAMME** (Promotor UGent)

LM-UGent: Laboratory of Microbiology  
Faculty of Sciences, Ghent University, Ghent, Belgium

**Prof. dr. ir. Luc DE VUYST** (Promotor VUB)

IMDO: Research Group of Industrial Microbiology and Food Biotechnology  
Faculty of Sciences and Bioengineering Sciences, Vrije Universiteit Brussel,  
Brussels, Belgium

**Prof. dr. Anita VAN LANDSCHOOT** (Promotor UGent)

Laboratory of Biochemistry and Brewing  
Faculty of Bioscience Engineering, Ghent University, Ghent, Belgium

**Prof. dr. Edilbert VAN DRIESSCHE** (Secretary)

SPRO: Research Group of Protein Chemistry  
Faculty of Sciences and Bioengineering Sciences, Vrije Universiteit Brussel,  
Brussels, Belgium

**Prof. dr. Tom COENYE**

LPM: Laboratory of Pharmaceutical Microbiology  
Faculty of Pharmaceutical Sciences, Ghent University, Ghent, Belgium

**Prof. dr. Anne WILLEMS**

LM-UGent: Laboratory of Microbiology  
Faculty of Sciences, Ghent University, Ghent, Belgium

**Prof. dr. ir. Ronnie WILLAERT**

SBB: Research Group of Structural Biology Brussels  
Faculty of Sciences and Bioengineering Sciences, Vrije Universiteit Brussel,  
Brussels, Belgium

**Prof. dr. Tom LENAERTS**

MLG: Machine Learning Group  
Faculty of Sciences, Université Libre de Bruxelles, Brussels, Belgium

AI lab: Artificial Intelligence lab

Faculty of Sciences and Bioengineering Sciences, Vrije Universiteit Brussel,  
Brussels, Belgium

**Prof. dr. Sebastiaan EELTINK**

Research Group of Chemical Engineering  
Faculty of Engineering, Vrije Universiteit Brussel, Brussels, Belgium

**Prof. dr. Rudi VOGEL**

Lehrstuhl für Technische Mikrobiologie  
Center of Life and Food Sciences Weihenstephan, Technische Universität  
München, München, Germany

# Dankwoord

Vijf jaar heb ik als een 'Bezig Bijtje' in de gangen van het laboratorium voor Microbiologie rondgecrosst. Al die jaren heb ik mezelf kunnen zijn en me zo maximaal kunnen ontplooien. Mijn dankbaarheid uitdrukken op papier is dus geen eenvoudige opdracht en is sowieso ontoereikend. Maar ik doe hier toch een poging.

*Professor Vandamme*, Peter, allereerst wil ik u bedanken omdat u me de kans hebt gegeven om dit onderzoek uit te voeren. Bedankt om vertrouwen te hebben in mijn kunnen, om mij vrijheid te geven in de invulling van het project en om mij naar vele workshops en congressen te laten gaan. Dankzij u heb ik veel kunnen bijleren. *Professor De Vuyst*, bedankt om mij als één van uw doctoraatstudenten te behandelen. Zo nodigde u me uit om samen te lunchen met het IMDO-team en zelfs bij het kerstetentje, waardoor ik me bijzonder welkom voelde. *Professor Van Landschoot*, dankzij uw connecties met de brouwerijen konden we de nodige stalen verzamelen. Simpelweg onmisbaar voor mijn onderzoek.

*Lieve buurvrouwtjes*, jullie waren mijn warme thuisbasis binnen het labo. Onze bureau was een plaatsje waar ik veel gelachen heb, waar ik raad kon vragen, waar ik iedere morgen goed gezind werd en waar ik ook even triest mocht zijn. Jullie zijn stuk voor stuk top-collega's en vriendinnen. *Charlie*: mijn partner in crime én mijn tweede brain. Nooit gedacht dat iemand mijn gek idee ook effectief mee ging uitvoeren: op een workshop in Detroit om 4 uur in de ochtend opstaan om nog voor de start te kunnen fitnessen, ontbijten én scriptjes schrijven. En dan s' avonds vroeg in bed, want er was toch niets te doen in Detroit. Gelukkig wel in Liverpool: de cafeetjes waren er zo gezellig en het bier zeer drinkbaar. Maar nog belangrijker: brainstormen met jou was pure synergie, we brachten elkaar steeds tot een hoger niveau. Dankzij jou was er geen probleem dat niet tot een oplossing kon komen.

*Anne, Timo, Jindrich, Freek, Jessy, Anneleen, Bram V., en Helen*, mijn maatjes, veel meer dan alleen collega's. Samen gaan eten, op café gaan en kletsen kon alle dipjes doen overwaaien en gaf mij weer moed. Heerlijk om jullie te leren kennen! *Gwen, Sofie, Joke*, en *Jonas*: wat deed het pijn om jullie één voor één te zien vertrekken na het beëindigen van jullie doctoraat. *Gwen*, zonder jou was ik zelfs nooit doctorandus geworden. Je hebt als mijn stagebegeleider je enthousiasme, gedrevenheid en relativeringsvermogen op mij geprojecteerd. Door jou kreeg ik definitief de microbiologie-microbe te pakken. *Sofie*, jij was en bent nog steeds een onuitputtelijke bron van ideeën voor leuke activiteiten. En *Joke* en *Jonas*, ook jullie bleef ik tot op vandaag missen. *Kim*, bij de start van mijn doctoraat had ik het gevoel had dat je niet volledig geloofde in mijn capaciteiten. Maar achteraf ben ik daar blij om. Want zo heb je me assertiever en sterker gemaakt, want ik wilde bewijzen dat ik het wel kon. Een waardevolle attitude die ik nu meeneem naar het bedrijfsleven. En ook je lieve en dankbare tips voor de combinatie werk én zorg voor een pittige dochter, zullen mij altijd bij blijven.

Het schrijven van dit dankwoord toont aan dat er zoveel warme persoonlijkheden bij LM-Ugent werken. Zoals al eerder gezegd: dit dankwoord is sowieso ontoereikend. Zo denk ik nog aan *Jeannine, Evie, Bart H., Liesbeth, Maarten, Leilei, Bart V., en Joke* en alle andere collega's van LM-Ugent. Bedankt voor de warme en hartelijke sfeer in het labo, ik was er elke dag opnieuw welkom.

Lieve IMDO collega's, als bezoeker in een 'vreemd' laboratorium komen is niet evident. Ik was plots weer de leek die niets wist liggen en iedereen lastig viel met mijn vragen. Maar ik kon steeds bij jullie terecht. Expliciet een merci aan *Tom, Audrey, Maria, David, Henning, Moens* en *Koen*. Na twee uur trainen van Brugge naar Etterbeek voelde ik me vaak suf (zelfs koffie hielp dan niet). Maar na aankomst, kregen jullie mij altijd weer vrolijk.

Het doctoraat mocht dan wel boeiend zijn, maar gelukkig is er in mijn leven ook nog meer. Lieve *Lindseys*, het is altijd heerlijk om samen met mijn vriendinnen wijntjes te drinken en te kletsen. Jullie zijn stuk voor stuk topwijken. *Ilse*, mijn capaciteit om te plannen en efficiënt te werken was even zoek geraakt de laatste twee maanden. Merci om mij toen weer op de rails te zetten. En ook jij bedankt, *mevrouw Mitchell*, omdat je gewoon dé meest attente persoon bent die ik ken.

*Mama en papa*, bedankt voor alle kansen die jullie mij gaven. Jullie bleven mij onvoorwaardelijk steunen en geloofden in mijn capaciteiten. Door jullie kon ik studeren wat ik maar wilde en kan ik nu terugkijken op een heerlijke studententijd. Kortom, zonder jullie liefde en vertrouwen kan ik niet. *Papa*, ooit lachte ik met je als academische 'nerd' en 'pietje precies', maar ik vrees dat ik deze eigenschappen zeker van je meekreeg. *Mama*, ik hoop dat ik van jou het 'mama-zijn' heb geërfd en voor mijn dochter Helena nu net zo'n supermama zal zijn zoals jij altijd voor mij bent geweest (en nog altijd bent).

Ook *Evelien* (mijn ideale zus), *Filip* (de eeuwige enthousiasteling), *Sanne* (R en latex kenner: heerlijk om iemand met deze vaardigheden in de familie te hebben), *Mathilde* (altijd paraat als babysit) en mijn schoonfamilie: - *Jan* (English Expert), *Hilde* (de onmisbare SOS oma), de gekke *Leens*, *Bert* en *Pieter* - : dikke merci!

Het zijn vijf leerrijke jaren geweest, maar toch ben ik enorm opgelucht dat ik eindelijk een punt achter dit werk mag zetten. Het is nu tijd om mij te focussen op de belangrijke dingen in het leven en dat zijn jullie -*Bram* en kleine *Helena*-. *Bram*, jij hebt gevoeld hoe zwaar het afwerken van het doctoraat op mijn schouders woog na de geboorte van ons klein spook. Ik heb je de voorbije maanden meer dan ooit nodig gehad en jij bent er steeds voor mij geweest. Jouw onvoorwaardelijke liefde is het mooiste dat ik heb mogen tegengekomen in mijn leven. *Lieve dochter*, mijn klein pittig vrouwtje, ik kijk ernaar uit om je te zien opgroeien en je heel erg veel liefde te geven.

*Gent, 17 november 2014*  
*Isabel Snauwaert*





# Table of Contents

Examination Committee	i
Dankwoord	iii
Table of Contents	vii
List of Figures	xi
List of Tables	xiii
Nederlandstalige Samenvatting	xxiii
English Summary	xxv
<b>I Introduction</b>	<b>1</b>
Background & Objectives	3
Outline	5
<b>II Literature Overview</b>	<b>7</b>
<b>1 Microbial Diversity Analyses of Food Fermentations</b>	<b>9</b>
1.1 Traditional Methods . . . . .	10
1.1.1 Plate Culturing . . . . .	11
1.1.2 Cultivation-Independent Molecular Techniques . . . . .	12
1.2 Microbial Diversity Analyses in the HT Sequencing Era . . . . .	14
1.2.1 Platforms . . . . .	15
1.2.2 (Meta)genomics . . . . .	17

<b>2</b>	<b>Importance of LAB in Food Fermentations</b>	<b>23</b>
<b>3</b>	<b>LAB Taxonomy</b>	<b>27</b>
3.1	Phylogenetics . . . . .	28
3.2	Polyphasic Taxonomy . . . . .	31
3.3	Genomic Taxonomy . . . . .	32
<b>III</b>	<b>Experimental Work</b>	<b>35</b>
<b>4</b>	<b>Traditional Workflow in Microbial Diversity Analyses</b>	<b>37</b>
4.1	Characterization of Strains of <i>Weissella fabalis</i> sp. nov. and <i>Fructobacillus tropaeoli</i> from Spontaneous Cocoa Bean Fermentations .	39
4.1.1	Introduction . . . . .	40
4.1.2	Methods, Results, and Discussion . . . . .	40
4.1.3	Conclusion . . . . .	49
4.1.4	Acknowledgements . . . . .	51
4.1.5	Supplementary Material . . . . .	52
4.2	<i>Leuconostoc rapi</i> sp. nov., Isolated from Sous-Vide Cooked Rutabaga	55
4.2.1	Introduction . . . . .	55
4.2.2	Methods, Results, and Discussion . . . . .	56
4.2.3	Conclusion . . . . .	61
4.2.4	Acknowledgements . . . . .	62
4.2.5	Supplementary Material . . . . .	62
4.3	<i>Carnobacterium iners</i> sp. nov., a Psychrophilic, Lactic Acid-Producing Bacterium from the Littoral Zone of an Antarctic Pond. . . . .	63
4.3.1	Introduction . . . . .	64
4.3.2	Methods, Results, and Discussion . . . . .	64
4.3.3	Conclusion . . . . .	71
4.3.4	Acknowledgements . . . . .	72
4.3.5	Supplementary Material . . . . .	72
<b>5</b>	<b>Microbial Diversity Analyses in the HT Sequencing Era</b>	<b>73</b>
5.1	Microbial Diversity and Metabolite Composition of Belgian Red-Brown Acidic Ales . . . . .	74
5.1.1	Introduction . . . . .	75
5.1.2	Materials and Methods . . . . .	76
5.1.3	Results . . . . .	82
5.1.4	Discussion . . . . .	95
5.1.5	Acknowledgments . . . . .	100
5.1.6	Supplementary Material . . . . .	100



5.1.7	Data Analysis Pipeline . . . . .	108
5.2	Comparative Genome Analysis of <i>Pediococcus damnosus</i> LMG 28219, a Strain Well-Adapted to the Beer Environment . . . . .	111
5.2.1	Introduction . . . . .	112
5.2.2	Materials and Methods . . . . .	113
5.2.3	Results and Discussion . . . . .	115
5.2.4	Conclusions . . . . .	129
5.2.5	Acknowledgments . . . . .	129
5.2.6	Supplementary Material . . . . .	129
<b>IV</b>	<b>General Discussion and Future Perspectives</b>	<b>143</b>
<b>6</b>	<b>Considerations for Microbial Diversity Analyses</b>	<b>147</b>
6.1	Traditional Methods . . . . .	147
6.2	Methods Involving HT Sequencing . . . . .	148
6.2.1	Pre-Sequencing Considerations . . . . .	148
6.2.2	Post-Sequencing Considerations . . . . .	152
6.3	Importance of Cultivation . . . . .	156
6.4	Integrative Ecosystems Biology . . . . .	159
<b>7</b>	<b>LAB Taxonomy in a HT Sequencing Era</b>	<b>161</b>
7.1	Polyphasic Taxonomy . . . . .	161
7.2	Genomic Taxonomy . . . . .	163
7.3	Integrating Genomics into LAB Taxonomy . . . . .	164
<b>8</b>	<b>Future Perspectives</b>	<b>167</b>
<b>V</b>	<b>Bibliography</b>	<b>171</b>
<b>VI</b>	<b>Curriculum Vitae</b>	<b>199</b>



## List of Figures

1	Overview of the techniques used for microbial community analyses of food samples . . . . .	10
2	HT sequencing platforms . . . . .	16
3	PCR as a target enrichment strategy . . . . .	21
4	Major pathways active in LAB . . . . .	24
5	Consensus dendrogram reflecting the phylogenetic relationships of a selection of LAB genera of the order <i>Lactobacillales</i> within the class <i>Bacilli</i> . . . . .	25
6	Schematic overview of the major analytical approaches to phylogenetic tree building. . . . .	29
7	ML tree based on 16S rRNA gene sequences showing the phylogenetic relationships of strains M75 <sup>T</sup> and M622 among the type strains of all species of the genera <i>Weissella</i> and <i>Fructobacillus</i> . .	41
8	ML tree based on concatenated <i>pheS</i> , <i>rpoA</i> , and <i>atpA</i> gene sequences showing the phylogenetic relationships of strains M56, M622, M710, M1190, and M1588 among other members of the genus <i>Fructobacillus</i> . . . . .	45
9	ML tree based on <i>pheS</i> gene sequences showing the phylogenetic relationships of strain M75 <sup>T</sup> among other members of the genus <i>Weissella</i> . . . . .	46
10	ML tree of the 16S rRNA gene sequence of strain LMG 27676 <sup>T</sup> with all type strains of the genus <i>Leuconostoc</i> . . . . .	57
11	ML tree of the concatenated <i>pheS</i> , <i>rpoA</i> , and <i>atpA</i> gene sequences of the closest relatives of strain LMG 27676 <sup>T</sup> . . . . .	59
12	ML tree based on 16S rRNA gene sequences showing the phylogenetic relationship of strains LMG 26642 <sup>T</sup> and LMG 26641 with type strains of species of the genus <i>Carnobacterium</i> . . . . .	65

13	ML tree based on concatenated <i>pheS</i> , <i>rpoA</i> , and <i>atpA</i> sequences showing the phylogenetic relationship of strains LMG 26642 <sup>T</sup> and LMG 26641 with the type strains of species of the genus <i>Carnobacterium</i> . . . . .	67
14	1/D of the bacterial (A) and fungal (B) datasets of brew samples from breweries 1, 2, and 3 . . . . .	84
15	Relative abundances of the most abundant OTUs (> 0.01% of the reads) of the bacterial (A) and fungal (B) communities . . . . .	86
16	PCA plot (A) and heatmap (B) of the metabolite concentrations of breweries 1, 2, and 3, determined in brew samples from the end of the maturation phase . . . . .	89
17	PCA plot (A) and heatmap (B) of the metabolite concentrations of the bottled beers from breweries 1, 2, and 3 . . . . .	93
18	Visualization of the OrthoMCL output comparing the number of unique and/or shared orthologs of <i>P. clausenii</i> ATCC BAA-344 <sup>T</sup> , <i>P. pentosaceus</i> ATCC 25745, and <i>P. damnosus</i> LMG 28219 . . . . .	121
19	Comparison of plasmid pPECL-8 from <i>P. clausenii</i> ATCC BAA 344 <sup>T</sup> and contigs 70, 14, and 90 of <i>P. damnosus</i> LMG 28219 . . . . .	122
20	Folate biosynthesis pathway reconstruction . . . . .	125
21	Overview of the functions enriched in the genomes of LAB of beer origin compared to those of LAB not originating from beer . . . . .	128
22	Schematic outlining of the general approaches to culturing microbial diversity . . . . .	158
23	Microbial diversity analyses in a HT sequencing era . . . . .	168

## List of Tables

1	Comparison of HT sequencing platforms . . . . .	18
2	<i>Weissella</i> and <i>Fructobacillus</i> strains included in the phylogenetic study and in the MALDI-TOF MS analysis. . . . .	43
3	Differential characteristics of strain M75 <sup>T</sup> ( <i>W. fabalis</i> sp. nov.) and other species of the genus <i>Weissella</i> . . . . .	50
4	Differential phenotypic characteristics of strain LMG 27676 <sup>T</sup> and closest related species of the genus <i>Leuconostoc</i> . . . . .	61
5	Amplification and sequencing primers used for the MLSA of <i>Carnobacterium</i> strains . . . . .	67
6	Members of the genus <i>Carnobacterium</i> examined in this study . . .	68
7	Differential characteristics of strain LMG 26642 <sup>T</sup> and members of the genus <i>Carnobacterium</i> that assimilate a limited range of carbohydrates . . . . .	70
8	Overview of the <i>P. damnosus</i> LMG 28219 draft genome. . . . .	117
9	Comparison of genome characteristics of <i>P. damnosus</i> LMG 28219 with <i>P. pentosaceus</i> ATCC 25745 and <i>P. claussenii</i> ATCC BAA-344 <sup>T</sup>	120



# List of Acronyms

## A

A	aerobic
AAB	acetic acid bacteria
AAI	average amino acid identity
ACA	American coolship ales
ADI	arginine deiminase pathway
ADP	adenosine diphosphate
AMP	adenosine monophosphate
ANI	average nucleotide identity
ANOSIM	analysis of similarity
ARISA	automated ribosomal intergenic spacer analysis
ATCC	American type culture collection
ATP	adenosine triphosphate
atpA	RNA polymerase $\alpha$ -subunit

## B

bcaT	branched chain amino acid aminotransferase
BLAST	basic local alignment tool
bp	base pair
BPW	buffered peptone water

## C

C.	<i>Carnobacterium</i>
CBA	columbia blood agar
CCA	canonical correlation analysis
CCMM	Moroccan coordinated collections of microorganisms
CDS	coding sequence
CECT	colección Española de cultivos tipo
CAMERA	community cyberinfrastructure for advanced marine microbial ecology research and analysis
CNS	coagulase-negative staphylococci
Cq	quantification cycle

## D

1/D	inverse Simpson diversity index
DDBJ	DNA data bank of Japan
DDH	DNA-DNA hybridization
DGGE	denaturing gradient gel electrophoresis
DNA	deoxyribonucleic acid
dNTP	deoxynucleotide
DSMZ	Deutsche sammlung von mikroorganismen und zellkulturen
dps	glucosyl transferase
DYPA	general yeast agar isolation medium
DYPAX	DYPA supplemented with 50 ppm cycloheximide

## E

EC	enzyme commission
ELSD	evaporative light-scattering detector
ENA	European nucleotide archive
EPS	exopolysaccharide
ERDF	European regional development fund



---

## F

F.	<i>Fructobacillus</i>
FAME	fatty acid methyl ester
FISH	fluorescence <i>in situ</i> hybridization
FWO	fund for scientific research-Flanders

## G

GC	gas chromatography
GEBA	genomic encyclopaedia of bacteria and archaea
GOLD	genomes online database
groEL	molecular chaperone <i>groEL</i> gene
GSC	genomic standards consortium
gtf	transmembrane glycosyl transferase
GTP	guanoside triphosphate
GTR	general time reversible

## H

HMM	hidden markov model
HPAEC	high-performance anion exchange chromatography
HPLC	high-performance liquid chromatography
HT	high-throughput

## I

IDT	integrated DNA technologies
IMG	integrated microbial genomes
IMG-ER	IMG expert review
IMG/M	IMG/metagenomes
INSDC	International nucleotide sequence databases collaboration

IS	insertion sequence
ITS	internal transcribed spacer

## J

JCM	Japan collection of microorganisms
-----	------------------------------------

## K

KCTC	Korean collection for type cultures
KEGG	kyoto encyclopedia of genes and genomes

## L

L.	<i>Leuconostoc</i>
LAB	lactic acid bacteria
LaCOG	<i>Lactobacillales</i> -specific clusters of orthologous group
Lb.	<i>Lactobacillus</i>
LMG	laboratory of microbiology, Ghent University

## M

mA	microaerobic
MALDI	matrix-assisted laser desorption/ionization
MEGA	molecular evolutionary genetics analysis
MEGAN4	metagenome analyzer
METAREP	meta gene annotator
MG	metagenome
MG-RAST	metagenomic rapid annotation using subsystems technology
MID	multiplex identifier
MiXS	minimum information about any sequence

---

ML	maximum-likelihood
MLSA	multilocus sequence analysis
MP	maximum parsimony
MRS	de Man–Rogosa–Sharpe
MS	mass spectrometry
MUM	maximal unique matches

## N

NADP(H)	nicotinamide adenine dinucleotide phosphate
NCBI	National center for biotechnology information
NCFB	National collection of food bacteria
NCIMB	National collections of industrial, food and marine bacteria
ND	no data available
NK	not known
NJ	neighbour joining
no.	number

## O

OTU	operational taxonomic unit
-----	----------------------------

## P

P.	<i>Pediococcus</i>
pABA	4-aminobenzoic acid
PATRIC	pathosystems resource integration center
PCA	principal component analysis
pcDNA	percentage of conserved DNA
PCoA	principal coordinate analysis
PCR	polymerase chain reaction
PE	paired-end
pepN	aminopeptidase N
pepX	X-prolyl dipeptidyl peptidase

PGAAP	prokaryotic genome automatic annotation pipeline
PHAST	phage search tool
pheS	phenylalanyl-tRNA synthase $\alpha$ subunit
P(P)i	(pyro)phosphate

## Q

QIIME	quantitative insights into microbial ecology
qPCR	quantitative PCR

## R

RAPD	randomly amplified polymorphic DNA
RAST	rapid annotation using subsystems technology
RDP	ribosomal database project
rep-PCR	repetitive element sequence-based PCR
RNA	ribonucleic acid
rpoA	RNA polymerase $\alpha$ subunit
rpoB	$\beta$ -subunit of the bacterial RNA polymerase
rRNA	ribosomal RNA

## S

SCFA	short-chain fatty acids
SE	standard error
SH	static-headspace
sp.	species
SRA	sequence read archive

## T

T	type strain
---	-------------

TLC	thin layer chromatography
TOF	time-of-flight
TRF	terminal restriction fragment
TRFLP	terminal restriction fragment length polymorphism
tRNA	transfer RNA
TSA	tryptone soya agar
TSBY	trypticase soy broth with yeast extract salt medium

## U

UPGMA	unweighted pair group method with arithmetic means
UV	ultra violet

## V

VBNC	viable but non-cultivable
------	---------------------------

## W

W.	<i>Weissella</i>
----	------------------



# Nederlandstalige Samenvatting

Micro-organismen, inclusief melkzuurbacteriën, dragen bij tot de smaak, het aroma, de textuur, en de houdbaarheid van gefermenteerde levensmiddelen en dranken, die deel uitmaken van het dagelijks dieet van de mens. Kennis over de microbiële gemeenschappen in deze fermenterende ecosystemen is nodig om veilige producten met gewenste eigenschappen te kunnen afleveren aan de consument. Het doel van deze thesis is het evalueren van de huidige methoden om de microbiële diversiteit van voedsel ecosystemen te analyseren en om nieuwe soorten van melkzuurbacteriën te beschrijven.

Een grote fractie aan micro-organismen kunnen niet gekweekt worden door middel van conventionele cultivatie-condities, wat leidt tot een vertekend beeld van de werkelijke microbiële diversiteit van een ecosysteem wanneer cultivatie-afhankelijke methoden worden aangewend. Cultivatie-onafhankelijke moleculaire technieken (zoals '*denaturing gradient gel electrophoresis*', '*terminal restriction fragment length polymorphism*', '*clone libraries*', enz.) zijn niet afhankelijk van de kweekbaarheid van micro-organismen in een gemeenschap, maar deze zijn ongeschikt door hun arbeidsintensiviteit, tijdrovende natuur, en lage-throughput. Om deze redenen zijn effectieve methoden essentieel die het documenteren van de microbiota in levensmiddelen mogelijk maken.

De ontwikkeling van high-throughput sequenceringsmethoden en de resulterende sequenceringskosten aan lage kost per base hebben het microbiologisch onderzoek de laatste decennia drastisch veranderd. In deze thesis werd high-throughput sequenceringsmethode toegepast om het ecosysteem van Belgische roodbruine bieren te bestuderen en om het genoom van het dominante bacteriële lid van deze gemeenschap te sequencen. Daaruit blijkt dat high-throughput sequenceringsmethoden waardevol zijn voor bestuderen van de diversiteit en het metabolische potentieel van micro-organismen. Toch, door fouten, geïntroduceerd tijdens microbiële diversiteitsstudies die high-throughput sequenceringsmethode toepassen, is het essentieel om deze te overwegen vooraleer een experiment aan te vatten en tijdens de analyse van de data.

De traditioneel toegepaste polyfasische taxonomische benadering werd in deze thesis aangewend om nieuwe soorten van melkzuurbacteriën te beschrijven. Daaruit blijkt dat deze benadering gehinderd is door haar arbeidsintensieve en tijdrovende natuur, wat contrasteert met de grote hoeveelheid stammen die momenteel wachten om beschreven te worden. Daarbovenop contrasteert de polyfasische taxonomische benadering met de opkomende 'culturomics' aanpak, die het potentieel heeft om een grote hoeveelheid tot op heden niet gekarakteriseerde stammen te genereren. Om bovenstaande redenen is de polyfasische taxonomische benadering verouderd en moet deze dringend herbekeken worden om een snelle beschrijving van nieuwe soorten van melkzuurbacteriën in de toekomst mogelijk te maken.

Omdat volledige genoom sequenties inzichten verwerven in de genetische natuur van microbiële species, kunnen ze gebruikt worden om bacteriële species af te lijnen en om hun fylogenie te bestuderen. De beschrijving van nieuwe soorten in het genomische tijdperk zou een genoom sequentie en een minimale beschrijving van fenotypische eigenschappen moeten bevatten, die samen kunnen fungeren als een biologische identiteitskaart die voldoende, kostenbesparend, en geschikt is voor de beschrijving van nieuwe species. De opslag en het onderhoud van deze biologische identiteitskaarten zal resulteren in taxonomische databanken van hoge kwaliteit die de wetenschappelijke wereld van dienst kunnen zijn om de kwaliteit van hun annotaties en taxonomische toewijzingen te verbeteren. Bovendien zouden bacteriële taxonomen kunnen bijdragen aan de inspanningen van het '*Genomic Encyclopaedia of Bacteria and Archaea*'-project om de afwijkende fylogenetische verdeling van de beschikbare genoom-sequenties in te perken.

Finaal zou het combineren van (meta)genoom-, (meta)transcriptoom-, (meta)proteoom-, en (meta)metabool-data met cultivatie nieuwe inzichten kunnen verwerven in de interacties tussen de leden van een microbiële gemeenschap. Bovendien zou deze geïntegreerde aanpak belangrijk kunnen zijn om interventies te ontwerpen die gefocust zijn op de functies van de gemeenschappen in levensmiddelen in plaats van de afzonderlijke species.



# English Summary

Microorganisms, including lactic acid bacteria, contribute to the taste, aroma, texture, and shelf life of fermented foods and beverages, which are part of the human diet globally. Knowledge on the microbial communities within these fermenting ecosystems is required to deliver safe products with desirable properties. The goals of this thesis were to evaluate the state-of-the-art approaches for microbial diversity analyses of food ecosystems and for the description of novel lactic acid bacteria taxa.

Because a large fraction of the microbial consortia cannot be cultured using conventional growth conditions, culture-dependent approaches lead to a biased view on the actual microbial diversity. Culture-independent molecular techniques (*i.e.*, denaturing gradient gel electrophoresis, terminal restriction fragment length polymorphism, clone libraries, and others) do not depend on the cultivability of microorganisms in a community, but they are inadequate for large-scale microbial diversity analyses because of their labor-intensity, low-throughput, and time-consuming nature. Therefore, effective tools are required enabling thorough documentation of the microorganisms in food ecosystems.

The advent of high-throughput sequencing technologies and the resulting sequencing at reduced cost per base have changed microbiological research drastically in the last decade, making high-throughput sequencing affordable for many research laboratories. In this thesis, high-throughput sequencing technologies were applied to unravel the microbial diversity of mature Belgian red-brown acidic ales and to sequence the genome of the dominant bacterial member of this beer ecosystem. These high-throughput sequencing technologies proved to be valuable for studying the diversity and the metabolic potential of environmental microorganisms. Nevertheless, because of the biases associated with high-throughput sequencing, it is crucial to consider these before launching an experiment and while handling the data.

The traditionally used polyphasic taxonomy approach was applied to describe novel lactic acid bacteria species and proved to be labor-intensive and thus unsuited to perform in high-throughput, which is contradictory with the huge amount of strains that are currently awaiting description and formal naming. Furthermore, this approach contrasts with the emerging culturomics method, which has the potential to generate a vast amount of not yet characterized strains. As a consequence, the polyphasic taxonomy approach is outdated and should urgently be reconsidered to enable fast description of novel lactic acid bacteria taxa.

Because whole-genome sequences provide insights into the genetic nature of microbial species, they can be used as a tool for delineating bacterial species and for studying their phylogeny. The genomic taxonomy approach should include a whole-genome sequence and a minimal description of phenotypic characteristics, which will serve as a basic biological identity card that be considered sufficient, cost-effective, and appropriate for species descriptions. These biological identity cards should be stored and maintained in high-quality, readily accessible, iterative, and adaptable taxonomic databases, which can serve the scientific community to improve the quality of their annotations and taxonomic assignments. Furthermore, bacterial taxonomists could contribute to the phylogenetically driven sequencing effort, initiated by the Genomic Encyclopaedia of Bacteria and Archaea project, and thereby reduce the biased phylogenetic distribution of the currently available genome sequences.

Finally, combining (meta)genome, (meta)transcriptome, (meta)proteome, and (meta)-metabolome approaches with cultivation could provide new insights into relations between members of a microbial community and will be important for designing interventions targeted at functions of the microbial community in food rather than specific constituent species.

# Part I

## Introduction





# Background & Objectives

Microorganisms, including lactic acid bacteria (LAB), contribute to the taste, aroma, texture, and shelf life of fermented foods and beverages, which are part of the human diet globally. Knowledge on the microbial communities within these fermenting ecosystems is required to deliver safe products with desirable properties. A large fraction of these microbial consortia cannot be cultured using conventional cultivation conditions, leading to a biased view on the actual microbial diversity when applying traditional approaches. Therefore, effective tools are required enabling documentation of the microorganisms in food ecosystems. Developments in high-throughput (HT) sequencing technologies provide a world of opportunities for investigating these ecosystems. Furthermore, the currently used polyphasic taxonomy approach is hampered by its labor-intensive and time-consuming nature, and microbiologists are therefore looking for alternatives based on the evolutionary information contained in whole-genome sequences.

The first goals of the present study were to evaluate (i) the state-of-the-art approaches for microbial diversity analyses applied to food ecosystems and (ii) the polyphasic taxonomy approach currently used for the description of novel species. Furthermore, (iii) the existing multilocus sequence analysis (MLSA) approach was extended to the genus *Carnobacterium*, which includes food-grade LAB species. Next, (iv) a HT sequencing approach, comprising targeted sequencing of the bacterial partial 16S ribosomal ribonucleic acid (rRNA) gene and the fungal internal transcribed spacer (ITS) region, was applied to study the ecosystem of mature Belgian red-brown acidic ales. The goal was to update and broaden the knowledge on a previously studied Belgian red acidic ale and to compare its microbial community diversity and metabolite composition with two additional Belgian red-brown acidic ales. Following, (v) whole-genome sequencing of *Pediococcus damnosus* LMG 28219, which was the dominant LAB member of the Belgian red-brown acidic ale ecosystem, and comparative genome analysis were used to investigate this strain's mechanisms to reside in the beer niche. Finally, (vi) the value of HT sequencing technologies in microbial diversity analyses was evaluated and the advantages of incorporating genome information into the field of taxonomy were explored.



# Outline

In **PART II** of the present thesis, a comprehensive LITERATURE OVERVIEW is given on microbial diversity analyses in a changing landscape, with special focus on LAB in food fermentations. **Chapter 1** describes the developments of molecular technologies for investigating the diversity of microorganisms in fermented foods and beverages, with a particular emphasis on the application of novel HT sequencing methods. **Chapters 2 and 3** review the importance of LAB in food fermentations and their taxonomy. **Chapter 3.1** provides background information on phylogenetics.

**PART III** presents the EXPERIMENTAL WORK performed during this thesis. In **Chapter 4**, the traditional workflow for microbial diversity analyses was applied to the spontaneous cocoa bean fermentation ecosystem and the sous-vide cooked rutabaga ecosystem. Novel LAB species (*i.e.*, *Weissella fabalis* sp. nov and *Leuconostoc rapi* sp. nov) discovered during these analyses, were described using a polyphasic taxonomy approach. Furthermore, the MLSA scheme for the genus *Carnobacterium* was completed, along with a polyphasic description of *Carnobacterium iners* sp. nov. **Chapter 5** comprises the application of HT sequencing technologies to unravel the microbial diversity of Belgian red-brown acidic ales, along with a comparative genome analysis of the *Pediococcus damnosus* LMG 28219 dominant LAB species, isolated from a Belgian red acidic ale, with other *Pediococcus* genome sequences.

**PART IV** presents a GENERAL DISCUSSION of the results and provides FUTURE PERSPECTIVES.





## Part II

### Literature Overview





# 1

## Microbial Diversity Analyses of Food Fermentations

Fermentation is a biotechnological method historically arisen from the need to preserve food. It can be carried out by bacteria, yeasts, and/or filamentous fungi, which convert fermentable carbohydrates mainly into organic acids, alcohols, and/or carbon dioxide (Leroy & De Vuyst, 2004; Ravyts *et al.*, 2012). LAB play a central role in the production of fermented foods and beverages, because of their production of lactic acid as a common end-metabolite (Stiles & Holzapfel, 1997). Next to LAB, other bacterial species have importance in food fermentations. For instance, acetic acid bacteria (AAB) contribute to the production of vinegar and cocoa (Raspor & Goranovic, 2008). Furthermore, coagulase-negative staphylococci (CNS), *Kocuria*, and *Micrococcus* species are indispensable natural microbiota of fermented meat products next to lactobacilli and pediococci (Leroy *et al.*, 2006) and can also be found in both hard and soft cheeses (Irlinger, 2008). Other bacterial groups with importance in food fermentations include brevibacteria, corynebacteria, and propionibacteria, which are used in cheese production, and *Bacillus subtilis* that is used for soybean fermentations (Ravyts *et al.*, 2012). In carbohydrate-rich environments, acidifying LAB frequently grow in association with yeasts that are present in lower numbers. Yeast species of *Saccharomyces*, *Candida*, *Torula*, *Hanseniaspora*, *Hansenula*, *Dekkera*, and others, can proliferate in these niches, causing spontaneous alcoholic fermentation of wine and beer (Fleet, 2007). Finally, filamentous fungi, such as *Aspergillus* and *Rhizopus* may play a role, such as in cheeses, fermented sausages, and soy-based fermentations (Cocolin *et al.*, 2013).

Searching for the presence, numbers, and type of microorganisms in foods is of great importance for the food industry. Whatever the primary objective of these microbial analyses (e.g., control of food quality or preservation, efficiency of starter cultures, monitoring of particular species/strains, detection of unwanted microorganisms), the taxonomic level of the microbial discrimination needed should be initially decided. An overview of the methods used in microbial diversity analyses is presented in Figure 1; the traditionally applied methods are discussed in Section 1.1, whereas those involving HT sequencing are elaborated in Section 1.2.




Figure 1: Overview of the techniques used for microbial community analyses of food samples

I: Shotgun sequencing strategy, II: Target enrichment strategy. Abbreviations are explained in Section 1.1.

## 1.1 Traditional Methods

A variety of culture-dependent and culture-independent techniques are traditionally used in microbial diversity analyses to study the phylogenetic and functional diversity of food ecosystems.

### 1.1.1 Plate Culturing

Microbial diversity analyses traditionally involve the cultivation of microorganisms onto a set of general and selective growth media and conditions. Following incubation, a dedicated percentage of colonies with morphologically distinct types are randomly picked up –whether or not statistically relevant– for further analysis. As bacteria are a convenient source of intrinsic marker proteins, matrix-assisted laser desorption/ionization time-of-flight mass spectrometry (MALDI-TOF MS) is an appropriate tool for the HT generation of strain-specific spectral profiles (Freiwald & Sauer, 2009; Ghyselinck *et al.*, 2011). By comparing these spectral profiles, a large set of isolates can easily be grouped, which allows the selection of a smaller set of representative strains, the so-called dereplication step (Ghyselinck *et al.*, 2011). Dereplication is important in large-scale isolation campaigns and screening programs, since it can significantly reduce labor intensity and time, and costs in further downstream analyses. Other tools used in the context of dereplication are repetitive element sequence-based polymerase chain reaction (rep-PCR) (De Clerck & De Vos, 2002), randomly amplified polymorphic deoxyribonucleic acid (RAPD) (Martín-Platero *et al.*, 2009), and fatty acid methyl ester (FAME) (Coorevits *et al.*, 2008) analyses. The resulting reduced set of isolates is assumed to be representative for the microbial communities studied and is subsequently identified using Sanger sequencing (Sanger *et al.*, 1992) of marker genes. Alternatively, a tentative identification can be obtained by querying MALDI-TOF MS protein profiles of this reduced set against an in-house developed or a commercial reference database containing high-quality spectra (Andres-Barrao *et al.*, 2013; Clark *et al.*, 2013; De Bruyne *et al.*, 2011; Dieckmann *et al.*, 2005; Duskova *et al.*, 2012; Wieme *et al.*, 2014). Examples of microbial diversity analyses of food ecosystems involving cultivation are numerous [as example (Spitaels *et al.*, 2014b; Papalexandratou *et al.*, 2011a; Laureys & De Vuyst, 2014; Pothakos *et al.*, 2014; Nguyen *et al.*, 2013)].

This labor-intensive plate culturing approach gives a biased view on the actual microbial diversity, because of the inability of certain community members to grow on the media or in the incubation conditions selected, and because of the challenging recovery of microorganisms, which are either stressed or entered into a viable but non-cultivable (VBNC) state (Fleet, 1999; Rappé & Giovannoni, 2003). The VBNC state is induced when adverse conditions such as nutrient depletion, low temperatures, and stresses cause cultivable cells to enter a phase, in which they are still capable of metabolic activity but do not produce colonies on media that normally would support their growth. As a consequence, the cultivation-based approach highlighted above is inadequate for the complete characterization of microbial communities.

### 1.1.2 Cultivation-Independent Molecular Techniques

Many of the cultivation-independent molecular methods for microbial diversity studies rely on the analysis of the 16S rRNA gene. This gene is part of the ribosomal operon and its size, nucleotide sequence, and secondary structure is conserved within bacterial species (Maidak *et al.*, 1997). It is composed of highly conserved sequence domains interspersed with hypervariable regions, which serve as primer docking sites and as phylogenetic markers, respectively. However, the presence of multiple rRNA operons in the bacterial genome of some species may lead to an overestimation of the microbial richness (Acinas *et al.*, 2004; Klappenbach *et al.*, 2000). Furthermore, the 16S rRNA gene does not reflect the remaining functional genome content (Robinson *et al.*, 2010). These restrictions have led to the consideration of alternative single-copy housekeeping genes to study the bacterial diversity of microbial ecosystems, such as the *rpoB* gene, encoding the  $\beta$ -subunit of the bacterial RNA polymerase (Case *et al.*, 2007). However, no gene is without limitations and the lack of universal primers and comprehensive databases for a wide spectrum of strains hampers their implementation in microbial community surveys. Similar to the bacterial 16S rRNA gene, the preferred molecular target for fungal phylogenetics is the ITS region of the nuclear DNA (Seifert, 2009).

Popular community profiling methods for food ecosystems are denaturing gradient gel electrophoresis (DGGE) and terminal restriction fragment length polymorphism (TRFLP) of amplified fragments. Although only sporadically used by food microbiologists, automated ribosomal intergenic spacer analysis (ARISA) can also be applied for community profiling. Those methods only provide a qualitative (*i.e.*, presence or absence) assessment of microbial communities. DGGE, TRFLP, and ARISA consist of the PCR amplification of a marker gene fragment of interest from the community DNA, followed by separation of the amplicons. Therefore, these methods are prone to amplification-related biases common to all PCR-based techniques. In the case of DGGE, the separation of PCR amplicons is based on a decreased electrophoretic mobility of PCR-amplified, partially melted, double-stranded DNA molecules in polyacrylamide gels containing a linear gradient of a DNA denaturant. Molecules with different sequences may have a different melting behavior and will stop migrating at different positions along the gel. The patterns generated could provide a preliminary ecological view of the predominant community members (Muyzer *et al.*, 1993). DGGE is applied by many researchers in the food microbiology field [as example (Gkatzionis *et al.*, 2014; Pangallo *et al.*, 2014; Papalexandratou *et al.*, 2011a; Papalexandratou & De Vuyst, 2011b; Papalexandratou *et al.*, 2011d)]. The drawbacks associated with this method are that it is time-consuming, has a low-throughput capacity, and is technically challenging, as the denaturing gradient gels are difficult to set and run properly (Papalexandratou *et al.*, 2011a). In addition, prokaryotic 16S rRNA genes may contain insufficient heterogeneity to reliably identify some bacterial species using the short amplicons necessary for DGGE separation. Another popular community profiling method to study microbial ecology

is TRFLP, which has received less attention by food microbiologists. This method differentiates microbial community members based on terminal restriction fragment (TRF) sizes of the amplified regions. Mixed DNA templates are amplified with fluorescence-labeled universal primers, digested with selected restriction enzymes, and separated by capillary electrophoresis according to fragment size. This method has been applied to study cheese and yogurt fermentation (Rademaker *et al.*, 2006), to analyze viable bacterial communities during cheese fermentations (Sanchez *et al.*, 2006), for species-level LAB differentiation (Bokulich & Mills, 2012c), to profile microbial successions during beer and wine fermentations (Bokulich *et al.*, 2012a), to analyze the bacterial composition of commercial probiotic strains (Marcobal *et al.*, 2008), and others. This method is HT and has a low technical demand, but it encounters the same problems as DGGE regarding the low taxonomic depth of the identifications obtained. Finally, ARISA is based on the use of a fluorescent primer for the amplification of microbial ribosomal intergenic spacers. The PCR products are analyzed in an automated capillary electrophoresis system that produces an electropherogram, the peaks of which correspond with discrete DNA fragments detected by a laser-based fluorescence detection system. The sensitivity of ARISA is very high and its reproducibility is reinforced by instrumental automation. This method is not widely used in food microbiology because the primer sets described are not optimized for food samples and the lack of large datasets comprising intergenic spacer sequences, limiting the identification of these fragments. It has been applied by Panelli and coworkers (2013) to diagnose the presence of *Clostridium tyrobutyricum* in raw milk before cheese making.

Although more expensive and time-consuming than the community fingerprinting techniques highlighted above, sequence analysis of clone libraries provides an unparalleled level of phylogenetic resolution due to the long read lengths generated by Sanger sequencing technology (Sanger *et al.*, 1992). This method involves PCR amplification of phylogenetic marker genes, followed by insertion of these PCR products into a plasmid vector, which is subsequently used for transformation of competent *Escherichia coli* cells. The DNA sequence of the insert is finally resolved using the Sanger sequencing technology. Clone libraries have been constructed in numerous studies on food fermentation, including the detection of cucumber fermentation spoilage bacteria (Breidt *et al.*, 2013), the assessment of the bacterial community structure in kimchi (a Korean fermented vegetable food) (Kim & Chun, 2005), sour cassava starch, cachaça, and minas cheese production (Lacerda *et al.*, 2011), Ghanaian and Brazilian cocoa bean fermentations (Garcia-Armisen *et al.*, 2010), and others. The drawbacks associated with clone libraries are that the method is time-consuming, has a low-throughput capacity, is labor-intensive, and is prone to PCR biases. Furthermore, it can only provide a qualitative assessment of the microbial communities.

The quantification deficit of the approaches discussed above can be counteracted by quantitative PCR (qPCR) analysis, a powerful extension of conventional PCR that allows reliable and accurate quantifications. qPCR has become an extremely

popular method in food microbial ecology studies, in which amplification is tracked in real-time using a fluorescent reporter molecule. A baseline threshold is determined at which sample fluorescence can be distinguished from background noise. This is used to determine the quantification cycle ( $C_q$ ) for each sample, at which fluorescence crosses the baseline threshold. A standard curve is constructed by plotting  $C_q$  against gene copy number or cell count and is used to quantify unknown samples based on  $C_q$ . Beyond cell quantification based on a taxonomic marker gene, the detection of functional genes provides interesting applications in food microbiology as well. This approach has been applied to detect biogenic amine-producing strains of LAB in wines (Arena *et al.*, 2011; Lucas *et al.*, 2008), ciders (Ladero *et al.*, 2011), and cheeses (Fernandez *et al.*, 2006; Torriani *et al.*, 2008), to quantify exopolysaccharide (EPS)-producing LAB (Ibarburu *et al.*, 2010), to track sulfide reductase genes in *Saccharomyces cerevisiae* that are involved in off-flavor production (Mendes-Ferreira *et al.*, 2010), and others.

Another frequently applied technique to detect and quantify bacteria in food samples is fluorescence *in situ* hybridization (FISH), which is used for directly visualizing microbial cells in a sample. This method utilizes fluorescence-labeled oligonucleotide probes targeting specific DNA sequences, unique for discrete microbial taxa. Cells are fixed and permeabilized *in situ*, incubated in the presence of probes to facilitate hybridization, and subsequently observed directly by fluorescence microscopy or counted via flow cytometry if suspended in a fluid. This technique has become very popular, as it avoids the challenges and biases of culturing and PCR assays and, additionally, it allows the observation of target cells within their native environment. The drawbacks of this method are its low-throughput capacity and its lower efficiency for enumerating cells compared to qPCR. FISH has been used to monitor microbial communities in a number of fermented foods, including yeasts and bacteria in wine (Andorra *et al.*, 2011; Blasco *et al.*, 2003; Xufre *et al.*, 2006), beer (Yasuhara *et al.*, 2001), and cheese (Babot *et al.*, 2011; Bottari *et al.*, 2010; Mounier *et al.*, 2009).

## 1.2 Microbial Diversity Analyses in the HT Sequencing Era

The traditional Sanger DNA-sequencing technology (Sanger *et al.*, 1992) can only sequence specimens individually and is therefore inadequate for processing complex microbial communities, especially in large-scale studies. Although conventional sequencing has provided large DNA reference databases, the number of individuals in a complex sample is beyond its ability. Recovering DNA sequences from complex and/or multiple samples requires the capacity to read DNA from multiple templates in parallel, which HT sequencing technologies do effectively at low costs.



### 1.2.1 Platforms

In 2005, the first HT sequencing technology emerged and drastically reduced the time and cost needed for sequencing (Metzker, 2005). The currently available HT sequencing technologies can be categorized into two broad groups, depending on the kind of template needed for the sequencing reactions (Loman *et al.*, 2012). The oldest group comprises the construction of a library of clonally amplified templates [e.g., the 454 (Roche Diagnostics), Ion Torrent (Life Technologies), Illumina (Illumina Inc.), SOLiD (Life Technologies), and Complete Genomics (Complete Genomics Inc.) platforms], whereas the more recently developed group of HT sequencing technologies involves the sequencing of single molecules [e.g., the HeliScope Single-Molecule (Helicos BioSciences), PacBio RS (Pacific Biosystems), and Oxford Nanopore (Oxford Nanopore Technologies) platforms] (Figure 2). Considerable variation in cost, time, throughput, read length, and error rate exists within these broad categories of HT sequencing technologies.

The first group of template amplification technologies involves a three-stage workflow of (i) library preparation, (ii) template amplification, and (iii) sequencing (Figure 2). Library preparation (i) for shotgun sequencing involves an initial (mechanical or enzymatic) fragmentation step to generate random, overlapping DNA fragments. Fragmentation is followed by the ligation of adapters to the fragments generated. Tagmentation is a promising transposase-based approach that, in a single step, fragments DNA and incorporates sequence tags. Currently, the only available implementation of tagmentation is within the Nextera® system, which is only available for the Illumina platform. Next to shotgun sequencing, all platforms can handle PCR products, allowing adapter sequences to be incorporated into the 5' ends of the primers. In preparation of amplification, template molecules are immobilized on a solid surface, which is a flow cell for sequencing with the Illumina platform or beads for other approaches (ii). Simultaneous solid-phase amplification of millions or billions of spatially separated fragments, prepares the way for massive parallel sequencing. For the Illumina platform, clusters are generated by bridge amplification on the surface of a flow cell. For platforms using bead-based immobilization (the SOLiD, 454, and Ion Torrent platforms), amplified template sequence libraries rely on an emulsion PCR, in which the beads are enclosed in aqueous-phase microreactors and are kept separately from each other into a water-oil emulsion.




Figure 2: HT sequencing platforms

The main HT sequencing platforms available and their associated sample preparation and template amplification procedures are shown. PGM: personal genome machine. Adapted from (Loman *et al.*, 2012). Adaptations to the original figure are highlighted with black squares.

The last part of the workflow in template amplification technologies is the sequencing (iii). Illumina, 454, and Ion Torrent technologies rely on the sequencing-by-synthesis design, whereas the SOLiD platform and the platform from Complete Genomics depend on sequencing-by-ligation. Illumina uses the Solexa chemistry that includes reversible termination of sequencing products, whereas in 454 and Ion Torrent a dNTP flows across the template in each cycle. More specifically, 454 uses the pyrosequencing approach, whereby the presence of pyrophosphate is signaled by visible light as the result of an enzyme cascade. In contrast, Ion Torrent does not rely on light emission, but uses a modified silicon chip to detect hydrogen ions that are released during base incorporation. The SOLiD and Complete Genomics platforms use an approach in which fluorescent probes undergo iterative steps of hybridization and ligation to complementary positions in the template strand followed by fluorescence imaging to identify the ligated probe.

The second group of single-molecule sequencing platforms does not comprise amplification and library preparation steps, thereby circumventing the artifact associated with these steps. The HeliScope Single-Molecule sequencer was the first entering the market in 2009. This technology applies one-color reversible terminator sequencing to unamplified single-molecule templates. Another more recent technology is the one of PacBio, in which dye-labeled nucleotides are continuously incorporated into a growing DNA strand by a specialized strand-displacing DNA polymerase that is attached to a solid surface. This technology allows continuous imaging of the labeled nucleotides as they enter the strand. Another promising single-molecule sequencing technology is the one of Oxford Nanopore, which is currently developing the 'strand sequencing' technology that exploits protein nanopores embedded in an industrially fabricated polymer membrane. This technology passes intact DNA polymers through a protein nanopore, allowing sequencing in real-time as the DNA translocates the pore.

Most companies have both high-end and bench-top instruments to comply with the needs of different users, which are large sequencing centers and research laboratories, respectively. The high-end machines include PacBio RS, the HiSeq instruments, Genome Analyser IIX, the SOLiD 5500 series, and the 454 GS FLX<sup>+</sup> system, whereas the bench-top instruments include the GS Junior, Ion PGM, and the Illumina MiSeq. An overview of these instruments and their performance is summarized in Table 1.

### 1.2.2 (Meta)genomics

The advent of HT sequencing technologies and the resulting sequencing at reduced cost per base have changed microbiological research drastically in the last decade, making genomics and metagenomics affordable for many research laboratories.

Table 1: Comparison of HT sequencing platforms

Machine (manufacturer)	Chemistry	Modal read length (bases)	Run time	Output per run	Advantages	Disadvantages
<b>High-end instruments</b> 454 GS FLX+ (Roche)  HiSeq2500 (Illumina)  NextSeq500 (Illumina)  PacBio RS (Pacific Biosystems) MinION, PromethION, GridION (Oxford Nanopore's)	Pyrosequencing	700 bp/450 bp	23 hours/10 hours	700 Mb/450 Mb	Long read lengths	Appreciable hands-on time, high reagent costs, high error rate in homopolymers
	Reversible terminator	2x250/2x125 bp	7-60 hours/<1 day - 6 days	10-300 Gb/50-1000 Gb	Cost-effectiveness, steadily improving read lengths, massive throughput, minimal hands-on time	Long run time, short read lengths
	Reversible terminator	2x150 bp	15-26 hours/12-30 hours	20-39 Gb/30-120 Gb	Cost-effectiveness, steadily improving read lengths, massive throughput, minimal hands-on time	Long run time, short read lengths
	Real-time sequencing	>14000 bp	30 -240 minutes	500 Mb-1 Gb	Simple sample preparation, low reagent costs, very long read lengths	high error rate, expensive system, difficult installation
	Real-time sequencing	not fixed	not fixed	not fixed	Very long read lengths	Instruments not available at the time of writing
<b>Bench-top instruments</b> 454 GS Junior + (Roche)  MiSeq (Illumina)  Ion Personal Genome Machine (Life Technologies)  Ion Proton (Life Technologies)	Pyrosequencing	700 bp	18 hours	70 Mb	Long read lengths	Appreciable hands-on time, high reagent costs, high error rate in homopolymers
	Reversible terminator	2x300 bp	5-55 hours	0.3-15 Gb	Cost-effectiveness, short run times, appropriate for microbial applications, minimal hands-on time	Read lengths too short for efficient assembly
	Proton detection	200 bp	Ion 314: 2.3-3.7 hours/Ion 318: 4.4 - 7.3 hours/Ion 316: 3.0 - 4.9 hours	2 Gb	Short run times, appropriate throughput for microbial applications	Appreciable hands-on time, high error rate in homopolymers
	Proton detection	200 bp	2-4 hours	10 Gb	Short run times, flexible chip reagents	Appreciable hands-on time, high error rate in homopolymers

Adapted from (Loman *et al.*, 2012)

Genomics refers to the study of the genomes of organisms. Species with defined genomes increase the value of many different biological studies by enabling the comparison of microbial evolution, metabolism, and ecology of organisms in the environment (DeLong & Karl, 2005). The first sequenced bacterial genome was that of *Haemophilus influenzae* in 1995 (Fleischmann *et al.*, 1995). Since then, the genomes online database (GOLD) (Bernal *et al.*, 2001) lists thousands of bacterial genomes, including both finished and permanent draft genomes. These bacterial genomes vary in terms of G+C content, gene content and density, genome size and number of plasmids. The accumulation of genome sequences is crucial for comparative genome analysis, for at least two related but distinct fundamental reasons: (i) identifying sets of orthologs [*i.e.*, genes that arise by speciation and tend to retain similar functions (Richardson & Watson, 2012)], and (ii) searching for genes that are absent in one but present in another genome sequence. The ability to delineate sets of orthologs and to search for missing genes is indispensable for genome-based reconstruction of an organism's metabolism and other functional systems and for reconstruction of genome evolution.

Bacterial genome sequence analysis involves assembly, gene prediction, and annotation. Genome assembly involves the process of reconstructing a genome sequence by joining short reads together up to the chromosomal level. Assembly procedures can be classified as either reference-guided genome assemblies or *de novo* genome assemblies. For the sake of computational memory saving and convenience of data inquiry, HT short reads data are always initially formatted to a specific data structure. Currently, the majority of existing data structures are graph-based. A well-known graph-based assembly tool is the Velvet *de novo* assembler, which builds a *de Bruijn* graph from short reads and removes errors from the graph (Zerbino & Birney, 2008). Following, gene prediction is the identification of open reading frames encrypted in the DNA sequence. Examples of gene finding tools are Glimmer (Delcher *et al.*, 2007) and GeneMark (Lukashin & Borodovsky, 1998), which are implemented in the rapid annotation using subsystems technology (RAST) (Aziz *et al.*, 2008) and the prokaryotic genome automatic annotation pipeline (PGAAP) (Angiuoli *et al.*, 2008), respectively. The next step is to take the set of predictions and search for hits against one or more protein and/or protein domain databases (*i.e.*, annotation) using BLAST (*i.e.*, basic local alignment tool) (Altschul *et al.*, 1997), HMMer (*i.e.*, hidden markov model) (Eddy, 1998) or other programs. For each gene with a significant match, the BLAST output together with the annotation of the hit can be used to assign a name and function to the corresponding protein. The accuracy of this step depends not only on the annotation software but also on the quality of the annotations already in the reference database. Many genomes of bacteria involved in food fermentations, including several LAB, have already been sequenced [as example (Illegghems *et al.*, 2013; Moreno Switt *et al.*, 2014; Makarova *et al.*, 2006)].

Contrasting to genomics, metagenomics refers to culture-independent studies of the collective set of genomes of mixed microbial populations and applies to explorations of all microbial genomes that reside in a certain niche of interest (Petrosino *et al.*, 2009). It starts with the extraction of genomic DNA from cellular organisms and/or viruses in a sample. This metagenomic DNA can either be sheared into fragments and sequenced with the platform of choice (shotgun sequencing strategy), or one can enrich and sequence a target region of interest (target enrichment strategy) (Figure 1). Metagenomics can be applied to study both qualitative and quantitative aspects of the microbial community of interest. For the qualitative approach, the recognition of the presence of a taxonomic unit or a metabolic function is used to describe the microbial diversity present in the ecosystem. The quantitative approach provides the relative proportion of the taxonomic unit or metabolic function relative to others in the same sample or to other samples or ecosystems. Nevertheless, caution must be taken when interpreting these data because sequencing depth, DNA extraction procedures, PCR biases, and others bias the outcome.

In the field of metagenomics, the shotgun sequencing strategy generates a huge amount of short reads, comprising the metagenome of the community studied. Similarly to genomics, data processing usually involves the assembly of short overlapping sequence reads into a consensus sequence, followed by the prediction and annotation of coding sequences (CDSs). These data enable researchers to look for functionalities of the communities and to identify microbial community members in a process called binning or classification. The metagenomic RAST (MG-RAST) server is an automated analysis platform providing phylogenetic and functional analyses of metagenomes (Meyer *et al.*, 2008). Examples of alternative tools are metaGenome analyzer (MEGAN4) (Huson & Mitra, 2012), integrated microbial genomes/metagenomes (IMG/M) (Markowitz *et al.*, 2014), community cyberinfrastructure for advanced marine microbial ecology research and analysis (CAMERA) (Seshadri *et al.*, 2007), metagenomics Reports (METAREP) (Goll *et al.*, 2010), and others. Food fermentations that have been analyzed using the shotgun sequencing strategy are a kimchi (Jung *et al.*, 2011) and cocoa bean fermentation processes (Illegghems *et al.*, 2012). Despite the increased throughput and resulting reduced prices of HT sequencing technologies, metagenome shotgun sequencing is still expensive and time-consuming, especially for complex metagenomes containing a large number of eukaryotic genomes and for the analysis of a large number of samples. Furthermore, the storage of these large datasets will place a substantial burden on a research center's bioinformatics infrastructure.

Consequently, considerable efforts have been devoted to develop target enrichment methods, in which genomic regions are selectively enriched from a community DNA sample before sequencing (Mamanova *et al.*, 2010). This approach is, compared to the metagenome shotgun sequencing strategy, more cost-effective and the resulting data are considerably less cumbersome to store and to analyze. Its major application is to identify microorganisms in complex communities by exploiting targets

containing both universal and conserved sequences, such as regions within 16S rRNA genes, which enable approximate identification of bacteria and archaea. The same principle holds for the identification of fungal communities by amplifying a fragment of the ITS region. Several tools for target enrichment have been developed, each with their own advantages and disadvantages (Mamanova *et al.*, 2010). For instance, PCR is compatible with any HT sequencing platform, though to make full use of the HT, a large number of amplicons must be sequenced together. However, PCR is difficult to perform in multiplex to any useful degree: the simultaneous use of many primer pairs can generate a high level of nonspecific amplifications, caused by interactions between the primer pairs, and furthermore, amplifications can fail (Cho *et al.*, 1999; Wang *et al.*, 1998). Clever derivatives of multiplex PCR have been developed (Meuzelaar *et al.*, 2007; Varley & Mitra, 2008; Simon *et al.*, 2007), but in practice, it is often more straightforward to perform PCRs in uniplex (Figure 3). Following uniplex amplification, the concentration of PCR products must be normalized before pooling. A convenient solution to many of the problems encountered in a standard PCR-based approach is the RainStorm platform, developed by RainDance Technologies (<http://raindancetech.com/>), which uses microdroplets, similar to emulsion PCR. Each droplet supports an independent PCR and contains a single primer pair along with genomic DNA and other reagents. The entire population of droplets represents many distinct primer pairs and is subjected to thermal cycling, after which this emulsion is broken and PCR products are recovered. The advantages of this technology are that different primer pairs cannot interact with each other, and that direct competition of multiplex PCRs for the same reagent pool is prevented, which should improve uniformity relative to conventional multiplex PCR. Even with an efficient, automatic PCR pipeline, it is not feasible to use conventional PCR to target genomic regions that are several megabases in size because of the high cost of primers and reagents and the DNA input requirements, particularly in large sample sets. Consequently, for very large or moderately sized target regions, other approaches to target enrichment should be used. Alternative strategies involve the usage of molecular inversion probes (Hardenbol *et al.*, 2003) and hybrid capture (Albert *et al.*, 2007).




Figure 3: PCR as a target enrichment strategy

(Mamanova *et al.*, 2010)

The use of target enrichment strategies is common in food microbiology. Pyrosequencing of PCR-amplified 16S rRNA genes has been applied to study the bacterial diversity of a number of fermented foods, including narezushi (fermented fish and rice) (Kiyohara *et al.*, 2012; Koyanagi *et al.*, 2011), nukadoko (fermented rice bran) (Sakamoto *et al.*, 2011), Chinese liquor fermentations (Li *et al.*, 2011), Danish raw-milk cheeses (Alegria *et al.*, 2012), Polish oscypek cheese (Alegria *et al.*, 2012), and pearl millet fermentations (Humblot & Guyot, 2009). Additionally, it has been used to analyze the bacterial and archaeal or fungal communities of fermented seafood (Roh *et al.*, 2010), makgeolli (rice beer) (Jung *et al.*, 2012), meju (Kim *et al.*, 2011), doenjang (soybean pastes) (Nam *et al.*, 2012a), kochujang (fermented red pepper condiment) (Nam *et al.*, 2012b), and kimchi (Park *et al.*, 2012). Because of the increasing read length, higher throughput and lower cost of Illumina sequencing, its application recently gained popularity in food microbiology. For instance, Bokulich and coworkers used Illumina sequencing to study botyriized wine fermentations (2012b) and the brewhouses resident microbiota responsible for the fermentation of American coolship ales (ACA) (2012a), beers that are characterized by a spontaneous fermentation process. Compared to Sanger sequencing of PCR clone libraries, these target enrichment strategies permit a much deeper sampling of microbial communities by providing orders of magnitude more sequence information. Moreover, performing PCR in multiplex enables different samples to be analyzed in parallel and makes the inclusion of technical, biological, and experimental replications feasible. These advantages will provide greater sensitivity than PCR clone libraries (Sogin *et al.*, 2006; Parameswaran *et al.*, 2007).



# 2

## Importance of LAB in Food Fermentations

LAB play a prominent role in the world food supply, performing the main bio-conversions in fermented dairy products, meats, cereals, and vegetables. They contribute to the taste and texture of fermented food products and inhibit food spoilage bacteria by producing growth-inhibiting substances and large amounts of lactic acid, thereby prolonging the shelf-life of fermented foods.

One of the key metabolic features of LAB is carbohydrate fermentation coupled to substrate-level phosphorylation, finally resulting in the reduction of pyruvate into lactate to regenerate  $\text{NAD}^+$  in a process catalyzed by lactate dehydrogenase (Figure 4). Based on differences in carbohydrate fermentation patterns, they are subdivided into homofermentative and heterofermentative strains (Stiles & Holzapfel, 1997). The homofermenters produce lactic acid as the major end-product of carbohydrate fermentation, whereas heterofermenters generate ethanol and  $\text{CO}_2$  next to lactic acid. Differences between both categories occur on the enzyme level, with aldolase being absent and phosphoketolase being present in heterofermenters, and vice versa in homofermenters. Aldolase is a key enzyme of the glycolysis, converting fructose 1,6-biphosphate into glyceraldehyde 3-phosphate and dihydroxy acetone phosphate, whereas phosphoketolase converts xylulose 5-phosphate into acetyl phosphate and glyceraldehyde 3-phosphate. The homofermentative metabolism is energetically favored to the heterofermentative one, although the latter has the competitive advantage of catabolizing pentoses next to hexoses,

whereas homofermenters can only utilize hexoses. Next to acidification due to the carbohydrate metabolism, LAB contribute to flavor due to their amino acid catabolism (Figure 4). In general, aminotransferases are involved in the conversion of amino acids (branched-chain amino acids, aromatic amino acids, and sulfur-containing amino acids) into the corresponding  $\alpha$ -keto acids, followed by decarboxylation into the corresponding aldehydes and further dehydrogenation into the corresponding alcohols.




Figure 4: Major pathways active in LAB

Left panel: carbohydrate metabolism via the glycolysis (homofermentative LAB; lower left) and the phosphoketolase pathway (heterofermentative LAB; lower middle). Upper right panel: generic conversion of branched-chain amino acids (valine, leucine, and isoleucine), aromatic amino acids (tyrosine, tryptophan, and phenylalanine), and sulphur-containing amino acids (methionine), as initiated by transamination in LAB. Lower right panel: arginine deiminase pathway (ADI). Key enzymes involved in these pathways are designated with Roman numbers: I, cell wall-bound fructosyltransferase; II,  $\beta$ -galactosidase; III, maltose phosphorylase; IV, phosphoketolase; V, lactate dehydrogenase; VI, acetate kinase; VII, aminotransferase; VIII, glutamate dehydrogenase; IX, decarboxylase; X, arginine deiminase; XI, ornithine transcarbamoylase; XII, carbamate kinase. (Ravyts *et al.*, 2012)

Current taxonomy classifies LAB genera in the order *Lactobacillales* within six families: *Aerococcaceae*, *Carnobacteriaceae*, *Enterococcaceae*, *Lactobacillaceae*, *Leuconostocaceae*, and *Streptococcaceae* (Figure 5). Food-grade LAB are generally restricted to the genera *Lactobacillus*, *Enterococcus*, *Leuconostoc*, *Fructobacillus*, *Weissella*, *Pediococcus*, *Lactococcus*, *Streptococcus*, *Carnobacterium*, *Oenococcus*,

and *Tetragenococcus*. They possess small genomes (approximately 2 Mb), encoding a range of biosynthetic capabilities that reflect both prototrophic and auxotrophic characters (Makarova *et al.*, 2006). Furthermore, their genomes encode a broad repertoire of transporters for efficient carbon and nitrogen acquisition from the nutritionally rich environments they inhabit (Makarova *et al.*, 2006). This suggests both extensive gene loss as well as acquisitions via horizontal gene transfer during the evolution of LAB within their habitats (Makarova *et al.*, 2006). Several strains have plasmids containing genes regulating the fermentation of carbohydrates and encoding resistances. Most LAB are free-living or live in beneficial or harmless associations with animals, although some are opportunistic pathogens.




Figure 5: Consensus dendrogram reflecting the phylogenetic relationships of a selection of LAB genera of the order Lactobacillales within the class Bacilli

(Garrity *et al.*, 2004). The genus *Leuconostoc* includes *Leuconostoc sensu stricto* and the genus *Fructobacillus* (Endo & Okada, 2008).

Examples of fermented foods, for which LAB are critical in their production are yogurt, cheese, butter, sour cream, sausage, pickles, olives, sauerkraut, cocoa, and others. For instance, LAB play a major role in cocoa bean fermentation, which is characterized by the successional growth of various species of yeasts, LAB, and AAB whereby LAB contribute to the fermentation of carbohydrates and citrate into lactate, acetate, and mannitol (De Vuyst *et al.*, 2010). A diversity of species within the genera *Lactobacillus*, *Streptococcus*, *Lactococcus*, *Oenococcus*, *Pediococcus*, *Weissella*, *Fructobacillus*, and *Leuconostoc* have been isolated from fermented cocoa beans from different locations (Papalexandratou *et al.*, 2011a). LAB also occur in the beer environment, with *Lactobacillus* and *Pediococcus* species being frequently reported as beer spoilers (Bokulich & Bamforth, 2013; Bergsveinson *et al.*, 2012; Pittet *et al.*, 2012). LAB are also known to contribute to the flavor development of certain spontaneously fermented beers, such as lambic beers (Spitaels *et al.*, 2014b) and beers produced by mixed-culture fermentation, such as Belgian red-brown acidic ales, including red acidic ales and red-brown acidic ales (Martens *et al.*, 1997). *Pediococcus damnosus* is a predominant microbial community member during the maturation of these beers, resulting in increased lactic acid concentrations and thereby contributing to the fresh and acidic taste of the bottled beers. Additionally, psychrotrophic LAB are increasingly reported as spoilers within the storage period of cold-stored packaged foods, causing alterations of the organoleptic properties of these food products (Pothakos *et al.*, 2014). Examples of LAB species that successfully proliferate under refrigeration temperatures are *Lactobacillus sakei*, *Lactobacillus curvatus*, *Leuconostoc mesenteroides*, *Leuconostoc gelidum* subsp. *gasicomitatum*, *Leuconostoc gelidum* subsp. *gelidum*, and *Lactococcus piscium* (Pothakos *et al.*, 2014).

An interesting application of LAB is their use as starter cultures, which are defined as microbial preparations of large numbers of cells of at least one microorganism that is added to a raw material to produce a fermented food by accelerating and steering its fermentation process (Leroy & De Vuyst, 2004). The advantages of their use are that a high degree of control over the fermentation process is achieved and standardization of the end-products is reachable. Additionally, LAB can be used as functional starter cultures, which are starters that possess at least one inherent functional property besides lactic acid production. Functional starter cultures can contribute to food safety and/or offer organoleptic, technological, nutritional, or health advantages (Leroy & De Vuyst, 2004; Leroy *et al.*, 2006). Examples are LAB strains that are able to produce antimicrobial substances, carbohydrates, polymers, sweeteners, aromatic compounds, useful enzymes, or nutraceuticals, or LAB with health-promoting properties, so called probiotic strains. Furthermore, progress has been made in the construction of food-grade genetically modified LAB for the improvement of fermented foods (Smid *et al.*, 2006), and for usage as mucosal delivery vectors for therapeutic proteins and DNA vaccines (Bermúdez-Humarán *et al.*, 2013; Sybesma *et al.*, 2006).

# 3

## LAB Taxonomy

The availability of an appropriate taxonomic framework in sequence databases is crucial for organizing and cataloging microbial diversity (Yilmaz *et al.*, 2013). Taxonomy relies on three key elements depending on each other: classification (*i.e.*, the orderly arrangement of organisms into taxonomic groups on the basis of similarity), nomenclature (*i.e.*, the labeling of taxonomic groups), and identification of unknowns (*i.e.*, the process of determining whether an organism belongs to one of the taxonomic groups or represents a novel taxon). While nomenclature is governed by the International Code of Nomenclature of Bacteria (1990 Revision) (Lapage *et al.*, 1992), the classification and identification of prokaryotes has been a changing area in the last 50 years. Phylogeny elucidates evolutionary relationships among organisms and is a tool for classification. The 16S rRNA gene is one of the best targets for phylogenetic studies of bacteria and its sequences therefore provide a backbone in taxonomy, as they yield a bacterial phylogenetic framework (Vandamme *et al.*, 1996). The bacterial species is an unit in biological classification, which is crucial for the construction of a useful taxonomic framework based on microbial evolution (Gevers *et al.*, 2005; Dijkshoorn *et al.*, 2000). Various species concepts, describing bacterial species and explaining how they are formed, have been suggested for microbial species (see below), but none have been generally accepted (Rossello-Mora & Amann, 2001). The definition of a bacterial species is how the concept is applied in practice and consists of a set of rules (see box below) (Konstantinidis *et al.*, 2006). While the definition of an eukaryotic species has focused on the biology of reproduction, the bacterial species has mainly been discussed in terms of similarity between strains.

**Species concepts in bacteriology (Achtman & Wagner, 2008)**

'A species could be described as a monophyletic and genomically coherent cluster of individual organisms that show a high degree of overall similarity in many independent characteristics, and is diagnosable by a discriminative phenotypic property (Rossello-Mora & Amann, 2001).'

'Species are considered to be an irreducible cluster of organisms diagnosably different from other clusters and within which there is a parental pattern of ancestry and descent (Staley, 2006).'

'A species is a group of individuals where the observed lateral gene transfer within the group is much greater than the transfer between groups (Dykhuizen, 2005).'

'Microbes do not form natural clusters to which the term 'species' can be universally and sensibly applied (Nesbo *et al.*, 2006).'

'Species are (segments of) metapopulation lineages (de Queiroz, 2005).'

### 3.1 Phylogenetics

The term 'phylogenetics' refers to the study of the evolutionary history of species, organisms, genes, or proteins through the construction and analysis of mathematical entities known as trees or phylogenies (Sleator, 2011). It involves the alignment of nucleic acid or protein sequences between extant organisms. An inference is then generated to explain the repartition of character states and the results are presented as a phylogenetic tree, which is a graphic representation of the computed results (Sleator, 2011; Hall, 2011).

While several methods exist to infer evolutionary relatedness, most can be classified as either distance- or character-based methods (Figure 6). Distance-based methods utilize an algorithm, incorporating a model of evolution, to compute a distance-matrix from which a phylogenetic tree is calculated by means of progressive clustering (Sleator, 2011). Distances in the matrix relate to the number of differences between each pair of DNA or protein sequences. The tree is constructed from the numerical data in the matrix, with the most closely related sequences occupying a position on the tree which is distant from the less closely related sequences. Both the neighbour-joining (NJ) and the unweighted pair group method using arithmetic averages (UPGMA) approaches employ distance-based methods (Figure 6). Although effective, distance-based methods have a number of disadvantages. For instance, NJ may compute different trees depending on the order in which the constituent sequences are added (Sleator, 2011). Furthermore, NJ provides only a single tree rather than a consensus tree (Sleator, 2011). Character-based methods, such as maximum parsimony (MP) and maximum likelihood (ML) methods, search for the most probable tree for a specific sequence set, based on characters at each

position of the sequence alignment (Sleator, 2011). Each character is considered one at a time to calculate the 'tree score', representing the minimum number of changes for maximum parsimony and the log-likelihood value for maximum likelihood. *MP* seeks to find the tree(s) that are compatible with the minimum number of substitutions among sequences, *i.e.*, the fewest evolutionary changes. An advantage of *MP* is that it provides diagnosable units for each clade and branch lengths in terms of the number of changes on each branch. However, a significant limitation of the *MP* approach is that it requires strict assumptions of consistency across sites and among lineages. *ML* methods are based on specific probabilistic models of evolution and search for the tree with maximum likelihood under these models. The major advantage of likelihood approaches is that they are based on powerful statistical theory. However, *ML* approaches are computationally intensive, limiting its use to a relatively small number of sequences. The distance-based and *ML* methods make use of a model of evolution and as such are model based, while *MP* does not have an explicit model.




Figure 6: Schematic overview of the major analytical approaches to phylogenetic tree building.

(Sleator, 2011). The Bayesian inference approach is not discussed in the text.

In order to reconstruct evolutionary trees, assumptions about the substitution process must be made by means of a model of evolution. A model of evolution makes assumptions on how DNA or amino acid substitutions occurred in the DNA or protein sequence since they last shared a common ancestor (Sleator, 2011; Hall, 2011). Because multiple substitutions can occur at the same site, there is a potential for underestimating the amount of change that has occurred along a branch. Each model of evolution represents an attempt to account for multiple hits and various other aspects of the substitution process. The easiest model to consider is one in which the probabilities of any nucleotide changing to any other nucleotide

are equal (*i.e.*, One-Parameter or Jukes-Cantor model. The One-Parameter Model is the simplest, but it is not very realistic because all changes do not occur at the same rates. Therefore, a variety of models have been proposed that allow to specify different rates. The most general model is one in which each different substitution can occur at a different rate, involving 12 free-parameters. A lot of important models are special cases of this general model. For instance, the Kimura 2-parameter model (Kimura, 1980) extends the Jukes-Cantor correction by taking into account the possibility that the rates at which transitions and transversions occur might be different. The Tamura 3-parameter model adds a correction for compositional bias (Tamura, 1992). The general time-reversible (GTR) model has six different substitution probabilities. It involves nine free parameters and the equilibrium base frequencies. The substitution process looks the same whether we observe the process with time running forward or backward (Hall, 2011).

The models listed above implicitly assume that the rates are the same at all sites, but it is also possible to include rate variation across sites in the models. An example of site-specific variation is that second positions in codons evolve most slowly, first positions evolve at an intermediate rate, and third positions evolve most rapidly. Furthermore, because proteins fold, those slowly evolving sites tend to be located in patches rather than in one specific region of the sequence. Estimating these individual rates for each site is often computationally impractical when many sequences are involved. Therefore, the alternative is to assume a well-behaved distribution of rates across sites. The distribution that is commonly used is the gamma distribution. Furthermore, some sites, such as initiation codons, may not be free to vary at all. This constitutes a specific case of site-specific variation, which we call the invariant sites (Hall, 2011).

Once an optimum tree is chosen, some statistical measure of internal support for clades must be obtained to ascertain whether the tree is sufficiently robust and biologically meaningful (Sleator, 2011). A method that is frequently used to verify the evolutionary reliability of trees is the bootstrap analysis that pseudoreplicates the collection of data as a method to estimate the reliability of the tree. A random site is taken from the alignment and is used as the first site in a pseudoalignment. Another random site is taken and used as the second site in the pseudoalignment, and this process is continued until the pseudoalignment contains the same number of sites as the original alignment. The tree is then constructed from the pseudoalignment using the same method and under the same parameter settings used to estimate the original tree (Hall, 2011).



## 3.2 Polyphasic Taxonomy

Developments in the field of sequencing of rRNA genes contributed to bacterial phylogeny and molecular fingerprinting techniques and resulted in the introduction of the polyphasic taxonomy concept (Vandamme *et al.*, 1996). Polyphasic taxonomy comprises the determination of the phylogenetic position of an isolate, based on measures of evolutionary relationships, which are in turn based on gene sequences (e.g., the 16S rRNA gene) and phenotypic properties to assess novelty. Within this framework of polyphasic taxonomy, strains belonging to the same species should form differentiable coherent groups, displaying similar phenotypes, genotypes, and chemotaxonomic features. The polyphasic taxonomy scheme defines a bacterial species as a group of strains having a certain degree of phenotypic consistency, > 70% DNA-DNA hybridization (DDH) similarity, and > 97% 16S rRNA gene sequence identity (Vandamme *et al.*, 1996).

In the case of LAB species, unraveling phylogenetic relationships is cumbersome because of the low taxonomic resolution of the 16S rRNA gene, resulting in only an approximate or tentative identification if high (> 97%) similarity values are found. Therefore, MLSA of different housekeeping genes is often needed to obtain a species level identification. The first comprehensive effort to make a large database for the classification and identification of LAB has been made by Naser and coworkers (2005a; 2005b; 2007). They evaluated the MLSA of three housekeeping genes [*i.e.*, *pheS* encoding phenylalanine transfer RNA (tRNA) synthetase, *rpoA* encoding RNA polymerase  $\alpha$ -chain, and *atpA* encoding RNA polymerase  $\alpha$ -subunit] as species identification tool within the genera *Enterococcus* and *Streptococcus*, and two of these genes (*i.e.*, *pheS* and *rpoA*) within the genus *Lactobacillus* (Naser *et al.*, 2005a). This MLSA scheme was expanded by De Bruyne and coworkers (2007), who constructed a MLSA scheme based on *pheS*, *rpoA*, and *atpA* for the genera *Pediococcus*, *Leuconostoc*, and *Fructobacillus*. Furthermore, a single-locus sequence analysis framework was established based on *pheS* for the genera *Weissella* and *Oenococcus* (De Bruyne *et al.*, 2008a). Furthermore, Rademaker and coworkers (2007) extended this list with MLSA for the genus *Lactococcus* based on the *pheS*, *rpoA*, *atpA*, *bcaT* (encoding branched chain amino acid aminotransferase), *pepN* (encoding aminopeptidase N), and *pepX* (encoding X-prolyl dipeptidyl peptidase) gene sequences.

If a LAB strain or a set of strains belong to a novel taxon, they should be characterized as comprehensive as possible (Tindall *et al.*, 2010). The goal of this characterization is to place them within the hierarchical framework laid down by the International Code of Nomenclature of Bacteria (Lapage *et al.*, 1992), as well as to provide a description of the novel taxon. The starting point for novel LAB species description is often the availability of a full-length 16S rRNA gene sequence with less than 97% sequence similarity towards its closest neighbor

(Stackebrandt & Goebel, 1994). When the 16S rRNA gene similarity exceeds 97%, MLSA and/or DDH are needed to determine whether the strain represents a novel taxon (Mattarelli *et al.*, 2014). For LAB species, the phenotypic description should also include the composition of the peptidoglycan of the cell wall and the ratio of D- to L-lactic acid production, as these are species-specific (Mattarelli *et al.*, 2014).

### 3.3 Genomic Taxonomy

Despite the widely used phylogenetic analysis based on the 16S rRNA gene, there has been considerable debate whether a tree based on any single gene can accurately represent the evolution of a species. Hence, the importance of the frequency of recombination, either in the form of homologous recombination (Fraser *et al.*, 2007) or lateral gene transfer (Doolittle & Papke, 2006), has challenged bacterial species concepts and raised many questions. Still, it has been shown that these forms of recombination are considered cohesive rather than disruptive forces in bacterial species (Konstantinidis & Tiedje, 2005). Furthermore, DDH experiments have several disadvantages, including the requirement of large quantities of high-quality DNA, its time-consuming and labor-intensive nature, and the inability to build databases (Gevers *et al.*, 2005). Because of these drawbacks, bacterial taxonomists are actively searching for alternative methods that can abandon DDH experiments (Gevers *et al.*, 2005; Coenye *et al.*, 2005; Cho & Tiedje, 2001).

DDH experiments were originally developed because the *ad hoc* committee on reconciliation of approaches to bacterial systematics (Wayne *et al.*, 1987) stated that taxonomy should be determined phylogenetically and that the complete genome sequence should therefore be the standard for species delineation. Particularly interesting within this regard is development of HT sequencing technologies, which caused the accumulation of whole-genome sequence data and allowed the establishment of taxonomic schemes based on evolutionary information contained in whole-genome sequences (Eisen, 2000; Wolf *et al.*, 2001).

Genomic taxonomy integrates comparative genomics in taxonomy and can include MLSA, phylogenetic analysis based on whole-genome sequences, average nucleotide identity (ANI) (Konstantinidis & Tiedje, 2005; Richter & Rossello-Mora, 2009), average amino acid identity (AAI), the percentage of conserved DNA (pcDNA) (Goris *et al.*, 2007), maximal unique matches (MUM) indices (Deloger *et al.*, 2009), core and pan genome analysis, genomic signatures, codon usage bias, metabolic pathway content, and others (Konstantinidis & Stackebrandt, 2013). Phylogenetic analysis based on whole-genome sequences are based on all genes of the core genome. The ANI and AAI are calculated among the nucleotide or amino acid sequences, respectively, of the conserved genes or DNA of a pair of genomes

(Konstantinidis & Tiedje, 2005; Goris *et al.*, 2007). Furthermore, the calculation of the pcDNA or MUM indices can be used as *in silico* imitations of DDH experiments, whereas the analysis of the ANI and the core genome are both natural expansions of MLSA. ANI represents a robust measure of the genetic and evolutionary distances between two sequenced strains, because it shows a strong correlation with 16S rRNA gene sequence similarity and the mutation rate of the genome, it is not affected by lateral gene transfer or variable recombination rates of single (or a few) genes, and it offers resolution at the subspecies level (Konstantinidis & Tiedje, 2005). The recommended cut-off of 70% DDH for species delineation corresponds to 95% ANI (Goris *et al.*, 2007). Finally, genomic signatures are compositional parameters, reflecting the di-, tri-, or tetranucleotide relative abundances (Karlin signature), which are similar between closely related species and dissimilar between non-related species.


The continuing accumulation of LAB genome sequences provides a good opportunity to study the evolutionary history of LAB species. The first LAB genome sequence was published in 2001, *i.e.*, that of *Lactococcus lactis* subsp. *lactis* IL 1403, a strain commonly used as cheese starter (Bolotin *et al.*, 2001). Nowadays, numerous LAB genome sequences are available, mainly belonging to the genera *Lactobacillus* and *Streptococcus*. Phylogenomic studies on LAB provide interesting new information but thus far include a rather limited number of species (Coenye & Vandamme, 2003; Makarova *et al.*, 2006; Zhang *et al.*, 2011; Claesson *et al.*, 2008).

Coenye and Vandamme (2003) analyzed 10 LAB genome sequences belonging to the genera *Streptococcus* (8 genomes), *Lactococcus* (1 genome), and *Lactobacillus* (1 genome). They showed that trees based on different kinds of phylogenetic information extracted from these genomes do not provide much additional information about the phylogenetic relationships among LAB taxa compared to more traditional alignment-based methods. Nevertheless, they expected that the study of the genome sequences would have their value in taxonomy by determining which genes are shared and when genes or sets of genes have been lost during evolution. Furthermore, these genome sequences could aid in detecting the presence of horizontally transferred genes and/or confirming or enhancing the phylogenetic signal derived from traditional methods. Next, Makarova and colleagues (2006) compared 12 complete genome sequences, covering the major families within the order *Lactobacillales* [*Leuconostocaceae* (2 genomes), *Lactobacillaceae* (7 genomes), and *Streptococcaceae* (3 genomes)]. Phylogenetic analysis of multiple protein CDSs showed that the streptococci-lactococci branch is basal in the *Lactobacillales* tree and that the *Pediococcus* group is a sister of the *Leuconostoc* group, which supports the paraphyly of the *Lactobacillus* genus. Lefébure and Stanhope (2007) applied comparative evolutionary analysis on 26 *Streptococcus* genomes to assess levels of recombination and positive selection in pathogen adaptation to their hosts. This study indicates that positive selection appears to be of main importance in species differentiation and adaptation to new hosts, whereas the process of recombination

has a role in strain evolution. Claesson and coworkers (2008) conducted a phylogenomic study on 12 *Lactobacillus* genomes and observed incongruence between the tree based on 141 core proteins and single-gene trees. The source of incongruence is so far unknown, but is likely due to different evolutionary rates among the genes, hidden paralogies, or horizontal gene transfer. Finally, Zhang and coworkers (2011) studied the phylogenetic relationships among 28 LAB (comprising 7 genera of 4 families) genome sequences based on 232 core genes. The concatenation of these genes allowed the recovery of a strongly supported phylogeny, providing a maximum and decisive resolution of the relationships among the LAB species examined.

## Part III

### Experimental Work





# 4

## Traditional Workflow in Microbial Diversity Analyses

Traditional methods for microbial diversity analyses were applied to two food ecosystems. The first ecosystem studied was the spontaneous cocoa bean fermentation ecosystem. Cocoa beans, which are the seeds of the fruit pods of the cocoa tree *Theobroma cacao*, undergo a fermentation prior to further processing. In the pods, the cocoa beans are embedded in a mass of mucilaginous white pulp, consisting of pectin, citric acid, and carbohydrates. After removal of both beans and pulp from the pods, they are piled into heaps, covered with plantain leaves, and a spontaneous fermentation starts. A succession of microbial activities takes place, encompassing a restricted species diversity of yeasts, LAB, and AAB. An isolation campaign was carried out by Papalexandratou and coworkers (2011a; 2011c; 2011d) from spontaneously cocoa bean fermentations carried out in Brasil, Ecuador, and Malaysia. The resulting isolates were subsequently dereplicated using rep-PCR fingerprinting. This dereplication gained a set of isolates that could not be identified on the species level. Therefore, the polyphasic taxonomy approach was applied to these isolates in Section 4.1, which led to the description of *Weissella fabalis* sp. nov. and the identification of several *Fructobacillus tropaeoli* strains.

The second ecosystem studied was rutabaga (*Brassica napobrassica*), which is a cruciferous plant with a thick bulbous edible yellow root. The edible yellow root was sous-vide cooked, packaged, and stored. Sous-vide cooking or vacuum

cooking is defined as: 'raw materials or raw materials with intermediate foods that are cooked under controlled conditions of temperature and time inside heat-stable vacuumized pouches'. Lower temperatures (usually under 100°C) and longer cooking times than those used in traditional cooking are typically applied. A Finnish laboratory isolated a set of strains from sous-vide cooked rutabaga. These isolates were dereplicated by means of MALDI-TOF MS and resulted in the description of a novel *Leuconostoc rapi* sp. nov. using the polyphasic taxonomy approach (Section 4.2).

Finally, Peeters and coworkers (2011) obtained several Antarctic isolates using plate culturing and subsequent dereplication using rep-PCR. This resulted in a set of unidentified isolates, which were subsequently identified in Section 4.3 as a novel species of the genus *Carnobacterium* (i.e., *Carnobacterium iners* sp. nov.) using the polyphasic taxonomy approach. Furthermore, this novel species description was accompanied by the construction of an MLSA scheme for the genus *Carnobacterium*, which includes food-grade LAB species.



## 4.1 Characterization of Strains of *Weissella fabalis* sp. nov. and *Fructobacillus tropaeoli* from Spontaneous Cocoa Bean Fermentations

I. Snauwaert, Z. Papalexandratou, L. De Vuyst, P. Vandamme

Redrafted from: International Journal of Systematic and Evolutionary Microbiology, 2013 (63), 1709-16.

Author contributions: IS performed the experiments and wrote the manuscript. ZP did the isolation and dereplication of the strains. PV and IS designed the experiment. LDV and PV proofread the manuscript.

Six facultatively anaerobic, non-motile LAB were isolated from spontaneous cocoa bean fermentations carried out in Brazil, Ecuador, and Malaysia. Phylogenetic analysis revealed that one of these strains, designated M75<sup>T</sup>, isolated from a Brazilian cocoa bean fermentation, had the highest 16S rRNA gene sequence similarity towards *Weissella fabaria* LMG 24289<sup>T</sup> (97.7%), *Weissella ghanensis* LMG 24286<sup>T</sup> (93.3%), and *Weissella beninensis* LMG 25373<sup>T</sup> (93.4%). The remaining LAB isolates, represented by strain M622, showed the highest 16S rRNA gene sequence similarity towards the type strain (T) of *Fructobacillus tropaeoli* (99.9%), a recently described species isolated from a flower in South Africa. *pheS* gene sequence analysis indicated that the former strain represented a novel species, whereas *pheS*, *rpoA*, and *atpA* gene sequence analysis indicated that the remaining five strains belonged to *F. tropaeoli*; these results were confirmed by DDH experiments towards their respective nearest phylogenetic neighbors. Additionally, MALDI-TOF MS proved successful for the identification of species of the genera *Weissella* and *Fructobacillus* and for the recognition of the novel species. We propose to classify strain M75<sup>T</sup> (=LMG 26217<sup>T</sup> =CCUG 61472<sup>T</sup>) as the type strain of the novel species *Weissella fabalis* sp. nov.

### 4.1.1 Introduction

Papalexandratou and colleagues (2011a; 2011c; 2011d) analyzed fermenting cocoa pulp-bean mass samples collected from traditional Brazilian (October–November 2006), Ecuadorian (April 2008), and Malaysian (April 2010) cocoa bean box and platform fermentations. In these studies, they isolated a large number of LAB that were first screened and identified by (GTG)<sub>5</sub>-PCR fingerprinting. However, this approach failed to identify one isolate, designated M75<sup>T</sup>, from a Brazilian cocoa bean box fermentation, with a unique (GTG)<sub>5</sub>-PCR fingerprint and another 63 LAB isolates (two, 58 and three isolates from Brazil, Ecuador, and Malaysia, respectively) with similar (GTG)<sub>5</sub>-PCR fingerprints. The precise taxonomic positions of six of these LAB, which belonged to five different rep clusters [clusters I–III, M75<sup>T</sup>, M56, and M622 (Brazil); cluster IV, M1588 and M1190 (Malaysia and Ecuador, respectively); cluster V, M710 (Ecuador)] were determined in the present study.

### 4.1.2 Methods, Results, and Discussion

Analysis of the 16S rRNA gene sequences of strains M75<sup>T</sup> and M622 was performed as described by De Bruyne *et al.* (2008a), except that sequencing reactions were purified using the BigDye xTerminator purification kit according to the protocol of the supplier (Applied Biosystems). The ARB software package (Ludwig *et al.*, 2004) and the corresponding SILVA SSURef 102 database (Pruesse *et al.*, 2007) were used to align the 16S rRNA gene sequences obtained and those of the type strains of all established species of the genera *Fructobacillus* and *Weissella*, their nearest phylogenetic neighbors (see below). These aligned sequences were imported into the software package molecular evolutionary genetics analysis (MEGA) version 5.0 (Tamura *et al.*, 2011) and analyzed using the neighbor-joining, maximum-likelihood (ML), and maximum-parsimony methods. The statistical reliability of tree topologies was evaluated by bootstrapping analysis based on 1000 tree replicates. The maximum-parsimony and neighbor-joining trees (not shown) revealed topologies similar to those obtained in the phylogenetic tree reconstructed using the ML approach (Figure 7). Sequence similarity calculations performed using the ARB software package indicated that the closest relatives of strain M75<sup>T</sup> were *W. fabaria* LMG 24289<sup>T</sup> (97.7%), *W. ghanensis* LMG 24286<sup>T</sup> (93.3%), and *W. beninensis* 2L24P13<sup>T</sup> (93.4%). Lower sequence similarities (< 89%) were found towards other species of the genus *Weissella* with validly published names. Strain M622 was most closely related to *F. tropaeoli* F214-1<sup>T</sup> (99.9%) and *F. pseudoficulneus* LC 51<sup>T</sup> (99.2%). Lower sequence similarities (< 98%) were found towards other species of the genus *Fructobacillus* with validly published names.




Figure 7: ML tree based on 16S rRNA gene sequences showing the phylogenetic relationships of strains M75<sup>T</sup> and M622 among the type strains of all species of the genera *Weissella* and *Fructobacillus*

Bootstrap values (%) based on 1000 replications are shown at branch points. The substitution model used was the general time reversible (GTR) model and the aligned sequence had a length of 1345 bp. Bar, 1% sequence divergence.

Sequence analysis of the housekeeping genes *pheS*, *rpoA*, and *atpA* correlates with species delineation as determined by DDH in the genus *Fructobacillus* (De Bruyne *et al.*, 2007). Therefore, *pheS*, *rpoA*, and *atpA* gene sequences of strains M56, M622, M710, M1190, and M1588 and *F. tropaeoli* LMG 26298<sup>T</sup>, for which no *pheS*, *rpoA*, or *atpA* sequences were available, were determined as described previously (De Bruyne *et al.*, 2007), except that sequencing reactions were purified using

the BigDye xTerminator purification kit as described above. Similarly, sequence analysis of the *pheS* gene has been used successfully to distinguish all established species of the genus *Weissella* (De Bruyne *et al.*, 2008a,0). Therefore, *pheS* gene sequences were determined for strain M75<sup>T</sup> and for *W. beninensis* LMG 25373<sup>T</sup> and *W. fabaria* M1160 and M1167, for which no *pheS* gene sequences were available. *pheS*, *rpoA*, and *atpA* gene sequences of all remaining *Fructobacillus* and *Weissella* reference strains were available from previous studies (De Bruyne *et al.*, 2007). Details of strains, depositors, and accession numbers are given in Table 2. SeaView version 4 was used to concatenate the *pheS*, *rpoA*, and *atpA* gene sequences (Gouy *et al.*, 2010) and the software package MEGA version 5.0 (Tamura *et al.*, 2011) was used to align the translated concatenated gene sequences and to analyze the nucleotide sequences as mentioned above.

The maximum-parsimony and neighbor-joining trees (not shown) revealed topologies similar to those obtained in the phylogenetic tree reconstructed using the ML approach (Figures 8 and 9) for both analyses. The concatenated *pheS*, *rpoA*, and *atpA* gene sequences of strains M56, M622, M710, M1190, and M1588 revealed high similarity to the concatenated sequence of *F. tropaeoli* LMG 26298<sup>T</sup> (98.4, 97.4, 99.2, 99.3, and 99.3% similarity, respectively). Lower gene sequence similarities were found (< 95%) towards other species of the genus *Fructobacillus* with validly published names. The phylogenetic tree of the concatenated *pheS*, *rpoA*, and *atpA* gene sequences confirmed the discriminatory power of these sequences for species identification within the genus *Fructobacillus*: all species were clearly delineated above 97% concatenated gene sequence similarity (Figure 8). Based on these results and the previously established correlation between MLSA of concatenated *pheS*, *rpoA*, and *atpA* gene sequences and levels of DDH, we concluded that strains M56, M622, M710, M1190, and M1588 belong to *F. tropaeoli*. In addition, pairwise *pheS* gene sequence similarity calculations, calculated using the MEGA software version 5.0, confirmed that the closest relatives of strain M75<sup>T</sup> were *W. fabaria* LMG 24289<sup>T</sup> (85.2%), *W. ghanensis* LMG 24286<sup>T</sup> (86.9%), and *W. beninensis* LMG 25373<sup>T</sup> (80%). Lower *pheS* gene sequence similarities (< 78%) were found towards other species of the genus *Weissella* with validly published names. With the exception of *Weissella viridescens*, De Bruyne *et al.* (2010; 2008a) demonstrated that strains belonging to the same species of the genus *Weissella* share *pheS* gene sequence similarity of at least 96.8%. Therefore, the present results suggest that strain M75<sup>T</sup> represents a novel species of the genus *Weissella* (Figure 9).

Table 2: *Weissella* and *Fructobacillus* strains included in the phylogenetic study and in the MALDI-TOF MS analysis.

Strain	Depositor	Source	Accession numbers		
			<i>pheS</i>	<i>rpoA</i>	<i>atpA</i>
<b><i>W. beninensis</i></b> LMG 25373 <sup>T</sup>	SW. Padonou	Submerged fermented cassava (Benin)	HE576797		
<b><i>W. cibaria</i></b> LMG 13587	L. Devriese	Dog, ear (Belgium)	FM202101		
LMG 17699 <sup>T</sup>	G. Rusul	Chili bo (Malaysia)	FM202102		
LMG 17704	G. Rusul	Chili bo (Malaysia)	FM202100		
LMG 18507	NCFB	Post-harvest deterioration of sugar cane	FM202098		
LMG 21843	KCTC	Partially fermented kimchi (Korea)	FM202099		
R-31690	CCMM	Fermented skimmed milk (Morocco)	FM202103		
<b><i>W. confusa</i></b> LMG 9497 <sup>T</sup>	A. Ledebøer	Sugar cane	FM202105		
LMG 11983	ATCC	Grass silage	FM202104		
LMG 14040	L. Devriese	Dog, ear (Belgium)	FM202107		
LMG 17718	G. Rusul	Chili bo (Malaysia)	FM202108		
LMG 18480	G. Rusul	Tapai (Malaysia)	FM202106		
<b><i>W. fabaria</i></b> LMG 24289 <sup>T</sup>	Own isolate	Fermenting cocoa pulp-bean mass (Ghana)	FM202097		
252 (R-34084)	Own isolate	Fermenting cocoa pulp-bean mass (Ghana)	FM202096		
M1160 (R-46910)	Own isolate	Fermenting cocoa pulp-bean mass (Ecuador)	HE577177		
M1167 (R-46912)	Own isolate	Fermenting cocoa pulp-bean mass (Ecuador)	HE577176		
<b><i>W. ghanensis</i></b> LMG 24286 <sup>T</sup>	Own isolate	Fermenting cocoa pulp-bean mass (Ghana)	FM202095		
194B (R-27442)	Own isolate	Fermenting cocoa pulp-bean mass (Ghana)	FM202094		
<b><i>W. halotolerans</i></b> LMG 9469 <sup>T</sup>	DSMZ	Sausage	FM202114		
<b><i>W. hellenica</i></b> LMG 15125 <sup>T</sup>	NCFB	Naturally fermented sausage (Greece)	FM202110		
<b><i>W. kandleri</i></b> LMG 18979 <sup>T</sup>	NCIMB	Desert spring (Namibia)	FM202116		
<b><i>W. koreensis</i></b> LMG 21853 <sup>T</sup>	KCTC	Kimchi (Korea)	FM202115		
<b><i>W. minor</i></b> LMG 9847 <sup>T</sup>	NCFB	Slime from milking machine	FM202117		
<b><i>W. paramesenteroides</i></b> LMG 9852 <sup>T</sup>	NCFB	Fermented dry salami	FM202111		
<b><i>W. soli</i></b> LMG 20113 <sup>T</sup>	S. Roos	Garden soil (Sweden)	FM202113		
LMG 20114	S. Roos	Garden soil (Sweden)	FM202112		
<b><i>W. thailandensis</i></b> LMG 19821 <sup>T</sup>	JCM	Fermented fish (Thailand)	FM202109		

Continued on next page

Table 2 – continued from previous page

Strain	Depositor	Source	Accession numbers		
			<i>pheS</i>	<i>rpoA</i>	<i>atpA</i>
<b><i>W. viridescens</i></b>					
LMG 3507 <sup>T</sup>	NCFB	Cured meat products	FM202120		
LMG 11497	NCFB	NK	FM202122		
LMG 12021	JCM	NK	FM202121		
LMG 13093	NCFB	Frankfurters	FM202119		
LMG 23120	J. Björkroth	Spanish blood sausage (Spain)	FM202118		
<b><i>Weissella</i> sp. nov.</b>					
M75 <sup>T</sup> (LMG 26217 <sup>T</sup> )	Own isolate	Fermenting cocoa pulp-bean mass (Brazil)	HE576796		
<b><i>F. durionis</i></b>					
LMG 22556 <sup>T</sup>	J. Leisner	Tempoyak made from durian fruit (Malaysia)	AM711166	AM711309	AM711205
LMG 22557	J. Leisner	Tempoyak made from durian fruit (Malaysia)	AM711140	AM711290	AM711171
LMG 22558	J. Leisner	Tempoyak made from durian fruit (Malaysia)	AM711141	AM711291	AM711172
<b><i>F. ficulneus</i></b>					
LMG 21928 <sup>T</sup>	DSMZ	Ripe fig (Portugal)	AM711151	AM711342	AM711183
<b><i>F. fructosus</i></b>					
LMG 9498 <sup>T</sup>	A.Ledeboer	Flower (Japan)	AM711194	AM711321	AM711174
<b><i>F. pseudoficulneus</i></b>					
LMG 23899 <sup>T</sup>	CECT	Ripe fig (Portugal)	AM711281	AM711355	AM711274
R-35156	R. Tenreiro	Ripe fig (Portugal)	AM711235	AM711335	AM711259
R-35157	R. Tenreiro	Ripe fig (Portugal)	AM711236	AM711336	AM711260
R-35158	R. Tenreiro	Ripe fig (Portugal)	AM711237	AM711337	AM711261
R-35159	R. Tenreiro	Ripe fig (Portugal)	AM711238	AM711338	AM711262
R-35160	R. Tenreiro	Ripe fig (Portugal)	AM711239	AM711333	AM711263
<b><i>F. tropaeoli</i></b>					
LMG 26298 <sup>T</sup>	L. Dicks	Flower (South Africa)	HE590678	HE590679	HE590680
M56 (R-46388)	Own isolate	Fermenting cocoa pulp-bean mass (Brazil)	HE590681	HE590682	HE590683
M622 (R-46389)	Own isolate	Fermenting cocoa pulp-bean mass (Brazil)	HE590684	HE590685	HE590686
M710 (R-46397)	Own isolate	Fermenting cocoa pulp-bean mass (Brazil)	HE590687	HE590688	HE590689
M1190 (R-46399)	Own isolate	Fermenting cocoa pulp-bean mass (Brazil)	HE590690	HE590691	HE590692
M1588 (R-46401)	Own isolate	Fermenting cocoa pulp-bean mass (Brazil)	HE590693	HE590694	HE590695

**Depositors listed in Table 2:** S. W. Padonou, Université d'Abomey-Calavi, Cotonou, Benin; L. Devriese, Ghent University, Ghent, Belgium; G. Rusul, University Putra, Malaysia; NCFB, National Collection of Food Bacteria (now NCIMB), Aberdeen, UK; KCTC, Korean Collection for Type Cultures, Yusong, Taejeon, Republic of Korea; CCMM, Moroccan Coordinated Collections of Microorganisms, Rabat, Morocco; A. Ledeboer, Unilever, Vlaardingen, The Netherlands; ATCC, American Type Culture Collection, Manassas, VA, USA; DSMZ, Deutsche Sammlung von Mikroorganismen und Zellkulturen, Braunschweig, Germany; NCIMB, National Collections of Industrial, Food and Marine Bacteria, Aberdeen, UK; S. Roos, Swedish University of Agricultural Sciences, Uppsala, Sweden; JCM, Japan Collection of Microorganisms, RIKEN BRC, Japan; J. Björkroth, University of Helsinki, Helsinki, Finland; J. Leisner, Royal Veterinary and Agricultural University, Frederiksberg, Copenhagen, Denmark; CECT, Colección Española de Cultivos Tipo, Valencia, Spain; R. Tenreiro, University of Lisbon, Lisbon, Portugal; L. Dicks, University of Stellenbosch, South Africa. NK, Not known.




Figure 8: ML tree based on concatenated *pheS*, *rpoA*, and *atpA* gene sequences showing the phylogenetic relationships of strains M56, M622, M710, M1190, and M1588 among other members of the genus *Fructobacillus*

Bootstrap values (%) based on 1000 replications are shown at branch points. The substitution model used was the GTR model and the aligned concatenated sequence had a length of 903 bp. Bar, 2% sequence divergence. Accession numbers are given in Table 2.




Figure 9: ML tree based on *pheS* gene sequences showing the phylogenetic relationships of strain M75<sup>T</sup> among other members of the genus *Weissella*

Bootstrap values (%) based on 1000 replications are shown at branch points. The substitution model used was the GTR model and the aligned concatenated sequence had a length of 369 bp. Bar, 2% sequence divergence. Accession numbers are given in Table 2



To confirm these results, DDH experiments between strains M75<sup>T</sup> and M1190 and their respective nearest neighbors were performed. Genomic DNA of strain M75<sup>T</sup>, *W. ghanensis* LMG 24286<sup>T</sup>, *W. fabaria* LMG 24289<sup>T</sup>, M1190, and *F. tropaeoli* LMG 26298<sup>T</sup> was extracted using the guanidine thiocyanate method described by Pitcher *et al.* (1989). DDHs were performed using the microplate method, with photobiotin for labeling of the DNA (Ezaki *et al.*, 1989), as modified by Goris *et al.* (1998). The hybridization levels of strain M75<sup>T</sup> towards *W. ghanensis* LMG 24286<sup>T</sup> and *W. fabaria* LMG 24289<sup>T</sup> were 44 and 36%, respectively, confirming that it represents a distinct species of the genus *Weissella*. The DDH between strain M1190 and *F. tropaeoli* LMG 26298<sup>T</sup> was 82%, which confirmed its MLSA-based identification.

DNA G+C content was determined according to the enzymatic DNA degradation method described previously (Mesbah & Whitman, 1989) using a Waters Breeze high-performance liquid chromatography (HPLC) system and XBridge Shield RP18 column. The solvent used was 0.02 M NH<sub>4</sub>H<sub>2</sub>PO<sub>4</sub> (pH 4.0)/1.5% (v/v) acetonitrile. Non-methylated lambda phage DNA (Sigma) was used as a calibration reference and *Escherichia coli* LMG 2093 DNA was included as a control. The DNA G+C contents of strains M75<sup>T</sup> and M1190 were 37 and 45 mol%, respectively, consistent with the G+C contents found previously in the genera *Weissella* (37–47 mol%; (Björkroth *et al.*, 2002; Collins *et al.*, 1993; De Bruyne *et al.*, 2008a; Lee *et al.*, 2002; Magnusson *et al.*, 2002; Padonou *et al.*, 2010; Tanasupawat *et al.*, 2000; Vela *et al.*, 2011) and *Fructobacillus* (42–45 mol%; (Endo & Okada, 2008; Endo *et al.*, 2010).

Preparation of peptidoglycan and analysis of peptidoglycan structure were performed according to published protocols (Schumann, 2011). The total hydrolysate (4 M HCl, 100 °C, 16 h) of the peptidoglycan contained the amino acids lysine, alanine, serine, glycine and glutamic acid in an approximate ratio of 1.0 : 1.7 : 0.9 : 0.2 : 2.6. Because the ratio of glutamic acid to lysine was < 1 and did not fit the expected A3 $\alpha$  peptidoglycan type, the peptidoglycan preparation was subjected to hydrofluoric acid treatment in order to remove contaminating polymers linked to the peptidoglycan via phosphodiester bonds before repeating the analysis. However, the molar ratio of the peptidoglycan amino acids changed only slightly after the hydrofluoric acid treatment. The identity of the amino acids was confirmed by gas chromatography coupled to mass spectrometry (GC-MS). The partial hydrolysate (4 M HCl, 100 °C, 45 min) contained, in addition to the amino acids, the peptides L-Ala-D-Glu-L-Lys-L-Ala, D-Ala-L-Lys, and D-Ala-L-Lys-L-Ala. The peptide Ala-Ala could not be found. Dinitrophenylation according to Schleifer (1985) revealed that serine represents the N terminus of the interpeptide bridge.

Though the quantitative amino acid ratio contained too much glutamic acid and too little alanine, the peptidoglycan type A3 $\alpha$  L-Lys-L-Ala-L-Ser is concluded from the N terminus of the interpeptide bridge and the 2D thin layer chromatography (TLC) patterns of peptides and amino acids. The high content of glutamic acid as well as the traces of glycine might be caused by residual polymer contamination.

MALDI-TOF MS has shown to be both rapid and accurate for species and sub-species classification of a broad spectrum of bacteria (Sauer & Kliem, 2010). To test whether MALDI-TOF MS analysis is a suitable tool for the identification of the LAB of the present study, mass spectra were generated from all *Weissella* and *Fructobacillus* strains (Table 2) except for *Weissella hellenica* LMG 15125<sup>T</sup>, *W. cibaria* R-31690, *W. confusa* LMG 18480, *W. fabaria* 252, and *F. pseudoficulneus* R-35156 and R-35158, from which good-quality spectra could not be obtained. Bacteria were grown under standardized conditions [28 °C, 24 h, de Man-Rogosa-Sharpe (MRS) agar (Oxoid) and an aerobic atmosphere] and subcultured twice prior to analysis. For each strain, a spectral profile was generated from three different generations to improve the robustness of the library of mass spectral fingerprints. Harvesting of cells, extract preparation, measuring, and data analysis were performed as described previously (Ghyselinck *et al.*, 2011). The similarity between the spectra of strains M75<sup>T</sup>, M56, M622, M710, M1190, and M1588 and the reference strains was calculated using Pearson's product moment correlation coefficient and clustering was performed using the unweighted pair group method with arithmetic means (UPGMA) clustering algorithm (Figures 4.1 and 4.2, available in Supplementary Material). The MALDI-TOF MS profile of strain M75<sup>T</sup> formed a separate cluster that was most similar to the protein profiles of recently described *W. ghanensis* and *W. fabaria* strains originating from Ghanaian cocoa bean fermentations. The remaining cocoa isolates showed high similarity to the recently described *F. tropaeoli*, isolated from a South African flower. For the genus *Fructobacillus*, the intraspecies diversity was consistently smaller than the interspecies divergence towards their nearest neighbors, as was the case for the genus *Weissella*.

Finally, a biochemical analysis of the cocoa isolates was performed to characterize the novel species and to compare the characteristics of the novel *F. tropaeoli* strains (cocoa origin) with those of the type (and currently only known) strain (Chambel *et al.*, 2006; Endo *et al.*, 2010). Cell and colony morphology of strains M56, M622, M710, M1190, and M1588 was verified after growth on MRS agar (Oxoid) supplemented with 1% (w/v) fructose and 24 h of aerobic incubation at 28 °C. For strain M75<sup>T</sup>, MRS agar (Oxoid) was used as the basal medium. Conventional bio-

chemical characteristics and enzyme activities were tested as described previously in triplicate, unless stated otherwise (De Bruyne *et al.*, 2007). Growth was tested at 4, 15, 20, 37, and 42°C and in the presence of 6.5, 8, and 10% NaCl (w/v). The production of gas from glucose was determined using inverted Durham tubes. The API 50 CHL *Lactobacillus* identification system (bioMérieux) proved useful for determining carbohydrate fermentation profiles. The production of D- and L-lactate from glucose was determined enzymatically (B-Biopharm). Unlike the *F. tropaeoli* isolates, strain M75<sup>T</sup> was able to produce dextran in MRS agar (Oxoid) in which glucose had been replaced with 5% sucrose. The physiological and biochemical characteristics of the *F. tropaeoli* isolates were identical to those of *F. tropaeoli* LMG 26298<sup>T</sup>, with the only difference that the isolates were also able to ferment sucrose, in addition to D-glucose, D-fructose, and D-mannitol. The results for strain M75<sup>T</sup> are given in the species description. Characteristics that differentiate strain M75<sup>T</sup> from other species of the genus *Weissella* are summarized in Table 3.

#### 4.1.3 Conclusion

In conclusion, the results of the present study demonstrate that strains M56, M622, M710, M1190, and M1588 belong to *F. tropaeoli*. These results and those of Papalexandratou (2011a; 2011c; 2011d) demonstrate that *F. tropaeoli* participates in the fermentation process of cocoa beans in Brazil, Ecuador, and Malaysia, which are major cocoa-producing countries. Additionally, the results of the present study demonstrate that strain M75<sup>T</sup> represents a novel species of the genus *Weissella* that is closely related to *W. fabaria*, *W. ghanensis*, and *W. beninensis* (96.9, 94.9, and 94.2% 16S rRNA gene sequence similarity towards the respective type strains), but which can be distinguished from these and other species of the genus *Weissella* by DDH, *pheS* gene sequence analysis, MALDI-TOF MS-based analysis, and biochemical characteristics. Based on these results, we propose to classify strain M75<sup>T</sup> as the type strain of the novel species *Weissella fabalis* sp. nov.

Table 3: Differential characteristics of strain *M75<sup>T</sup>* (*W. fabalis* sp. nov.) and other species of the genus *Weissella*

Characteristic	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
<b>Acid from:</b>															
Arabinose	-	-	-	-	-	-	+	-	+	+	-	+	+	d	-
Cellobiose	+	+	+	-	+	-	-	-	-	+	+	-	-	d	d
Fructose	+	+	+	+	+	+	-	+	ND	+	+	+	+	+	+
Galactose	-	-	-	-	-	-	-	+	-	-	+	+	-	+	+
Maltose	+	-	+	+	+	+	+	-	-	+	+	+	+	+	+
Melibiose	-	-	-	-	-	-	+	-	-	-	-	+	-	+	+
Raffinose	-	-	-	-	-	-	+	-	-	-	-	+	-	d	+
Ribose	-	-	-	+	+	-	+	+	+	-	+	+	-	d	d
Salicin	-	-	+	-	-	-	+	-	ND	+	+	-	-	-	d
Sucrose	-	-	d	-	+	d	+	-	-	+	+	d	+	+	+
Trehalose	+	+	+	-	+	d	+	-	-	-	-	d	+	+	d
Xylose	-	-	-	-	-	-	+	-	+	+	+	-	-	d	-
<b>Hydrolysis from esculin</b>	+	+	+	-	+	-	+	-	-	+	+	-	ND	d	+
<b>NH<sub>3</sub> from arginine</b>	+	+	+	+	+	-	+	+	+	+	+	-	-	-	+
<b>Dextran formation</b>	+	+	+	ND	-	ND	-	+	+	+	+	-	-	-	ND
<b>Lactic acid configuration</b>	D	DL	DL	DL	DL	DL	D	D	D	DL	D	D	D	D	D
<b>Growth at/in:</b>															
15°C	+	+	+	ND	ND	ND	+	ND	+	+	ND	ND	ND	ND	+
37°C	+	+	+	ND	-	ND	-	+	+	+	+	-	-	-	ND
42°C	+	+	+	ND	ND	ND	+	ND	+	+	ND	+	ND	ND	ND
6.5% NaCl	-	-	-	ND	ND	ND	-	ND	-	+	ND	-	ND	ND	-
8.0% NaCl	-	-	-	ND	ND	ND	ND	ND	ND	+	ND	ND	ND	ND	ND
10.0% NaCl	-	-	-	ND	ND	ND	ND	ND	-	-	ND	ND	ND	ND	ND
<b>DNA G+C content (mol%)</b>	37	38	40	44	44	41-44	43	39	37	44-45	45-47	38-41	39-40	37-38	37

Species: 1, *W. fabalis* sp. nov.; 2, *W. fabaria*; 3, *W. ghanensis*; 4, *W. halotolerans*; 5, *W. minor*; 6, *W. viridescens*; 7, *W. soli*; 8, *W. kandleri*; 9, *W. koreensis*; 10, *W. cibaria*; 11, *W. confusa*; 12, *W. thailandensis*; 13, *W. hellenica*; 14, *W. paramesenteroides*; 15, *W. beninensis*. +, 90% or more strains positive; 2, 90% or more strains negative; d, 11-89% of strains positive; ND, no data available. Data partially adapted from (Collins *et al.*, 1993; Tanasupawat *et al.*, 2000; De Bruyne *et al.*, 2008a,b; Padonou *et al.*, 2010).

*Description of Weissella fabalis* sp. nov.

*Weissella fabalis* (fa.ba'lis. L. fem. adj. *fabalis* belonging to beans. Cells are Gram-stain-positive, catalase-negative, facultatively anaerobic, and non-motile. Cells are coccoid, approximately 1.0  $\mu\text{m}$  wide and 1.5  $\mu\text{m}$  long, and occur singly, in pairs or in short chains. Colonies grown for 2 days on MRS agar at 28°C are approximately 1 mm in diameter, beige, opaque, smooth, and circular, with a low-convex elevation. Gas is produced from glucose, indicating the heterofermentative character of the type strain. Produces D-lactic acid. Growth occurs at 15–37°C and in the presence of 5–6% NaCl but not in the presence of 7–8% NaCl. Arginine is hydrolyzed. Acid is produced from glucose, fructose, mannose, N-acetylglucosamine, aesculin, cellobiose, maltose, trehalose, and gentiobiose. Acid is not produced from glycerol, erythritol, D- or L-arabinose, ribose, D- or L-xylose, adonitol, methyl  $\beta$ -D-xylopyranoside, galactose, sorbose, rhamnose, dulcitol, inositol, mannitol, sorbitol, methyl  $\alpha$ -D-mannopyranoside, methyl  $\alpha$ -D-glucopyranoside, amygdalin, arbutin, salicin, lactose, melibiose, sucrose, inulin, melezitose, raffinose, starch, glycogen, xylitol, turanose, D-lyxose, D-tagatose, D- or L-fucose, D- or L-arabitol, potassium gluconate, potassium 2-ketogluconate, or potassium 5-ketogluconate. The DNA G+C content of the type strain is 37 mol%.


The type strain, M75<sup>T</sup> (=LMG 26217<sup>T</sup> =CCUG 61217<sup>T</sup>), was isolated from a Brazilian cocoa bean box fermentation carried out in Ilhéus, Bahia, Brazil, in 2007.

#### 4.1.4 Acknowledgements

We acknowledge financial support from the Fund for Scientific Research (FWO)-Flanders and thank Anneleen Wieme and Freek Spitaels for their contributions to the development of the MALDI-TOF MS protocol. Financial support from the Research Council of the Vrije Universiteit Brussel and Barry Callebaut N.V. is also acknowledged.


### 4.1.5 Supplementary Material

The GenBank/EMBL/DDBJ (DNA Data Bank of Japan) accession numbers for the 16S rRNA and *pheS* gene sequences of strain M75<sup>T</sup> are HE576795 and HE576796. Accession numbers of other sequences obtained in this study are detailed in Table 2.



Supplementary Material 4.1: MALDI-TOF MS dendrogram constructed using the Pearson product moment correlation coefficient and the UPGMA clustering algorithm, showing the similarity of strain M75<sup>T</sup> with other strains of the genus *Weissella*.

All strains of Table 2 are incorporated in the dendrogram, with the exception of LMG 15125<sup>T</sup>, R-31690, and LMG 18480.



Supplementary Material 4.2: MALDI-TOF MS dendrogram constructed using the Pearson product moment correlation coefficient and the UPGMA clustering algorithm, showing the similarity of strains M56, M622, M710, M1190, and M1588 with other strains of the genus *Fructobacillus*.

All strains of Table 2 are incorporated in the dendrogram, with the exception of R-35156 and R-35158.





## 4.2 *Leuconostoc rapi* sp. nov., Isolated from Sous-Vide Cooked Rutabaga

U. Lyhs, I. Snauwaert, S. Pihlajaviita, L. De Vuyst, P. Vandamme

Submitted to: International Journal of Systematic and Evolutionary Microbiology, 2014

Author contributions: UL wrote the manuscript. IS performed the experiments. SP did the isolation of the strains. PV and IS designed the experiment. PV and LDV proofread the manuscript.

A Gram-stain-positive, ovoid LAB, strain LMG 27676<sup>T</sup>, was isolated from a spoiled sous-vide cooked rutabaga. 16S rRNA gene sequence analysis indicated that the novel strain belongs to the genus *Leuconostoc*, with *Leuconostoc kimchi* and *Leuconostoc miyukkimchii* as nearest neighbors (99.1 and 98.8% 16S rRNA gene sequence similarity towards the type strain, respectively). Phylogenetic analysis of the 16S rRNA gene, MLSA of the *pheS*, *rpoA*, and *atpA* genes, and biochemical and genotypic characterization allowed to differentiate strain LMG 27676<sup>T</sup> from all established *Leuconostoc* species. Strain LMG 27676<sup>T</sup> (= DSMZ 27776<sup>T</sup>) therefore represents a new species, for which the name *Leuconostoc rapi* sp. nov. is proposed.

### 4.2.1 Introduction

The genus *Leuconostoc*, which at the time of writing comprises 12 species, belongs phylogenetically to the phylum *Firmicutes*, class *Bacilli*, order *Lactobacillales*. *Leuconostoc* cells are Gram-stain-positive, small, regular, ovoid cocci, occurring in pairs or short chains. The cells are non-motile and do not form spores. *Leuconostoc* species are considered psychrotrophic mesophiles with optimal growth at 14–30°C. The temperature limits for growth vary among species and strains, ranging from 1–10°C to 30–40°C (Säde, 2011). In 2008, *L. fructosum*, *L. durionis*, *L. ficulneum*, and *L. pseudoficulneum* were reclassified on the basis of phylogenetic, physiological, and morphological differences into the genus *Fructobacillus* gen. nov. (Endo & Okada, 2008).

Species of the genus *Leuconostoc* are associated with the fermentation of vegetables (Di Cagno *et al.*, 2013). Latest described *Leuconostoc* species originating

from fermented vegetables include *L. inhae* and *L. kimchii* from kimchi (Kim *et al.*, 2000) and *L. miyukkimchii* from fermented miyukkimchii made of a brown algae (*Undaria pinnatifida*) and is a regional kimchi in Korea (Lee *et al.*, 2012). Other novel species include *L. palmae* from palm wine, an alcoholic beverage produced from the sap of various palm tree species (Ehrmann *et al.*, 2009), and *L. holzapfelii*, from Ethiopian coffee fermentation (De Bruyne *et al.*, 2007). The occurrence of leuconostocs including *L. gelidum*, *L. gasicomitatum* (both of which were recently reclassified as *L. gelidum* subsp. *gelidum* and *L. gelidum* subsp. *gasicomitatum*, respectively (Rahkila *et al.*, 2014)), *L. citreum*, and *L. mesenteroides*, in other vegetables products has also been reported (García-Gimeno & Zurera-Cosano, 1997; Lyhs *et al.*, 2004; Vihavainen *et al.*, 2008).

Strain LMG 27676<sup>T</sup> was isolated from packaged sous-vide cooked rutabaga (*Brassica napobrassica*) during an investigation of the spoilage microbiota in sous-vide cooked vegetables in Finland. For microbiological analyses, 10 g rutabaga sample was aseptically weighed into 90 ml of buffered peptone water (BPW) (LabM, Lancashire, UK) in a sterile plastic bag and then blended for 60 s using a Dilumat automatic system (bioMérieux). Tenfold serial dilutions were used for microbiological analyses. The number of LAB was determined on MRS agar (Oxoid) supplemented with sorbic acid (pH 6.4). All plates were incubated under anaerobic conditions at 20°C for 7 days. Three of the isolates obtained, LMG 27676<sup>T</sup>, R-50028, and R-50030, exhibited identical RAPD fingerprints as determined using the primers RAPD-270 (5'-TGCGCGCGGG-3') and RAPD-272 (5'-AGCGGGCCAA-3') (Mahenthiralingam *et al.*, 1996) (data not shown).

#### 4.2.2 Methods, Results, and Discussion

Nearly complete 16S rRNA gene sequence analysis was performed as described by Vancanneyt *et al.* (2004) for strain LMG 27676<sup>T</sup>. PCR products were purified using a Nucleofast 96 PCR clean up membrane system (Machery-Nagel, Düren, Germany). The sequencing primers were those listed by Coenye *et al.* (1999) and the fragments obtained were cleaned with a BigDye XTerminator Purification kit (Applied Biosystems, Life Technologies, Carlsbad, CA, USA), according to the manufacturer's instructions. The mothur software package and the corresponding SILVA reference alignment were used to align the 16S rRNA gene sequences of strain LMG 27676<sup>T</sup> (1507 bp) with sequences of type strains of all established species of the genus *Leuconostoc*. These aligned sequences were imported into MEGA version 5.0 (Tamura *et al.*, 2011) and analyzed using the neighbor-joining,

ML, and maximum-parsimony methods. The statistical reliability of tree topologies was evaluated by bootstrapping analysis based on 1000 replicates. The neighbor-joining and the maximum-parsimony trees revealed topologies similar to those obtained in the ML tree (Figure 10). 16S rRNA gene sequence similarities were calculated using MEGA version 5.0 (Tamura *et al.*, 2011) and the closest relatives of strain LMG 27676<sup>T</sup> were *L. kimchii* IH25<sup>T</sup> (99.1%) and *L. miyukkimchii* M2<sup>T</sup> (98.8%). Lower sequence similarities (< 98.6%) were found with other members of the genus *Leuconostoc*.




Figure 10: ML tree of the 16S rRNA gene sequence of strain LMG 27676<sup>T</sup> with all type strains of the genus *Leuconostoc*

The evolutionary history was inferred by using the ML method based on the Kimura 2-paramter model. The tree with the highest log likelihood is shown. There were a total of 1379 positions in the final dataset.

Sequence analysis of the housekeeping genes encoding *pheS*, *rpoA*, and *atpA* correlates with a species delineation as determined by DDH in the genus *Leuconostoc* (De Bruyne *et al.*, 2007). Therefore, *pheS*, *rpoA*, and *atpA* gene sequences of strains *Leuconostoc rapi* sp. nov. LMG 27676<sup>T</sup>, *Leuconostoc kimchii* KCTC 2386<sup>T</sup>, and *Leuconostoc miyukkimchii* LMG 27695<sup>T</sup>, for which no *pheS*, *rpoA*, or *atpA* sequences were available, were determined as described previously (De Bruyne *et al.*, 2007), except that sequencing reactions were purified using the BigDye xTerminator purification kit as described above. The *pheS*, *rpoA*, and *atpA* gene

sequences of the closest relatives (*L. carnosum* LMG 23898<sup>T</sup>, *L. inhae* LMG 22919<sup>T</sup>, *L. gelidum* subsp. *gasicomitatum* LMG 18811<sup>T</sup>, and *L. gelidum* subsp. *gelidum* LMG 18297<sup>T</sup>, and *L. gelidum* subsp. *aenigmaticum* LMG 27840<sup>T</sup>) of strain LMG 27676<sup>T</sup> were available from previous studies (De Bruyne *et al.*, 2007; Rahkila *et al.*, 2014). The software package MEGA version 5.0 (Tamura *et al.*, 2011) was used to align the translated concatenated gene sequences and to analyze the nucleotide sequences as mentioned above. The maximum-parsimony and neighbor-joining trees (not shown) revealed topologies similar to those obtained in a phylogenetic tree reconstructed using the ML approach (Figure 11) for both analyses. The concatenated *pheS*, *rpoA*, and *atpA* gene sequences of strain LMG 27676<sup>T</sup> revealed high similarity to the concatenated sequence of *L. kimchii* KCTC 2386<sup>T</sup> (94.2%). Lower concatenated sequence similarities (< 88.1%) were found towards other species of the genus *Leuconostoc* with validly published names. The *pheS*, *rpoA*, and *atpA* gene sequences of strain LMG 27676<sup>T</sup> revealed high similarities to the sequences of *L. kimchii* KCTC 2386<sup>T</sup> (91.3, 96.4, and 94.4% similarity, respectively). Lower gene sequence similarities were found (< 84.5, 89 and 91.5%, respectively) towards other species of the genus *Leuconostoc* with validly published names. According to De Bruyne and coworkers (2007), species now classified in the genera *Leuconostoc* and *Fructobacillus* are delineated above 93, 98, and 98% *pheS*, *rpoA*, and *atpA* gene sequence similarity, respectively. Therefore, these results suggest that strain LMG 27676<sup>T</sup> represents a novel species of the genus *Leuconostoc*. To confirm this, a DDH experiment between strain LMG 27676<sup>T</sup> and *L. kimchii* strains KCTC 2386<sup>T</sup> and LMG 23786 was performed. Genomic DNA was extracted using the guanidine thiocyanate method described by Pitcher *et al.* (1989). The DDH experiment was performed using the microplate method, with photobiotin for labeling of the DNA (Ezaki *et al.*, 1989), as modified by Goris *et al.* (1998). The hybridization levels of strain LMG 27676<sup>T</sup> towards *L. kimchii* KCTC 2386<sup>T</sup> was 51% (the reciprocal hybridization values were 53 and 49%); the hybridization level between *L. kimchii* KCTC 2386<sup>T</sup> and *L. kimchii* LMG 23786 was 85% (the reciprocal hybridization values were 72 and 97%); and the hybridization level between strain LMG 27676<sup>T</sup> and *L. kimchii* LMG 23786 was 64% (the reciprocal hybridization values were 70 and 57%). The hybridization values between strain LMG 27676<sup>T</sup> and both *L. kimchi* strains were clearly below the species delineation threshold.

DNA G+C content was determined according to the enzymatic DNA degradation method described previously (Mesbah & Whitman, 1989), using a Waters Breeze HPLC system and XBridge Shield RP18 column. The solvent used was 0.02 M NH<sub>4</sub>H<sub>2</sub>PO<sub>4</sub> (pH 4.0)/1.5% (v/v) acetonitrile. Non-methylated lambda phage DNA (Sigma-Aldrich, St. Louis, MO, USA) was used as a calibration reference and *Escherichia coli* LMG 2093 DNA was included as a control. The DNA G+C content




Figure 11: ML tree of the concatenated *pheS*, *rpoA*, and *atpA* gene sequences of the closest relatives of strain LMG 27676<sup>T</sup>

The evolutionary history was inferred by using the ML method based on the Tamura 3-parameter model. The tree with the highest log likelihood is shown. There were a total of 1482 positions in the final dataset. The type strain of *L. kimchii* was not available and was consequently replaced by strain LMG 23787

of strain LMG 27676<sup>T</sup> was 38 mol% which is consistent with the G+C contents found previously in the genus *Leuconostoc* (38–44 mol%; (De Bruyne *et al.*, 2007).

Preparation of peptidoglycan and analysis of the peptidoglycan structure were performed according to published protocols (Schumann, 2011). The total hydrolysate (4 N HCl, 100°C, 16 h) contained the amino acids alanine, glutamic acid and lysine, and analysis of partial hydrolysates (4 N HCl, 100°C, 45 min) revealed the presence of the peptidoglycan type A3α L-Lys-Ala<sub>2</sub> or type A11.5 according to [www.peptidoglycan-types.info](http://www.peptidoglycan-types.info).

Finally, a biochemical analysis of LMG 27676<sup>T</sup>, KCTC 2386<sup>T</sup>, and LMG 23786 was performed to characterize the novel species. Cell and colony morphology of

strain LMG 27676<sup>T</sup> was verified after growth on MRS agar (Oxoid) and 96 h of aerobic incubation at 28°C. Conventional biochemical characteristics and enzyme activities were tested as described previously in triplicate, unless stated otherwise (De Bruyne *et al.*, 2007). Growth was tested at 4, 20, 30, 37, and 45°C and in the presence of 0, 2, 4, 6, 8, and 10% NaCl (w/v). The production of gas from glucose was determined using inverted Durham tubes and carbohydrate fermentation profiles were determined using the API 50 CHL *Lactobacillus* identification system (bioMérieux). Furthermore, tests for the production of ammonia from arginine were performed as described previously (De Bruyne *et al.*, 2007). To test for the production of D- and L-lactate from glucose, cells were grown in MRS broth with glucose as the sole energy source. Proteins were removed by adding an equal volume of acetonitrile (Sigma-Aldrich, St. Louis, MO, USA); samples were subsequently microcentrifuged (14000 rpm for 8 min), filtered (0.2 µm, Sartorius AG, Göttingen, Germany) and transferred to an appropriate vessel prior to injection. The amount of D- and L-lactic acid in the supernatant was determined using HPLC with ultra violet (UV) detection. To this end, a Waters chromatograph (Waters Corporation, Milford, MA, USA) was used, equipped with a 486 UV detector, a 600S controller, a 717Plus autosampler, and a Shodex-Column Orpak CRX-853; 50 x 8 mm (Showa Denko KK, Tokyo, Japan). The mobile phase, at a flow rate of 1 ml min<sup>-1</sup>, consisted of a 1 mM CuSO<sub>4</sub> solution in ultrapure water with 20% acetonitrile (Sigma-Aldrich). D- and L-lactic acid were eluted isocratically and were detected by measuring the absorption at 253 nm. The results for strain LMG 27676<sup>T</sup> are given in the species description below. Characteristics that differentiate strain LMG 27676<sup>T</sup> from its nearest phylogenetic neighbors are summarized in Table 4. The biochemical characteristics of KCTC 2386<sup>T</sup> and LMG 23786 were identical with the following exceptions: only KCTC 2386<sup>T</sup> produces acid from L-arabinose; LMG 23786 produces acid from arbutin, which is not the case for KCTC 2386<sup>T</sup>.

Table 4: Differential phenotypic characteristics of strain LMG 27676<sup>T</sup> and closest related species of the genus *Leuconostoc*

Characteristic	1	2	3
Source	Rutabaga	Kimchi	Brown algae kimchi
Morphology	Ovoid	Coccus	Ovoid
NaCl (%) range for growth	0–6	0–7	0–6
Growth at 4°C	–	+	–
Growth at 37°C	–	+	+
Acid from:			
L-Arabinose	+	+	–
Arbutin	–	– <sup>a</sup>	+
Starch	–	–	+
D-xylose	–	–	+
DNA G+C content (mol%)	38	37	42.5

1: *L. rapi* sp. nov. LMG 27676<sup>T</sup>, 2: *L. kimchii* KCTC 2386<sup>TT</sup> and 3: *L. miyukkimchii* LMG 27965<sup>T</sup>.

<sup>a</sup>: This biochemical characteristic differs from what was previously reported (Kim *et al.*, 2000).

### 4.2.3 Conclusion

In conclusion, the results of the present study demonstrate that strain LMG 27676<sup>T</sup> represents a taxon that can be differentiated from the present *Leuconostoc* species by a range of genotypic methods and by several biochemical characteristics including the absence of growth at 37°C. Although the genotypic divergence in terms of DDH towards its nearest neighbor *L. kimchii* KCTC 2386<sup>T</sup> yielded a value that was close to the threshold level for species delineation, the *pheS*, *rpoA*, and *atpA* gene sequences clearly differentiated both taxa genotypically. We therefore classify the taxon represented by strain LMG 27676<sup>T</sup> as a novel *Leuconostoc* species, for which we propose the name *Leuconostoc rapi* sp. nov.

#### *Description of Leuconostoc rapi* sp. nov.

*Leuconostoc rapi* (ra'pi. L. gen. n. rapi of a turnip, referring to rutabaga, a turnip-like vegetable). Cells are Gram-stain-positive, catalase-negative, facultatively anaerobic, and non-motile. Cells are ovoid and occur singly, in pairs or in short chains. Colonies grown for 3 days on MRS agar at 28°C are approximately 1 mm in diameter, white/creamy, convex, and circular with smooth margins. Gas is produced from glucose, indicating the heterofermentative character of the type

strain. Produces L- and D-lactic acid in a ratio of 2:7. Growth occurs at 20–30°C and in the presence of 0–6% NaCl but not in the presence of 8–10% NaCl. Arginine is not hydrolyzed. Acid is produced from glucose, fructose, mannose, L-arabinose, D-ribose, mannitol, methyl- $\alpha$ -D-glucopyranoside, N-acetylglucosamine, amygdalin, esculin ferric citrate, salicin, D-cellobiose, sucrose, trehalose, gentiobiose, turanose, potassium gluconate, and potassium 2-ketogluconate. Acid is not produced from glycerol, erythritol, D-arabinose, D- and L-xylose, D-adonitol, methyl- $\beta$ -D-xylopyranoside, D-galactose, L-sorbose, L-rhamnose, dulcitol, inositol, D-sorbitol, methyl- $\alpha$ -D-mannopyranoside, arbutin, D-maltose, D-lactose, D-melibiose, inulin, D-melezitose, D-raffinose, starch, glycogen, xylitol, D-lyxose, D-tagatose, D- and L-fucose, and D- and L-arabitol. The DNA G+C content of the type strain is 38 mol% and its peptidoglycan type is A3 $\alpha$  L-Lys-Ala<sub>2</sub>. The type strain, LMG 27676<sup>T</sup> (= DSMZ 27776<sup>T</sup>) was isolated from a spoiled sous-vide cooked rutabaga produced by a Finnish manufacturer.

#### 4.2.4 Acknowledgements

The authors would like to thank Kauhajoki Food Laboratory and Pauliina Isohanni (Ruralia Institute, Seinäjoki Unit, University of Helsinki) for their technical assistance. Prof. Anu Hopia (Functional Foods Forum, University of Turku) and Prof. Per Saris (Department of Applied Chemistry and Microbiology, Faculty of Agriculture and Forestry, University of Helsinki) are acknowledged. The strain was isolated in a research project 'Novel food technologies - Special focus on quality and sustainable processing' that was financed by European Regional Development Fund (ERDF) via Tekes (The Finnish Funding Agency for Innovation), University Foundation of South Ostrobothnia and participating companies.

#### 4.2.5 Supplementary Material

The GenBank/EMBL/DDBJ accession numbers for the 16S rRNA, *pheS*, *rpoA*, and *atpA* gene of strain LMG 27676<sup>T</sup> are HG515542, HG515543, HG515544, and HG515545. The accession numbers for the *pheS*, *rpoA*, and *atpA* gene of strain LMG 27695<sup>T</sup> (*L. miyukkimchii*) and KCTC 2386<sup>T</sup> (*L. kimchii*) are HG515546–HG515548 and LN650690–LN650692, respectively.



### 4.3 *Carnobacterium iners* sp. nov., a Psychrophilic, Lactic Acid-Producing Bacterium from the Littoral Zone of an Antarctic Pond.

I. Snauwaert, B. Hoste, K. De Bruyne, K. Peeters, L. De Vuyst, A. Willems, P. Vandamme

Redrafted from: International Journal of Systematic and Evolutionary Microbiology, 2013 (63), 1370-75

Author contributions: IS performed the experiments and wrote the manuscript. BH and KDB designed the primers for the MLSA scheme construction. KP did the isolation and dereplication of the strains. PV and IS designed the experiment. AW, LDV, and PV proofread the manuscript.

Two lactic acid-producing, Gram-stain-positive rods were isolated from a microbial mat actively growing in the littoral zone of an Antarctic lake (Forlidas Pond) in the Pensacola mountains and studied using a polyphasic taxonomic approach. The isolates were examined by phylogenetic analysis of the 16S rRNA gene, MLSA of *pheS*, *rpoA*, and *atpA*, and biochemical and genotypic characteristics. One strain, designated LMG 26641, belonged to *Carnobacterium alterfunditum* and the other strain, designated LMG 26642<sup>T</sup>, could be assigned to a novel species, with *Carnobacterium funditum* DSM 5970<sup>T</sup> as its closest phylogenetic neighbor (99.2% 16S rRNA gene sequence similarity). *Carnobacterium iners* sp. nov. could be distinguished biochemically from other members of the genus *Carnobacterium* by the lack of acid production from most carbohydrates. DNA-DNA relatedness confirmed that strain LMG 26642<sup>T</sup> represented a novel species, for which we propose the name *Carnobacterium iners* sp. nov. (type strain is LMG 26642<sup>T</sup> = CCUG 62000<sup>T</sup>).

### 4.3.1 Introduction

The genus *Carnobacterium* was created to accommodate heterofermentative, facultatively anaerobic, psychrotolerant, rod-shaped LAB that produce L-lactic acid from glucose. At the time of writing, the genus *Carnobacterium* comprises 10 species with validly published names. Most of them were isolated from food of animal origin and/or living fish, except for *Carnobacterium alterfunditum*, *Carnobacterium funditum* and *Carnobacterium pleistocenium* which were isolated from cold environments with low nutrient contents, such as Antarctic ice lakes and permafrost ice (Franzmann *et al.*, 1991; Pikuta *et al.*, 2005).

Recently, Peeters *et al.* (2011) examined the heterotrophic bacterial diversity in a microbial mat sample originating from the littoral zone of a continental Antarctic lake (Forlidas Pond) in the Pensacola mountains. A large number of bacteria were isolated and characterized through rep-PCR fingerprinting and phylogenetic analysis of partial 16S rRNA gene sequences. Thirty LAB belonging to two main rep-clusters were isolated on marine agar 2216 (BD Difco) at 4–20°C in an aerobic or anaerobic atmosphere. Representatives of rep-cluster I (LMG 26642<sup>T</sup>, R-36994, R-37000, and LMG 26896) and rep-cluster II (LMG 26641 and LMG 26897) were grown aerobically on trypticase soy broth with yeast extract salt medium [TSBY salt; 3% trypticase soy broth (Oxoid), 0.3% yeast extract (Oxoid), 1.3% NaCl (Merck), 0.034% KCl (UCB), 0.4% MgCl<sub>2</sub>·6H<sub>2</sub>O (Sigma-Aldrich), 0.345% MgSO<sub>4</sub>·7H<sub>2</sub>O (UCB), 0.025% NH<sub>4</sub>Cl (Merck), 0.014% CaCl<sub>2</sub>·2H<sub>2</sub>O (Merck), 2% agar no. 2 (LabM), pH 7.2] at 4°C for 10 days and 20°C for 4 days, respectively. The clonality of these isolates was investigated by RAPD fingerprinting using primers RAPD-270 (5'-TGCGCGCGGG-3') and RAPD-272 (5'-AGCGGGCCAA-3'), as described by Mahenthiralingam *et al.* (1996). The fingerprints indicated that isolates LMG 26642<sup>T</sup>, LMG 26896, R-37000, and R-36994 were genetically identical, as was the case for LMG 26641 and LMG 26897 (data not shown).

### 4.3.2 Methods, Results, and Discussion

Nearly complete 16S rRNA gene sequence analysis was performed as described by Vancanneyt *et al.* (2004) for strains LMG 26642<sup>T</sup> and LMG 26641. PCR products were purified using a Nucleofast 96 PCR clean up membrane system (Macherey-Nagel, Germany). The sequencing primers were those listed by Coenye *et al.* (1999) and the obtained fragments were cleaned with a BigDye xTerminator Purification kit (Applied Biosystems, USA), according to the manufacturer's in-

structions. ARB (Ludwig *et al.*, 2004) and the corresponding SILVA SSURef 102 database (Pruesse *et al.*, 2007) were used to align the 16S rRNA gene sequences of strains LMG 26642<sup>T</sup> (1512 bp) and LMG 26641 (1433 bp) with sequences of type strains of all established species of the genus *Carnobacterium*. These aligned sequences were imported into MEGA version 5.0 (Tamura *et al.*, 2011) and analyzed using the neighbor-joining, ML and maximum-parsimony methods. The statistical reliability of tree topologies was evaluated by bootstrapping analysis based on 1000 replicates. The neighbor-joining and the maximum-parsimony trees revealed topologies similar to that obtained in the ML tree (Figure 12). 16S rRNA gene sequence similarities were calculated using ARB and the closest relatives of strain LMG 26642<sup>T</sup> were *C. funditum* DSM 5970<sup>T</sup> (99.2%) and *Carnobacterium mobile* DSM 4848<sup>T</sup> (97.0%). Strain LMG 26641 showed 100 and 99.8% 16S rRNA gene sequence similarity with *C. pleistocenium* FTR1<sup>T</sup> and *C. alterfunditum* ACAM 311, respectively. Lower sequence similarities (< 99.2%) were found with other members of the genus *Carnobacterium*.




Figure 12: ML tree based on 16S rRNA gene sequences showing the phylogenetic relationship of strains LMG 26642<sup>T</sup> and LMG 26641 with type strains of species of the genus *Carnobacterium*

Bootstrap values (> 50%) based on 1000 replications are shown at branch nodes. Bar, 0.5% sequence divergence. The substitution model used was the GTR model and the aligned sequence had a length of 1396 bp.

Because of the failure of 16S rRNA gene sequence similarity to reflect the phylogeny of closely related LAB, MLSA was performed to obtain a higher taxonomic

resolution. MLSA of the housekeeping genes encoding *pheS*, *rpoA*, and *atpA* has been successfully applied for the identification of different LAB (De Bruyne *et al.*, 2007; Naser *et al.*, 2005a). The primer combinations used for the amplification and sequencing of *pheS*, *rpoA*, and *atpA* are listed Table 5. Primer combinations *pheS*-24/25-F/*pheS*-26/27-R, *rpoA*-24-F/*rpoA*-25/26-R, and *atpA*-29-F/*atpA*-35-R amplified the target genes of all strains except for *atpA* of *C. alterfunditum*, *C. pleistocenium*, and *C. jeotgali*; for these, an alternative primer combination (*atpA*-30-F/*atpA*-33/34-R) was used. Amplification conditions and sequencing reactions were performed as described by Naser *et al.* (2005a) with the following modifications: (i) the annealing temperature was set at 50°C but in a few cases this value was raised to 55°C to avoid aspecific amplicons and (ii) sequencing reactions were purified using a BigDye xTerminator Purification kit. To assess the inter- and intraspecies variation, multiple strains per species were included if available; in total, 23 strains were studied. The strains examined and their depositors, sources and accession numbers are listed in Table 6. SeaView version 4 was used to concatenate the gene sequences of *pheS*, *rpoA*, and *atpA* (Gouy *et al.*, 2010). MEGA version 5.0 (Tamura *et al.*, 2011) was used to align the translated concatenated gene sequences and analyze the nucleotide sequences. The neighbor-joining and maximum-parsimony tree revealed topologies similar to that obtained in the ML tree (Figure 13). The phylogenetic tree of the concatenated *pheS*, *rpoA*, and *atpA* sequences proved the discriminatory power of these genes for species identification within the genus *Carnobacterium* and was roughly in agreement with the 16S rRNA gene-based analysis. All species were clearly delineated and had < 98% concatenated gene sequence similarity. Pairwise sequence similarities of the concatenated *pheS*, *rpoA*, and *atpA* sequences confirmed that the closest relatives of strain LMG 26642<sup>T</sup> were *C. funditum* LMG 14461<sup>T</sup> (93%) and *C. mobile* LMG 9842<sup>T</sup> (88.6%). MLSA also indicated that the closest relatives of LMG 26641 were *C. alterfunditum* LMG 14462<sup>T</sup> (98.1%) and *C. pleistocenium* LMG 23663<sup>T</sup> (93.8%). Lower sequence similarities (< 92.5%) were found towards the other strains tested. Based on MLSA using *pheS*, *rpoA*, and *atpA*, it can be concluded that strain LMG 26641 belongs to *C. alterfunditum* and that strain LMG 26642<sup>T</sup> can be assigned to a novel species, with *C. funditum* as its closest neighbor.

Table 5: Amplification and sequencing primers used for the MLSA of *Carnobacterium* strains

Primer name	Sequence (5'→3')	Position
<i>pheS</i> -24F	GAAGACTTGTTTATTGGTATGG	461
<i>pheS</i> -25F	CATCCAGCACGTGATATGC	557
<i>pheS</i> -26R	CCTAATCCAAAGGCAWATCC	1031
<i>pheS</i> -27R	GGATGAACVAYCCWGCRC	968
<i>rpoA</i> -24F	ATGATCGAAATTGAAAAACC	1
<i>rpoA</i> -25R	ACTTTAATCATTTCTGCTTC	844
<i>rpoA</i> -26R	ACTGAATTAATACCAGCACG	802
<i>atpA</i> -29F	TATRTCGGAGACGGGATTGC	97
<i>atpA</i> -30F	GCNCCWGGMGWTWATGSAWCG	397
<i>atpA</i> -33R	GCTTGTTTYGWTAAATCATC	781
<i>atpA</i> -34R	TTCATSGCTTTAATTTGWGC	1108
<i>atpA</i> -35R	CCACGATTTAATTTAGCTTG	1219




Figure 13: ML tree based on concatenated *pheS*, *rpoA*, and *atpA* sequences showing the phylogenetic relationship of strains LMG 26642<sup>T</sup> and LMG 26641 with the type strains of species of the genus *Carnobacterium*.

Bootstrap values (> 50%) based on 1000 replications are shown at branch nodes. Accession numbers are given in Table 6. Bar, 2% sequence divergence. The substitution model used was the GTR model and the aligned sequence had a length of 957 bp.

Table 6: Members of the genus *Carnobacterium* examined in this study

Strain	Depositor*	Source	Growth conditions	<i>pheS</i>	<i>rpoA</i>	<i>atpA</i>
<b><i>C. alterfunditum</i></b>						
LMG 14462 <sup>T</sup>	DSMZ	Anoxic water (Antarctica)	TSBY salt medium, 20°C, A	HE590697	HE590698	HE590699
LMG 26641	Own isolate	Microbial mat (Antarctica)	TSBY salt medium, 20°C, A	HE590759	HE590760	HE590761
<b><i>C. divergens</i></b>						
LMG 9199 <sup>T</sup>	NCIMB	Vacuum-packed refrigerated minced beef (South-Africa)	CBA, 37°C, MA	HE590700	HE590701	HE590702
LMG 22903	J. Leisner	Spoiled pork (Denmark)	CBA, 37°C, MA	HE590703	HE590704	HE590705
LMG 22900	J. Leisner	Beef (Denmark)	CBA, 37°C, MA	HE590706	HE590707	HE590708
R-22020	J. Leisner	Vacuum packed pork (ND)	CBA, 37°C, MA	HE590709	HE590710	HE590711
LMG 19743	J. Leisner	Modified-atmosphere-packaged shrimps (Denmark)	CBA, 37°C, MA	HE590712	HE590713	HE590714
<b><i>C. funditum</i></b>						
LMG 14461 <sup>T</sup>	DSMZ	Anoxic water (Antarctica)	TSBY salt medium, 20°C, A	HE590696	HE590715	HE590716
<b><i>C. gallinarum</i></b>						
LMG 9841 <sup>T</sup>	NCFB	Ice slush around chicken carcasses (ND)	MRS agar, 30°C, A	HE590717	HE590718	HE590719
<b><i>C. inhibens</i></b>						
LMG 23655 <sup>T</sup>	DSMZ	Gastrointestinal tract of an Atlantic salmon (Sweden)	TSA, 30°C, A	HE590720	HE590721	HE590722
<b><i>C. jeotgali</i></b>						
LMG 25668 <sup>T</sup>	JCM	Jeotgal (Korea)	TSA, 30°C, A	HE590723	HE590724	HE590725
<b><i>C. maltaromaticum</i></b>						
LMG 22899	J. Leisner	Modified-atmosphere packaged thawed cod (Denmark)	TSA, 30°C, A	HE590726	HE590727	HE590728
LMG 11393	DSMZ	Vacuum-packed beef (ND)	TSA, 30°C, A	HE590729	HE590730	HE590731
LMG 6903 <sup>T</sup>	NCIMB	Raw milk (ND)	TSA, 30°C, A	HE590732	HE590733	HE590734
LMG 9839	NCFB	Kidney of a diseased adult cutthroat trout (USA)	TSA, 30°C, A	HE590735	HE590736	HE590737
LMG 22902	J. Leisner	Modified atmosphere packaged beef (Denmark)	TSA, 30°C, A	HE590738	HE590739	HE590740
LMG 9840	NCFB	Refrigerated vacuum-packaged meat (ND)	TSA, 30°C, A	HE590741	HE590742	HE590743
<b><i>C. mobile</i></b>						
LMG 9842 <sup>T</sup>	NCFB	Irradiated chicken meat (ND)	TSA, 30°C, A	HE590744	HE590745	HE590746
LMG 21341	J. Leisner	Modified-atmosphere-packaged shrimps (Denmark)	TSA, 30°C, A	HE590747	HE590748	HE590749
LMG 21342	J. Leisner	Modified-atmosphere-packaged shrimps (Denmark)	TSA, 30°C, A	HE590750	HE590751	HE590752
<b><i>C. pleistocenium</i></b>						
LMG 23663 <sup>T</sup>	DSMZ	Ice core sample of a pleistocene thermokarst pond (Alaska)	CBA, 20°C, A	HE590753	HE590754	HE590755
<b><i>C. viridans</i></b>						
LMG 23657 <sup>T</sup>	DSMZ	Vacuum-packed ham (Canada)	TSA, 30°C, A	HE590756	HE590757	HE590758
<b><i>C. iners</i> sp. nov.</b>						
LMG 26642 <sup>T</sup>	Own isolate	Microbial mat (Antarctica)	TSBY salt medium, 4°C, A	HE578182	HE578183	HE578184

CBA, Columbia blood agar (Oxoid); TSA, tryptone soya agar (Oxoid); A, aerobic; mA, microaerobic, ND, no data available.  
 \* RIKEN BioResource Center, Japan; J. Leisner, Department of Veterinary Disease Biology, Faculty of Life Sciences, University of Copenhagen, Denmark.

To confirm the identification obtained by MLSA, DDH between strain LMG 26642<sup>T</sup> and its nearest phylogenetic neighbor, *C. funditum* LMG 14461<sup>T</sup>, was performed. Genomic DNA was extracted using the guanidinium thiocyanate method described by Pitcher *et al.* (1989). DDH was performed using the microplate method, with photobiotin for labeling the DNA (Ezaki *et al.*, 1989), as modified by Goris *et al.* (1998). DNA-DNA relatedness between strain LMG 26642<sup>T</sup> and *C. funditum* LMG 14461<sup>T</sup> was 18%, which confirmed that strain LMG 26642<sup>T</sup> represents a novel species.

The G+C content was determined according to the enzymatic DNA degradation method described by (1989), using a Waters Breeze HPLC system and XBridge Shield RP18 column. The solvent used was 0.02 M NH<sub>4</sub>H<sub>2</sub>PO<sub>4</sub> (pH 4.0)/1.5% (v/v) acetonitrile. Non-methylated lambda phage DNA (Sigma) was used as the calibration reference and *Escherichia coli* LMG 2093 DNA was included as the control. The G+C content of strain LMG 26642<sup>T</sup> was 34 mol%, which is within the range for the genus *Carnobacterium* (32–44 mol%). Strain LMG 26641 had a G+C content of 36 mol%, which corresponded with the G+C content of *C. alterfunditum* (32–36 mol%).

Preparation of the peptidoglycan and analysis of its structure were performed according to Schumann (2011). The diagnostic *meso*-diaminopimelic acid was found in addition to alanine and glutamic acid in the total hydrolysate (4 M HCl, 100°C, 15 h). The partial hydrolysate (4 M HCl, 100°C, 15 min) contained the peptides L-Ala-D-Glu and diaminopimelic acid-D-Ala. From these data, it was concluded that strain LMG 26642<sup>T</sup> contains the peptidoglycan type A1 $\gamma$ .

Physiological and biochemical tests were performed using strain LMG 26642<sup>T</sup>. Cell and colony morphology and the presence of catalase and oxidase was investigated after growth on TSBY salt agar for 10 days, incubated aerobically at 4°C. Standard biochemical tests and growth experiments were determined in TSBY salt broth and performed in triplicate, unless stated otherwise. Growth was tested at 4, 15, 20, and 37°C and in the presence of 2–10% (w/v) NaCl (at intervals of 2%). Production of gas from glucose was tested using inverted Durham tubes and peptone yeast extract as the basal medium (Holdeman & Moore, 1977). The production of D- and L-lactate from glucose was determined enzymatically (R-Biopharm). The API 50 CHL *Lactobacillus* identification system (bioMérieux) and GEN III Omilog ID system (Biolog) were used to determine the carbohydrate fermentation profile of the isolate and a positive control (*C. funditum* LMG 14461<sup>T</sup>). For both identification systems, strain LMG 26642<sup>T</sup> scored negative for all carbohydrates tested. The

same result was obtained when peptone yeast extract medium was used as the basal medium (Holdeman & Moore, 1977). The morphological and physiological characteristics of the isolate are given in the species description and an overview of the physiological differences between the isolate and its closest relatives is presented in Table 7.

*Table 7: Differential characteristics of strain LMG 26642<sup>T</sup> and members of the genus Carnobacterium that assimilate a limited range of carbohydrates*

Characteristic	1	2	3
<b>Acid from:</b>			
Galactose	-	w	-
Mannitol	-	+	+
Salicin	-	+	ND
Glycerol	-	w	-
D-Cellobiose	-	+	-
D-Fructose	-	+	+
Maltose	-	+	-
D- Mannose	-	+	-
D-Trehalose	-	+	-
D-Ribose	-	+	-
Sucrose	-	+	-
<b>Hydrolysis from aesculin</b>	+	w	+
<b>Lactic acid configuration</b>	L	L(+)	ND
<b>Growth in:</b>			
2% NaCl	+	ND*	+
4% NaCl	-	+	+
<b>DNA G+C content (mol%)</b>	34	32-34	43.9

Strains: 1, *C. iners* sp. nov. LMG 26642<sup>T</sup>; 2, *C. funditum* LMG 14461<sup>T</sup> (Franzmann *et al.*, 1991); 3, *C. jeotgali* LMG 25668<sup>T</sup> (Kim *et al.*, 2009). +, Positive; W, weakly positive; -, negative. \*

Sodium is required for growth with optimal growth at 1.7%



### 4.3.3 Conclusion

The colony and cell morphology of strain LMG 26642<sup>T</sup>, its motility, catalase activity, peptidoglycan type, and G+C content were in agreement with the description of the genus *Carnobacterium*. Additionally, strain LMG 26642<sup>T</sup> was closely related to *C. funditum* LMG 14461<sup>T</sup> (99.2% 16S rRNA gene sequence similarity) but it could be distinguished from this reference strain and other members of the genus *Carnobacterium* on the basis of MLSA and biochemical characteristics. Based on these results, we propose to classify strain LMG 26642<sup>T</sup> in a novel species, *Carnobacterium iners* sp. nov.

#### *Description of Carnobacterium iners* sp. nov.

*Carnobacterium iners* sp. nov. (in'ers. L. neut. adj. iners inactive, lazy). Cells are psychrophilic, Gram-stain-positive, catalase-negative, facultatively anaerobic, and motile rods, approximately 1.5  $\mu\text{m}$  wide and 3–6  $\mu\text{m}$  long, occurring singly or in pairs or short chains. Colonies grown for 10 days on TSBY salt agar at 4°C are approximately 0.8 mm in diameter, white, opaque, smooth and circular with undulate margins and an umbonate elevation. No gas is produced from glucose. Produces D- and L-isomers of lactic acid in a ratio of 1:9. Optimal growth is observed at 4°C; grows at 15 and 20°C. Grows with 1–2% NaCl, but not with 4–10% NaCl. Does not produce acid from glucose, fructose, mannose, N-acetylglucosamine, aesculin, cellobiose, maltose, trehalose, gentiobiose, glycerol, erythritol, D- or L-arabinose, ribose, D- or L-xylose, adonitol, methyl  $\beta$ -D-xylopyranoside, galactose, sorbose, rhamnose, dulcitol, inositol, mannitol, sorbitol, methyl  $\alpha$ -D-mannopyranoside, methyl  $\alpha$ -D-glucopyranoside, amygdalin, arbutin, salicin, lactose, melibiose, sucrose, inulin, melezitose, raffinose, starch, glycogen, xylitol, turanose, D-lyxose, D-tagatose, D- or L-fucose, D- or L-arabitol, potassium gluconate, potassium 2-ketogluconate, or potassium 5-ketogluconate. The cell wall contains *meso*-diaminopimelic acid.

The type strain, LMG 26642<sup>T</sup> (=CCUG 62000<sup>T</sup>), was isolated from a cyanobacterial mat growing in the littoral zone of a continental Antarctic lake (Forlidas Pond, Pensacola mountains) in December 2003. The DNA G+C content of the type strain is 34 mol%.

#### 4.3.4 Acknowledgements

We acknowledge financial support from the Research Foundation-Flanders and the Belgian Science Policy Office (BelSPO AMBIO project). We thank Dominic Hodgson and Peter Convey (British Antarctic Survey) who provided the environmental sample from which the isolate was obtained.

#### 4.3.5 Supplementary Material

The GenBank/EMBL/DDBJ accession numbers for the 16S rRNA gene, *pheS*, *rpoA*, and *atpA* sequences of strain LMG 26642<sup>T</sup> are HE583595 and HE578182-HE578184, respectively.

# 5

## Microbial Diversity Analyses in the HT Sequencing Era

This Chapter comprises the application of HT sequencing technologies to unravel the microbial diversity in an acidic beer ecosystem, along with a whole-genome sequence analysis of a LAB strain, isolated from this ecosystem. In Section 5.1, a HT sequencing approach, comprising targeted sequencing of the bacterial partial 16S rRNA gene and the fungal ITS region was applied to study the ecosystem of mature Belgian red-brown acidic ales. Belgian red-brown acidic ales are sour and alcoholic fermented beers, which are produced by mixed-culture fermentation. *Saccharomyces* dominates the ethanolic fermentation phase, after which lactobacilli produce lactic acid. *Dekkera* species, pediococci, and AAB prevail during the maturation phase, which takes place in oak barrels for about two years, after which the mature beer is blended with young, non-aged beer. The goal was to compare the microbial community diversity and metabolite composition of three Belgian red-brown acidic ales. Following, whole-genome sequencing of *Pediococcus damnosus* LMG 28219, which was the dominant LAB member of the Belgian red-brown acidic ale ecosystem, and comparative genome analysis were used in Section 5.2 to investigate this strains' mechanisms to reside in the beer niche.

## 5.1 Microbial Diversity and Metabolite Composition of Belgian Red-Brown Acidic Ales

I. Snauwaert, S. P. Roels, F. Van Nieuwerburg, A. Van Landschoot, L. De Vuyst, P. Vandamme

Submitted to: Applied and Environmental Microbiology, 2014

Author contributions: IS carried out all experiments and performed bioinformatics analyses. SPR performed the statistical analysis. FVN participated in the 454 pyrosequencing. AVL was involved in the collection of the samples. IS, LDV, and PV wrote the manuscript. All authors read and approved the final manuscript.

Belgian red-brown acidic ales, encompassing both red and red-brown acidic ales, are sour and alcoholic fermented beers, which are produced by mixed-culture fermentation. These beers are aged in oak barrels for about two years, after which mature beer is blended with young, non-aged beer to obtain the end-products. The microbial ecology of a Belgian red acidic ale has previously been characterized, setting the foundations of current knowledge on mixed-culture fermentations. The objective was to evaluate the microbial community diversity of Belgian red-brown acidic ales at the end of the maturation phase of three breweries and of three fermentation batches per brewery. The microbial diversity is compared with the metabolite composition at the end of the maturation phase. In this work, brew samples were subjected to 454 pyrosequencing of the 16S rRNA gene (bacteria) and the ITS region (yeasts) and a broad range of metabolites was quantified. The most important microbial species present in the Belgian red acidic ales were *Pediococcus damnosus*, *Dekkera bruxellensis*, and *Acetobacter pasteurianus*. In addition, culture-independent analysis revealed operational taxonomic units (OTUs) that were assigned to an unclassified fungal community member, *Candida*, and *Lactobacillus*. The main metabolites were L-lactic acid, D-lactic acid, and ethanol, whereas acetic acid was produced in lower quantities. The most prevailing aroma compounds were ethyl acetate, isoamyl acetate, ethyl hexanoate, and ethyl octanoate, which might be of impact on the aroma of the end-products.

### 5.1.1 Introduction

Beer production is generally a monoculture fermentation process, with the yeast *Saccharomyces cerevisiae* as the most important player. However, the fermentation and/or maturation of some Belgian acidic beers is determined by autochthonous nonstarter microorganisms or uncharacterized mixed starter cultures (Verachtert & Iserentant, 1995; Verachtert & Derdelinckx, 2005). These beers represent culturally important alcoholic beverages, for which the microbial activities play critical roles not only in their production but also regarding quality characteristics. Indeed, the end-products of these fermentations have a unique taste, for which they are gaining popularity worldwide.

Well-known Belgian acidic beers that are the result of the activities of an autochthonous microbiota are the lambic beers, which are characterized by a spontaneous fermentation process involving enterobacteria (the first phase), *S. cerevisiae*, and *Saccharomyces pastorianus* (the main fermentation phase), and *Dekkera bruxellensis* and *Pediococcus damnosus* (the maturation phase) (Verachtert & Iserentant, 1995; Verachtert & Derdelinckx, 2005; Spitaels *et al.*, 2014b). Other examples are the ACAs, for which production methods similar to those of lambic beers are used (Bokulich *et al.*, 2012a). The maturation phase of the production of these beers is dominated by the yeast *D. bruxellensis*, which produces a range of flavor compounds, including octanoic acid and decanoic acid and their respective ethyl esters (Bokulich & Bamforth, 2013). LAB are the second most prominent group of microorganisms in these beers, whose activities result in increased lactic acid concentrations.

Much less studied however are the Belgian red-brown acidic ales, which are produced by a mixed-culture fermentation (Martens *et al.*, 1997). Nowadays, these beers are produced by only a few breweries in West- and East- Flanders (Belgium) that produce these beers for years and by some American craft breweries that aim to produce a comparable style. Two types of Belgian red-brown acidic ales exist, namely the red acidic ales of South-West-Flanders and the red-brown acidic ales that are produced in South-West- and South-East-Flanders.

Martens and colleagues monitored the microbial communities during the industrial production of a Belgian red acidic ale and described a tandem fermentation process (1997). *Saccharomyces* dominates the ethanolic fermentation phase, after which lactobacilli start the production of lactic acid. *Dekkera* species, pediococci, and AAB prevail during the maturation phase, which takes place in oak barrels.

After two years of aging, the pH drops to 3.2–3.5 and lactic acid and acetic acid concentrations increase to 6000 mg/L and 1600 mg/L, respectively, are reached (Martens *et al.*, 1997). Finally, the end-product is obtained by blending the sour mature beer and young beer that did not undergo aging in oak vessels.

The current study attempted to update and extend the existing data on Belgian red-brown acidic ales by reanalyzing their microbial diversity and metabolite composition at the end of the maturation phase. In this study the microbial diversity and metabolite composition of Belgian red-brown acidic ales are compared.

## 5.1.2 Materials and Methods

### The Experimental Material

Three subsequent brews (samples referred to as A, B, and C) of three Belgian red-brown acidic ale breweries (referred to as no. 1, 2, and 3) were sampled at the end of their maturation phase. Red acidic ales and red-brown acidic ales were produced by brewery 1 and breweries 2 and 3, respectively. All microorganisms present were collected through microcentrifugation (8000 rpm, 20 min, 4°C) and the cell pellets were washed with resuspension buffer (0.15 M NaCl, 0.1 M EDTA, pH 8.0). The cell pellets and supernatants were stored separately at –20°C until further analysis. In addition, samples of the bottled beers produced from these brews were stored in triplicate at –20°C.

### Microbial Diversity Analysis of the Brew Samples

#### *Total DNA Extraction*

Total bacterial and fungal DNA was extracted using the procedure described by Gevers and colleagues (2001), with several modifications, as follows. (i) The thawed pellet was washed in 1 ml TES buffer (6.7% sucrose, 50 mM Tris-HCl, pH 8.0, 1 mM EDTA) and resuspended in 275 µL STET buffer (8.0% sucrose, 5.0% Triton X-100, 50 mM Tris-HCl, pH 8.0, 50 mM EDTA). (ii) Ninety µL lysozyme-mutanolysin-proteinase K solution (TES buffer containing 1667 U/mL mutanolysin, 33 mg/mL lysozyme, and 2.78 mg/mL proteinase K) was added and the suspension was incubated at 37°C for 1 h. (iii) Prior to extraction with a phenol/chloroform/isoamyl

alcohol (49.5:49.5:1.0) solution (Sigma-Aldrich, St. Louis, MO, USA), proteins were precipitated by adding 250  $\mu$ L ammonium acetate. (iv) Five  $\mu$ L RNase (2 mg/L) was added to the DNA solution, which was incubated at 37°C for 60 min and subsequently stored at -20°C. The integrity, concentration, and purity of the DNA isolated was evaluated using 1% (wt/vol) agarose gels, stained in ethidium bromide, and by spectrophotometric measurements at 234, 260, and 280 nm. Per brew sample, DNA extracts were prepared in triplicate and pooled. Pooled DNA solutions were diluted to a concentration of 50 ng/ $\mu$ L prior to amplification.

#### *Multiplex 454 Pyrosequencing of a Bacterial and Fungal Marker*

Amplification was performed using a two-step PCR procedure that increases the reproducibility and consistently recovers a higher genetic diversity in pyrosequencing libraries (Berry *et al.*, 2011). To analyze the bacterial communities, a 16S rRNA gene fragment encompassing the variable regions V1 and V3 was amplified using the forward primer pA 5'-AGAGTTTGATCCTGGCTCAG-3' (Sigma-Aldrich) and the reverse primer 518R 5'-ATTACCGCGGCTGCTGG-3' (Sigma-Aldrich). The thermal program used for the amplification of the V1-3 region of the 16S rRNA gene started with a denaturation at 95°C for 5 min, followed by 20 cycles of amplification (45 s at 95°C, 45 s at 55°C, and 30 s at 72°C), and a final elongation at 72°C for 8 min. The fungal ITS region was amplified using the forward primer ITS1F (5'-CTTGGTCATTTAGAGGAAGTAA-3'; Sigma-Aldrich) (Gardes & Bruns, 1993) and the reverse primer ITS4 (5'-CCTCCGCTTATTGATATGC-3'; Sigma-Aldrich) (White *et al.*, 1990). The following thermal program was applied for fungal DNA amplification: denaturation at 95°C for 5 min, 3 cycles of pre-amplification (1 min at 95°C, 2 min 15 s at 55°C, and 1 min 15 s at 72°C) followed by 30 amplification cycles (1 min at 95°C, 1 min 15 s at 55°C, and 1 min 15 s at 72°C), and elongation at 72°C for 7 min. All PCR assays were conducted in triplicate, using the following reagents: 1.25 U/reaction FastStart High Fidelity Enzyme blend (Roche Diagnostics, Basel, Switzerland), GeneAmp® dNTP's (200  $\mu$ M of each; Life Technologies, Carlsbad, CA, USA), 0.4  $\mu$ M of the forward and reverse primers, and FastStart High Fidelity Reaction buffer containing 1.8 mM MgCl<sub>2</sub> (Roche Diagnostics). The Veriti® 96-Well Fast Thermal Cycler (Life Technologies) was used for all amplification reactions. The PCR products were evaluated using 1% (wt/vol) agarose gels, stained in ethidium bromide, and they were pooled per brew sample afterwards. Subsequently, sequencing constructs (primers of the second PCR step) of the GS FLX Titanium chemistry were added to the amplicons, using the thermal program and PCR mixture applied before, with the following modifications. (i) The number of cycles for the pre-amplification and amplification runs was reduced to one

and five, respectively. (ii) The sequencing constructs were obtained from Integrated DNA technologies (IDT, Leuven, Belgium), as recommended by Roche Diagnostics, and their concentrations in the PCR mixture were reduced to 0.1  $\mu$ M. The constructs are compatible with Lib-L chemistry and consisted of a Titanium adaptor, an eight base pair (bp) multiplex identifier (MID), and a template-specific PCR primer. The PCR products were purified using a MinElute PCR Purification kit (Qiagen, Venlo, The Netherlands) and checked using the 2100 BioAnalyzer (Agilent Technologies, Santa Clara, CA, USA) with the DNA 7500 kit (Agilent Technologies). A fluorescent stain base kit (Qubit®dsDNA Broad range assay kit; Life Technologies) was used for DNA quantification. Subsequently, the PCR products of different samples were pooled in equimolar concentrations and sequenced unidirectionally on a GS FLX instrument (half run), using Titanium chemistry (Roche Diagnostics), at Beckman Coulter Genomics SA (Grenoble, France). The 16S rRNA gene amplicons were sequenced in the reverse direction, whereas the fungal ITS amplicons were sequenced in the forward direction. A mock community containing both bacterial as well as fungal DNA and a generous donor sample (*i.e.*, a sample that was sequenced before) were included in the analysis to enable error analysis.

### *Processing of the Sequence Data*

The raw .sff 454 pyrosequencing file was processed using a 16S rRNA gene pipeline constructed by Schloss and coworkers ([http://www.mothur.org/wiki/454\\_SOP](http://www.mothur.org/wiki/454_SOP)) (2011a), with several modifications, as follows. (i) No nucleotide mismatches to the sample barcode and adjacent PCR primer were allowed for sample multiplex barcode deconvolution and primer trimming. (ii) Sequences with an ambiguous base call or a homopolymer stretch longer than six nucleotides were removed from subsequent analysis. (iii) The Mothur implementation of the SeqNoise algorithm was used to further reduce the overall error rate (Quince *et al.*, 2011). Additional modifications were needed for the analysis of the fungal ITS region. (i) The full ITS1 of the nuclear ITS region was extracted using the V-Xtractor package (Hartmann *et al.*, 2010) and fungal HMMs developed by Nilsson and coworkers (2010). (ii) Sequences were aligned to each other using the needleman alignment method (a gap was only penalized once, terminal gaps were penalized, and all distances were calculated), and uncorrected pairwise distances between sequences were calculated. (iii) An in-house developed training set, containing ITS sequences retrieved from the YeastIP database (Weiss *et al.*, 2013), was used for taxonomic assignments. All identifications were reviewed by means of the Mycobank identification tool (Crous *et al.*, 2004).



The bacterial and fungal datasets were normalized (normalized value = relative abundance \* norm; norm = the number of sequences in the smallest group) and binned into OTUs with a cut-off of 97%, and rarefaction analyses were performed. The community diversity was measured using the inverse Simpson diversity index (1/D). Furthermore, differences in community structure and membership were not only explored using the Bray-Curtis and Jaccard index, but also using weighted and unweighted UniFrac (Lozupone *et al.*, 2007). These differences were quantified by means of analysis of similarity (ANOSIM) and visualized using principal coordinate analysis (PCoA). The R Studio software package (version 0.97.551) was used for heatmap construction and PCoA.

### *Isolation of Dominant Community Members from Brew Samples*

A mature brew sample from brewery 1 was serially diluted in 0.86% (wt/vol) saline and 50  $\mu$ L of each dilution was plated on different agar media in triplicate, followed by incubation at 28°C in an aerobic atmosphere for 4-6 days. The dominant LAB and AAB community members were isolated using MRS (Oxoid) and mDMS (Papalexandratou *et al.*, 2013) agars, respectively, supplemented with 5 ppm amphotericin B (Sigma-Aldrich) and 200 ppm cycloheximide (Sigma-Aldrich). DYP A [2% glucose, 0.5% yeast extract, 1% peptone, 100 ppm chloramphenicol (Sigma-Aldrich), and 1.5% agar (wt/vol)] was used as a general yeast agar isolation medium or was supplemented with 50 ppm cycloheximide (DYPAX) to favor the slow-growing *Dekkera* yeast (Abbott *et al.*, 2005; Suárez *et al.*, 2007). Isolates were subcultured twice and subjected to MALDI-TOF MS for dereplication, as described previously (Ghyselinck *et al.*, 2011). As sequence analysis of protein-encoding genes has proven to be superior for accurate species level identification of a variety of LAB and AAB (Cleenwerck *et al.*, 2010; De Bruyne *et al.*, 2007; Naser *et al.*, 2005a; Snauwaert *et al.*, 2013a), representative bacterial isolates were identified through sequence analysis of the *pheS* gene or the molecular chaperone (*groEL*) gene, for *Pediococcus* and *Acetobacter*, respectively, in addition to sequencing of the 16S rRNA gene. Representative fungal isolates were identified by sequencing of the ITS region and D1/D2 region of the 26S rRNA gene (Kurtzman & Robnett, 1998).

## Metabolite Target Analysis of the Brew Samples and Bottled Beers

### *Determination of Carbohydrate Concentrations*

High-performance anion exchange chromatography (HPAEC) with pulsed amperometric detection was used to determine the concentrations of glucose, fructose, sucrose, and mannitol. An ICS 3000 chromatograph (Dionex, Sunnyvale, CA, USA) with a CarboPac<sup>TM</sup> PA10 column (Dionex) was used for this purpose. The mobile phase, at a flow rate of 1.0 ml/min, consisted of ultrapure water (eluent A), 167 mM NaOH (eluent B), and 500 mM NaOH (eluent C), with the following gradient: 0.0 min, 87% A and 13% B; 20.0 min, 87% A and 13% B; 25 min, 57% A, 13% B, and 30% C; 26 min, 100% C; 30 min, 100% C; 31 min, 87% A and 13% B; and 35 min, 87% A and 13% B (Janssens *et al.*, 2012). To remove proteins, 150  $\mu$ L of Carrez A reagent [3.6% (wt/vol)  $\text{K}_4\text{Fe}(\text{CN})_6 \cdot 3\text{H}_2\text{O}$ ] and Carrez B reagent [7.2% (wt/vol)  $\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$ ] were added to 300  $\mu$ L of the tenfold diluted samples. HPLC with an evaporative light-scattering detector (ELSD) was used for the determination of the concentrations of maltose, maltotriose, maltotetraose, maltopentaose, and maltohexaose. To this end, a Waters chromatograph (Waters Corporation, Milford, MA, USA) was used, equipped with a 2424 ELS Detector (gain, 100; drift tube, 65°C; gas pressure, 40 psi; nebulizer cooling), a 600S controller, a 717Plus autosampler, and a Grace Prevail Carbohydrate ES column (250 x 4.6 mm, 5  $\mu$ m, 35°C; Grace Discovery Sciences, Deerfield, IL, USA). The mobile phase, at a flow rate of 1.0 ml/min, consisted of ultrapure water (eluent A) and acetonitrile (eluent B), with the following gradient: 0.0 min, 25% A and 75% B; 50.0 min, 40% A and 60% B; 51 min, 25% A and 75% B; and 60 min, 25% A and 75% B. Proteins were removed by adding 500  $\mu$ L acetonitrile (Sigma-Aldrich) to 500  $\mu$ L sample. All samples were microcentrifuged (14000 rpm for 8 min), filtered (0.2  $\mu$ m filters, Sartorius AG, Göttingen, Germany), and transferred to an appropriate vessel prior to injection. Calibration was performed using external standards and samples were analyzed in triplicate.

### *Determination of the Concentrations of D- and L-Lactic Acid*

Concentrations of D- and L-lactic acid were determined using HPLC-UV. To this end, a Waters chromatograph (Waters Corporation) was used, equipped with a 486 UV detector, a 600S controller, a 717Plus autosampler, and a Shodex-Column Orpak CRX-853 (50 x 8 mm; Showa Denko KK, Tokyo, Japan). The mobile phase, at a flow rate of 1.0 ml/min, consisted of a 1 mM  $\text{CuSO}_4$  solution in ultrapure water with 20% acetonitrile (Sigma-Aldrich). D- and L-lactic acid were eluted isocratically. Detec-

tion was performed by measuring the absorption at 253 nm. Sample preparation, calibration, and number of technical replicates were as described above.

### *Determination of Volatiles*

Static-headspace (SH)-GC-MS was used for the identification and quantification of flavor compounds. Samples were prepared in headspace vials containing 5 ml sample as well as a known amount of 1-butanol (Sigma-Aldrich) as internal standard. The same procedures and instrument control parameters were applied as reported by Ravyts and colleagues (2009), except that sample equilibration was done by agitation at 60°C for 30 min prior to injection into the column. Using this method, isobutyl alcohol, isoamyl acetate, isoamyl alcohol, ethyl hexanoate, ethyl octanoate, ethyl decanoate, 2-phenylethyl acetate, and 2-phenyl ethanol could be successfully identified. Ethyl acetate and 1-propanol could be analyzed successfully when splitting the injection port. The amount of volatiles present in the samples was calculated using external standards and all samples were analyzed in triplicate.

### *Determination of the Concentrations of Ethanol and Short-Chain Fatty Acids*

The concentrations of ethanol and short-chain fatty acids (SCFAs) in the samples were measured by gas chromatography, using a Focus gas chromatograph (Interscience, Breda, The Netherlands) equipped with a Stabilwax-DA column (Restek, Bellefonte, PA, USA), a flame ionization detector, and an AS 3000 autosampler. Hydrogen gas was used as carrier gas with a constant flow rate of 1.0 ml/min; nitrogen gas was used as make-up gas. The injector and detector temperatures were set to 240°C and 250°C, respectively, and the following temperature program was used: 0.0 min at 40°C, 10.0 min at 140°C, 12.0 min at 230°C, and 22.0 min at 230°C. Nine hundred  $\mu\text{L}$  of a mixture [acetonitrile (Sigma-Aldrich), 1% (vol/vol) formic acid (Merck, Darmstadt, Germany), and 2.842 g/L 2-butanol (Merck)] was added to a 300  $\mu\text{L}$  non-diluted or ten times diluted sample for the measurement of SCFAs and ethanol, respectively. Prior to injection, the sample was microcentrifuged (14000 rpm for 8 min) and filtered (0.2  $\mu\text{m}$ , Sartorius AG). A volume of 1.0  $\mu\text{L}$  was injected using a split ratio of 40:1. The quantification was done in triplicate using external standards, as described above.

## Statistical Analysis

Missing data were excluded prior to statistical analysis of the metabolite data. A correlation matrix was calculated using the Pearson correlation coefficient. Principal component analysis (PCA) was performed to visualize possible relationships within the data matrix. Statistical analysis was performed using the R Studio software package (version 0.97.551).

A multivariate approach to unravel possible interactions between the metabolite concentrations and the microbial communities was performed, as described previously (Wang *et al.*, 2012), with several modifications. Outlying brew samples (2B and 2C) and non-informative variables (metabolites and microorganisms absent in more than 3 out of 7 samples) were excluded prior to analysis. Canonical correlation analysis (CCA) was used to examine possible correlations between individual external variables and overall microbial communities plus correlations between all external variables and each individual OTU. CCA was performed using the *cca* function in the Vegan package (R Studio software package, version 0.97.551); the *envfit* function was used to calculate the p-value of each correlation between each OTU and all variables. For this dataset, the cut-off for significance was set at  $p < 0.15$ . Variables and community members without any correlations were excluded from the multivariate approach. Afterwards, the selected variables were subjected to a multiple linear model (function *lm* and ANOVA). For the small dataset, the statistical significance was set at a p-value  $< 0.05$ . Finally, the direction and size of the correlations between variables and OTUs were derived from a linear regression model.

### 5.1.3 Results

#### Analysis of the 454 Pyrosequencing Data of the Community DNA from the Brew Samples

Sequencing of the amplified fragments yielded a total of 544,508 reads with a mean length of 507 bp, which corresponded roughly with 20,000 reads per sample. Multiplex barcode deconvolution, primer trimming, and denoising reduced the total number of bacterial and fungal sequences to 199,481 (395 unique sequences, mean read length of 245 bp) and 232,921 (475 unique sequences, mean read length of 178 bp), respectively. Mock community analysis showed that singleton OTUs represented sequencing errors and these OTUs were therefore excluded for further

analysis. After quality control, the normalized number of bacterial and fungal reads per sample ranged from 9,781 to 9,796 [mean  $\pm$  standard error (SE) of  $9792 \pm 1.5$ ] and 13,339 to 13,360 ( $13,355 \pm 2.3$ ), respectively. A total of 45 and 60 bacterial and fungal OTUs, respectively, remained, with a few OTUs representing a large fraction of all reads (see below). Rarefaction analysis of the bacterial and fungal datasets (Supplementary Material 5.1) indicated that all samples were more or less reaching the plateau phase and hence no additional sequencing was performed.

Bacterial communities predominantly consisted of the divisions *Firmicutes* (representing 81.64% of all reads, 21 OTUs) and *Proteobacteria* (18.31%, 15 OTUs). Only a minor fraction of the reads belonged to *Actinobacteria* (0.02%, 4 OTUs), *Deinococcus-Thermus* (0.01%, 1 OTU), and *Bacteroidetes* (0.02%, 4 OTUs). All *Firmicutes* reads were assigned to the order *Lactobacillales*, with the exception of 4 *Bacillales* OTUs (0.02%). The *Lactobacillales* OTUs belonged to the following genera: *Pediococcus* (74.49%, 1 OTU), *Lactobacillus* (7.66%, 8 OTUs), *Leuconostoc* (0.22%, 1 OTU), *Weissella* (0.23%, 1 OTU), and *Lactococcus* (0.01%, 2 OTUs). A major fraction of the *Proteobacteria* reads were assigned to the order *Rhodospirales* (18.22%, 1 OTU), more specifically to a member of the family *Acetobacteraceae*. *Dekkera* was the dominant genus in the fungal dataset (59.26%, 2 OTUs), with smaller fractions of *Pichia* and *Candida* representing 7.93% (5 OTUs) and 5.72% (4 OTUs) of the fungal reads, respectively. Only one OTU was assigned to *Kregervanrija*, *Debaryomyces*, *Priceomyces*, *Hyphopichia*, and *Wickerhamomyces*, consisting of 0.24, 0.08, 0.07, 0.03, and less than 0.01% of the reads, respectively. Additionally, 26.45% (40 OTUs) of the fungal sequences remained unclassified.

### Comparison of the Microbial Communities of the Brew Samples

The highest median bacterial and fungal 1/D indices were found in the brew samples of breweries 2 and 3, respectively, whereas the lowest were found in the brew samples of breweries 3 and 1, respectively. Additional distributional information can be found in the boxplots, as depicted in Figure 14. Differences in community structure were explored and visualized by means of PCoA, based on the Bray-Curtis and weighted UniFrac indices (Supplementary Material 5.2). The PCoA ordination of the bacterial dataset based on the Bray-Curtis index showed a separation of brew samples 2B and 2C from the remaining samples along the first axis. This trend was confirmed (but less pronounced) when incorporating phylogenetic distances using weighted UniFrac indices. PCoA based on the Bray-Curtis index of the fungal dataset showed that brew sample 2C and brew samples

3A and 3C were separated from the remaining samples along the first axis, whereas brew samples 2A and 2B and brew sample 3A were separated from the remaining ones along the second axis. Differences in community structure were quantified by means of ANOSIM. The community structure varied across brew samples of different breweries, exerting a significant impact on the bacterial (Bray-Curtis  $R_{\text{ANOSIM}} = 0.506$ ,  $p < 0.005$ ) but not on the fungal (Bray-Curtis  $R_{\text{ANOSIM}} = 0.095$ ,  $p > 0.005$ ) genetic diversity. Incorporating phylogenetic distances did not reveal significant differences in community structure across brew samples of different breweries. In addition, differences in community membership were explored using PCoA, based on the Jaccard and unweighted UniFrac indices (Supplementary Material 5.3). PCoA of the bacterial dataset based on the Jaccard and unweighted UniFrac indices both revealed a separation of brew samples 2A and 2B from the remaining brew samples along the first axis, whereas no separation was apparent along the second axis. PCoA of the fungal dataset based on the Jaccard and unweighted UniFrac index showed a separation of the three brew samples from brewery 2 from those from breweries 1 and 3 along the first axis. The community membership varied across brew samples of different breweries, exerting a significant impact on the bacterial (Jaccard  $R_{\text{ANOSIM}} = 0.091$ ,  $P > 0.005$ ) and the fungal (Jaccard  $R_{\text{ANOSIM}} = 0.062$ ,  $P > 0.005$ ) genetic diversity. Incorporating phylogenetic distances did not reveal significant differences in community membership across brew samples of different breweries.




Figure 14:  $1/D$  of the bacterial (A) and fungal (B) datasets of brew samples from breweries 1, 2, and 3

The relative abundances of the most abundant bacterial and fungal OTUs were visualized by means of a heatmap, in which the brew samples were grouped based on their Bray–Curtis distances (Figure 15 and Supplementary Material 5.4). Brew samples A, B, and C from breweries 1 and 3 comprised rather stable bacterial communities, whereas more variation was apparent in the bacterial communities of brew samples from brewery 2 (Figure 15A). The bacterial communities of all brew samples were dominated by *Pediococcus* (bacterial OTU 1), except for brew samples 2B and 2C, in which an *Acetobacteraceae* OTU (bacterial OTU 2) was predominant. Several OTUs that were assigned to the genus *Lactobacillus* represented smaller fractions. Those were OTU 5, being present in all brew samples except for brew sample 2A, and OTU 3 and OTU 4, accounting for 33.44% and 7.18% of the reads in brew sample 2B, respectively, but being less abundant in the remaining brew samples. *Weissella* (bacterial OTU 7) and *Leuconostoc* (bacterial OTU 8) were both found in brew sample 2B, whereas brew sample 3C only contained a minor fraction of OTU 7 (*Weissella*). Brew samples A, B, and C from brewery 1 comprised rather stable fungal communities, whereas more variation was apparent in the fungal communities of brew samples from breweries 2 and 3 (Figure 15B). Fungal OTU 1, which was assigned to the genus *Dekkera*, was the predominant member in the fungal communities of all brew samples, except for brew sample 2C and brew samples 3A and 3C, which were dominated by an unclassified OTU (fungal OTU 3). Two fungal OTUs were assigned to the genus *Pichia* (OTUs 4 and 5) and *Candida* (OTUs 3 and 7), respectively, whereas one OTU (OTU 10) was assigned to the genus *Kregervanrija*.




Figure 15: Relative abundances of the most abundant OTUs ( $> 0.01\%$  of the reads) of the bacterial (A) and fungal (B) communities

The OTUs and sample names (breweries 1, 2, or 3; samples A, B, or C) are shown on the x- and y-axis, respectively. The color gradient ranges from black to white with decreasing relative abundances. The Bray-Curtis distances between the brew samples were calculated and an average-linkage clustering was used for the construction of the dendrograms.



### Analysis of the Community Diversity Shared by the Brew Samples of the Different Breweries and Isolation of Dominant Community Members


The bacterial community members that were present in brew samples of each of the three breweries were OTU 1 (*Pediococcus*), OTU 2 (*Acetobacteraceae* unclassified), OTU 3 (*Lactobacillus*), OTU 4 (*Lactobacillus*), and OTU 5 (*Lactobacillus*). Similarly, fungal OTU 1 (*Dekkera*), OTU 2 (unclassified), OTU 4 (*Pichia*), and OTU 5 (*Pichia*) were found in all three breweries. To confirm these results and to obtain a more accurate identification of this microbiota, a targeted isolation of each of the dominant community members was conducted on a brew sample from brewery 1. Dereplication analysis of 78, 39, and 85 isolates from aerobically incubated MRS, mDMS, and DYPAX agars, respectively, yielded for these media a single MALDI-TOF MS profile (data not shown). Firstly, Sanger sequencing of the 16S rRNA and *pheS* genes assigned the taxon isolated from MRS agar to *Pediococcus damnosus*; the 251 nucleotide long stretch of the 16S rRNA gene sequence, which corresponded with the 454 pyrosequence fragment randomly picked up from bacterial OTU 1 (*Pediococcus*), shared 99.37% of its sequence. Secondly, the taxon isolated from mDMS agar was identified as *Acetobacter pasteurianus* based on 16S rRNA and *rpoB* gene sequencing; the 226 nucleotide long stretch of 16S rRNA gene sequence, which corresponded with the 454 pyrosequence fragment randomly picked up from bacterial OTU 2 (*Acetobacteraceae*), shared 100% of its sequence. Finally, Sanger sequencing of the ITS and D1/D2 regions assigned the taxon isolated from DYPAX to *Dekkera bruxellensis*; the 175 nucleotide long stretch of the ITS sequence, which corresponded with the 454 pyrosequence fragment randomly picked up from fungal OTU 1 (*Dekkera*), shared 97.74% of its sequence. No growth was obtained on DYPA medium.

### Comparison of the Metabolite Data of the Brew Samples

The sucrose, fructose, and mannitol concentrations were  $5.68 \pm 1.79$  mg/L,  $154.47 \pm 6.83$  mg/L, and  $2.27 \pm 0.59$  mg/L, respectively. None of the brew samples contained glucose. The concentrations of the remaining carbohydrates were  $540.62 \pm 48.50$  mg/L maltose,  $877.72 \pm 72.43$  mg/L maltotriose,  $1760.99 \pm 69.56$  mg/L maltotetraose,  $1342.56 \pm 41.59$  mg/L maltopentaose, and  $1545.37 \pm 42.55$  mg/L maltohexaose. The average D- and L-lactic acid concentrations at the end of the maturation phase of all brew samples were  $2497.35 \pm 145.40$  mg/L (mean  $\pm$  SE) and  $2175.47 \pm 125.99$  mg/L, respectively (details on the spread of the data are presented in Supplementary Material 5.5). The average acetic acid concentration of the whole dataset was  $1376.78 \pm 75.84$  mg/L and only a limited number of brew

samples contained propionic acid and isobutyric acid ( $3.70 \pm 0.86$  mg/L and  $9.95 \pm 1.75$  mg/L, respectively). The average concentrations of ethyl acetate, isobutyl alcohol, isoamyl acetate, isoamyl alcohol, and ethyl hexanoate were  $151.07 \pm 0.61$  mg/L,  $14.69 \pm 0.61$  mg/L,  $0.88 \pm 0.08$  mg/L,  $77.89 \pm 2.92$  mg/L, and  $9.43 \pm 1.22$  mg/L, respectively. No 2-phenylethyl acetate, 2-phenyl ethanol, and only minor fractions of ethyl octanoate were found. An average ethanol concentration of  $6.46 \pm 0.14\%$  (vol/vol) was measured.

To explore and visualize differences in metabolite composition among all brew samples analyzed, a PCA was performed (Figure 16A). The data points from brew samples of breweries 1 and 3 showed overlap along both axes, whereas those from brewery 2 grouped separately. Along axis 2, there was a subdivision of the data points from brew samples of brewery 2 into two clusters. In addition, the metabolite data from brew samples of breweries 1, 2, and 3 were compared by means of a heatmap (Figure 16B). The median L-lactic acid concentration was the highest in brew samples from brewery 3 and the lowest in brew samples from brewery 2, whereas the median acetic acid concentration was the highest in brewery 1 and the lowest in brewery 3. Brew samples from breweries 1 and 3 contained higher median D-lactic acid concentrations compared to those from brewery 2. The fructose concentrations were comparable in brew samples of the three breweries. The median maltose concentrations in brew samples from breweries 1 and 2 were similar, whereas the median maltose concentration in beer samples from brewery 3 was lower. Brew samples from brewery 1 contained the highest median maltotriose, maltotetraose, and maltopentaose concentrations, whereas brew samples from brewery 2 had the lowest median maltotetraose and maltopentaose concentrations. The lowest median maltotriose concentration was found in brew samples from brewery 3. No clear differences in median maltohexaose concentrations could be found. Sucrose was only found in small amounts in brew sample 3B ( $51.12 \pm 0.62$  mg/L). The median ethyl acetate concentrations were higher in brew samples from breweries 1 and 2 compared to the median ethyl acetate concentration in brewery 3. No differences in the median concentrations of isoamyl alcohol among the brew samples of the different breweries were apparent. Brew samples from brewery 3 contained less isoamyl acetate compared to the quantities found in brew samples from breweries 1 and 2. The median isobutyl alcohol and ethyl hexanoate concentrations were the highest in brew samples from brewery 2, whereas lower concentrations were found in brew samples from breweries 1 and 3. No apparent differences were found in the median concentrations of mannitol, propionic acid, and isobutyric acid.



*Figure 16: PCA plot (A) and heatmap (B) of the metabolite concentrations of breweries 1, 2, and 3, determined in brew samples from the end of the maturation phase*

(A) In the PCA plot, the positions of the samples along the two first principal component axes are illustrated, along with the percentages of variation explained by each axis. A total of 81 data points were analyzed, including both biological and technical replicates.

Figure 16 is continued on the next page

B	mg/l																					v/v %
	L-lactic acid	D-lactic acid	ethyl acetate	isobutyl alcohol	isoamyl acetate	isoamyl alcohol	ethyl hexanoate	ethyl octanoate	mannitol	fructose	sucrose	maltose	maltotriose	maltotetraose	maltopentaose	maltohexaose	acetic acid	propionic acid	isobutyric acid	ethanol		
Brewery 1	A	2745.03	2436.39	153.88	9.88	1.17	60.95	6.04	0.06	0.00	145.64	0.00	662.53	1841.10	2513.59	1663.24	1472.07	1958.90	0.00	4.55	6.00	
	B	2419.26	2447.36	161.60	12.16	1.40	68.95	4.48	0.06	0.00	172.85	0.00	688.80	1593.63	2129.07	1682.59	1275.25	2062.68	0.00	17.96	5.00	
	C	2135.65	2270.74	258.21	9.88	1.14	93.16	0.00	0.06	0.00	60.08	0.00	866.34	1132.26	1356.63	1524.47	1886.33	2345.45	11.17	3.28	6.00	
Brewery 2	A	2393.79	1508.18	146.37	21.74	1.29	86.18	36.40	0.08	4.81	199.27	0.00	825.27	724.63	1243.98	902.53	1015.00	1723.17	12.09	43.99	7.00	
	B	711.78	876.90	25.73	20.70	0.29	69.44	18.98	0.06	0.00	160.34	0.00	736.65	988.93	1179.66	826.26	1482.70	306.31	0.00	0.00	7.00	
	C	1119.75	830.33	275.34	22.99	1.70	98.31	2.93	0.06	0.00	82.97	0.00	0.00	1109.93	1284.67	1207.45	1801.76	1308.60	0.32	0.00	7.00	
Brewery 3	A	5029.78	2329.56	68.96	8.76	0.00	57.56	0.00	0.00	14.69	262.08	0.00	1264.30	713.04	2916.97	1384.45	1328.79	1055.88	0.00	0.00	7.00	
	B	2356.33	2166.11	82.33	12.27	0.58	59.25	0.00	0.06	0.00	170.35	0.00	0.00	0.00	1432.26	1781.04	2254.31	1047.57	0.00	13.93	7.00	
	C	4489.14	5307.34	86.60	11.76	0.00	98.45	15.15	0.06	0.00	137.07	50.87	0.00	0.00	1665.40	1020.29	1244.38	493.17	2.78	0.00	4.00	

(B) The median concentrations of all compounds are shown and colored with a gradient that ranges from black (high median concentration) to white (low median concentration). All concentrations are expressed in mg/L, except for ethanol (in vol/vol%)


### Correlations Between Microbial Community Members and Metabolite Compositions

A CCA on 20 metabolites and 37 OTUs reduced the whole dataset to 6 metabolites and 9 OTUs. In total, 16 significant correlations between individual metabolites and OTUs were found through a multivariate analysis, namely 8 positive and 8 negative correlations. The fold change of the significant correlations at OTU level is depicted in Supplementary Material 5.6. The dominant *Pediococcus* (bacterial OTU 1) and *Dekkera* (fungal OTU 1) members were both negatively correlated with L-lactic acid, whereas *Pediococcus* was also negatively correlated with acetic acid but positively correlated with isobutyl alcohol. Bacterial OTU 2 (*Acetobacteraceae*) and OTU 5 (*Lactobacillus*) were negatively correlated with L-lactic acid and isoamyl acetate, respectively, whereas a positive correlation was found with isobutyl alcohol. *Pichia* OTU 4 showed a negative correlation with maltopentaose and a positive correlation with isobutyl alcohol and isobutyric acid, whereas *Pichia* OTU 5 was positively correlated with L-lactic acid. The unclassified fungal OTU 2 was negatively correlated with acetic acid, whereas OTU 29 was negatively correlated with isobutyric acid and maltopentaose. Finally, fungal OTU 24 (*Dekkera*) was positively correlated with isoamyl acetate.

### Metabolite compositions of the bottled beers

The glucose, fructose, and sucrose concentrations of the bottled beers were  $155.74 \pm 31.17$  mg/L,  $2612.15 \pm 227.79$  mg/L, and  $16.34 \pm 2.77$  mg/L, respectively. No mannitol could be found. The quantities of the remaining carbohydrates were  $989.38 \pm 74.99$  mg/L maltose,  $1481.94 \pm 90.31$  mg/L maltotriose,  $3468.06 \pm 96.50$  mg/L maltotetraose,  $1609.91 \pm 64.40$  mg/L maltopentaose, and  $1628.55 \pm 58.89$  mg/L maltohexaose. The average L- and D-lactic acid concentrations of the bottled beers were  $871.42 \pm 48.69$  mg/L (mean  $\pm$  SE) and  $560.57 \pm 44.06$  mg/L, respectively (details on the spread of the data points are presented in Supplementary Material 5.7). The average acetic acid concentration of the whole dataset was  $673.66 \pm 46.30$  mg/L and only a limited number of samples from the bottled beers contained propionic acid ( $8.40 \pm 1.88$  mg/L). No isobutyric acid could be found. The average concentrations and SEs of ethyl acetate, isobutyl alcohol, isoamyl acetate, isoamyl alcohol, and ethyl hexanoate were  $70.69 \pm 4.15$  mg/L,  $13.64 \pm 0.59$  mg/L,  $4.62 \pm 0.28$  mg/L,  $78.04 \pm 4.31$  mg/L, and  $0.69 \pm 0.11$  mg/L, respectively. No 2-phenylethyl acetate, 2-phenyl ethanol, and ethyl octanoate could be found. An average ethanol concentration of  $4.78 \pm 0.23\%$  (vol/vol) was measured.

To explore and visualize differences in metabolite compositions, a PCA and cluster analysis were performed (Figure 17A). The data points from bottled beers from breweries 1 and 3 showed overlap along both axes, whereas those from brewery 2 showed no overlap. The metabolite compositions of the bottled beers from breweries 1, 2, and 3 were compared by means of a heatmap (Figure 17B). The median concentrations of L-lactic acid and acetic acid were the highest in bottled beers from brewery 2, whereas the median concentrations of D-lactic acid were the lowest in bottled beers from brewery 2. The highest median glucose concentration was found in bottled beers from brewery 1. High fructose concentrations were found in bottled beers from breweries 1 and 3 compared to those from brewery 2. Sucrose was only found in bottled beers from breweries 1 and 3, with average concentrations of  $47.31 \pm 11.08$  mg/L and  $47.14 \pm 0.79$  mg/L, respectively. No large differences were found in the median maltose concentrations across the bottled beers of the different breweries. Brewery 3 had the highest median maltotriose concentration but the lowest median maltotetraose, -pentaose, and -hexaose concentrations. The median maltotriose, -tetraose, -pentaose, and -hexaose concentrations in bottled beers from brewery 2 were slightly higher compared to those from brewery 1. The median ethyl acetate and isoamyl alcohol concentrations in bottled beers from brewery 3 were lower compared to those from breweries 1 and 2. Bottled beers from brewery 2 contained a higher median isobutyl alcohol concentration compared to those from breweries 1 and 3. No apparent differences were found in the median concentrations of isoamyl acetate, ethyl hexanoate, and propionic acid.



*Figure 17: PCA plot (A) and heatmap (B) of the metabolite concentrations of the bottled beers from breweries 1, 2, and 3*

(A) In the PCA plot, the positions of the samples along the two first principal component axes are illustrated, along with the percentages of variation explained by each axis. A total of 81 data points were analyzed, including both biological and technical replicates.

Figure 17 is continued on the next page

B	mg/l																			v/v %
	L-lactic acid	D-lactic acid	ethyl acetate	isobutyl alcohol	isoamyl acetate	isoamyl alcohol	ethyl hexanoate	ethyl octanoate	glucose	fructose	sucrose	maltose		maltotriose	maltotetraose	maltohexaose	acetic acid	propionic acid	ethanol	
												maltopentaoase	maltotriose							
A	550.41	434.93	44.96	10.07	3.06	64.75	2.11	0.05	784.81	5852.54	0.00	1307.60	962.09	4399.48	1739.36	1749.03	379.83	0.00	5.00	
Brewery 1	B	368.99	398.48	81.52	10.05	1.88	88.73	0.00	0.00	601.11	5012.52	0.00	0.00	0.00	3081.70	1471.25	1643.19	553.63	51.57	5.00
	C	841.76	1047.53	108.32	10.58	3.14	149.84	0.00	0.00	0.00	3449.37	36.12	1173.84	1920.94	3692.03	1806.75	1840.93	608.01	0.00	2.00
Brewery 2	A	740.34	262.42	89.14	19.50	5.99	74.58	1.92	0.08	0.00	183.55	0.00	1356.19	1522.33	4362.03	2161.87	1887.32	1386.97	0.00	7.00
	B	1412.01	347.24	79.39	16.06	5.10	56.65	2.11	0.00	0.00	128.21	0.00	1168.93	1437.35	4098.63	2166.53	1860.15	1375.82	0.00	6.00
Brewery 3	C	982.38	169.96	57.00	21.29	3.78	86.92	0.00	0.00	156.18	538.42	0.00	0.00	0.00	3849.02	2223.10	2288.50	742.49	31.78	7.00
	A	713.33	485.43	38.21	8.05	3.93	31.27	0.00	0.00	0.00	4637.44	0.00	1096.63	2348.85	2651.69	607.27	614.16	485.67	0.00	5.00
Brewery 3	B	783.24	1020.20	55.18	14.11	6.34	74.74	0.00	0.00	0.00	2278.57	47.00	2037.26	2108.86	2645.38	1295.17	1565.89	295.04	0.00	2.00
	C	1165.87	1417.25	38.32	12.10	4.95	61.02	0.00	0.00	0.00	2402.66	48.58	564.03	1670.10	2087.77	937.20	1050.42	205.82	3.43	2.00

(B) The median concentrations of all compounds are shown and colored with a gradient that ranges from black (high median concentration) to white (low median concentration). All concentrations are expressed in mg/L, except for ethanol (in vol/vol%)



### 5.1.4 Discussion

The present study reanalyzed the microbial species diversity and metabolite composition of a Belgian red acidic ale previously characterized by traditional, culture-dependent techniques along with limited chemical analyses (Martens *et al.*, 1997), in comparison with two other Belgian red-brown acidic ales. Each of these beers was produced through mixed-culture fermentation and maturation in oak barrels (Martens *et al.*, 1997).

Viewed at higher taxonomic levels (*i.e.*, judging from partial 16S rRNA gene and ITS1 region sequence analysis), the microbial diversity of the brew samples of brewery 1 (*i.e.*, red acidic ale) was similar to that reported by Martens and coworkers (1997). Indeed, *Dekkera* and *Pediococcus* predominated during red acidic ale maturation, with *Dekkera* being mainly responsible for super-attenuation of the beer, *i.e.*, the depletion of oligo- and polysaccharides, and *Pediococcus* for lactic acid production. This coexistence of yeasts and LAB at the end of the maturation phase is a known phenomenon during gueuze beer fermentation (Spitaels *et al.*, 2014b; Van Oevelen *et al.*, 1977; Kumara *et al.*, 1993). Also, previously reported mixed culture experiments indicate that super-attenuation by *Dekkera* is less pronounced when pediococci are absent and suggest that growing cells of *Dekkera* can hydrolyse the EPSs produced by pediococci (Kumara & Verachtert, 1991; Van Oevelen & Verachtert, 1979). Besides these dominant members, the present study revealed the presence of an *Acetobacteraceae* member, an unclassified fungus, *Candida* and *Lactobacillus* in the brew samples of this brewery. The presence of AAB agrees with the findings of Martens and coworkers (1997), who isolated a minor fraction of these bacteria and probably reflects the presence of oxygenated niches in the vertical maturation casks. As the unclassified fungus was most abundant in one brew sample only, further research is needed to reveal its identity and function. The presence of *Candida*, which is known to be capable of growing in beer and to occur at the air-beer interface (van der Aa Kuhle & Jespersen, 1998; Timke *et al.*, 2008), contrasted with the previously reported findings for Belgian red acidic ales (Martens *et al.*, 1997), in which *Candida* species could not be recovered at the end of the maturation phase. Finally, the red acidic ale microbiota of this brewery also comprised a *Lactobacillus*. Indeed, several lactobacilli reside in beer (Bokulich & Bamforth, 2013) and were recovered during primary and secondary fermentation of the previously studied red acidic ale, but never from the mature product (Martens *et al.*, 1997).

At present, HT sequencing cannot reliably differentiate many closely related bacterial species because of the limited read length of HT sequencing technologies. Hence, HT sequencing cannot be considered fully comprehensive if detailed species level dynamics must be revealed, which is common practice in the study of food fermentations (Bokulich & Mills, 2012d). Therefore, the targeted isolation of microorganisms of interest, which were previously revealed by HT sequencing, is an appealing strategy for subsequent in-depth analysis of the prevailing microbiota at the species level. The community members, isolated from the mature brew samples of brewery 1, were identified as *P. damnosus*, *A. pasteurianus* and *D. bruxellensis*, and their SSU rRNA gene sequences showed high similarities (> 97%) towards the pyrosequences of the respective most abundant OTUs. This species composition seems different from that reported by Martens and coworkers (1997), who mainly isolated *P. parvulus*, *D. lambicus*, *D. bruxellensis*, and some AAB (which were not identified further) at the end of the maturation phase.

*P. damnosus* is however phylogenetically closely related to *P. parvulus* and this species shares a lot of phenotypic characteristics with the latter (Holzapfel *et al.*, 2006). Martens and coworkers (1997) identified *P. parvulus* based on phenotypic and biochemical characteristics only, distinguishing *P. parvulus* from *P. damnosus* because of the inability of the latter to grow at 35°C. Yet, both species have been associated with the brewery environment or with environments in which alcoholic beverages are produced, with *P. damnosus* being the most frequently encountered in brewery environments (Sakamoto & Konings, 2003). Most likely, the fairly high concentrations of L-lactic acid and D-lactic acid found in the brew samples of the present study were associated with the presence of *P. damnosus*, which is able to metabolize glucose homofermentatively to both L- and D-lactic acid, and which creates a desired tartness. The total concentrations of lactic acid in the bottled beers measured in the present study were lower compared to those reported by Martens and coworkers (1997); these researchers also found a higher concentration of D-lactic acid compared to the concentration of L-lactic acid in the brew samples and bottled beers. Similarly to *P. damnosus*, *P. parvulus* produces both L-lactic acid and D-lactic acid but, to our knowledge, there is no information on the ratio of L-lactic acid to D-lactic acid produced by these species in the beer environment. At present, the reason for the observed difference remains unclear, although it is not excluded that the higher L-lactic acid concentration originates from acidification in the brewing process.

*A. pasteurianus* was isolated from a mature brew sample, which agrees with the occurrence of AAB such as *A. aceti*, *A. pasteurianus*, *A. lambicus*, and *Gluconobacter oxydans* (Bokulich & Bamforth, 2013; Spitaels *et al.*, 2014a) at the air-beer inter-

face and possibly in the porous wooden walls of the barrels (Bokulich *et al.*, 2012a; Martens *et al.*, 1997), and which are responsible for a change in beer flavor through the oxidation of ethanol into acetate. The high acetic acid concentrations detected in the present study are likely associated with the action of *A. pasteurianus*. Martens and coworkers (1997) found similar acetic acid concentrations in the mature brew samples, but they reported higher concentrations in the bottled beers compared to the findings of the present study. However, the blending protocol of this brewery has changed over time from a strongly acidic beer to a less sour alcoholic beverage (less lactic acid and acetic acid), following consumers' preferences.

Furthermore, taxonomic studies revealed that *D. lambicus* is a synonym of *D. bruxellensis* (Smith *et al.*, 1990), which confirms that *D. bruxellensis* is the dominant fungal member of this Belgian red acidic ale. *Dekkera bruxellensis* outcompetes *S. cerevisiae* in environments with low carbohydrate concentrations (Tiukova *et al.*, 2013). Metabolite target analysis showed that glucose was completely depleted at the end of the maturation phase, whereas substantial amounts of maltose and several oligosaccharides (maltotriose, -tetraose, -pentaose, and -hexaose) were still present. This contrasted with the results obtained for bottled beers, where the glucose and fructose concentrations were much higher and indicated that the end-product was sweetened by blending and possibly carbohydrate addition to comply with the consumers' preferences. *Saccharomyces cerevisiae* produces, along with ethanol and carbon dioxide, low molecular-mass flavor compounds, such as higher alcohols, aldehydes, ketones, organic acids, and esters that soften the sour taste and add fruity notes to the beers (Verstrepen *et al.*, 2003) as well as organic sulfides (Lodolo *et al.*, 2008). In the case of *D. bruxellensis*, it is known that this yeast is indispensable for the typical lambic beer flavor and that it produces several metabolites that are characteristic for lambic beers (Verachtert & Iserentant, 1995; Shanta Kumara & Verachtert, 1991; Spaepen *et al.*, 1978,9; Spaepen & Verachtert, 1982; Van Oevelen *et al.*, 1976). Metabolite target analysis of brew samples at the end of the maturation phase showed the presence of isoamyl alcohol and isoamyl acetate, which contrasted with the absence of 2-phenyl ethanol and 2-phenylethyl acetate. These metabolites were also found in the bottled beers. Propionic acid and isobutyric acid levels were low, as was described previously by Van Oevelen and colleagues (1976). The ethyl acetate (*i.e.*, solvent or fruity flavor) levels found in the brew samples at the end of the maturation phase and in the bottled beers of the present study were higher compared to what has been measured previously (Martens *et al.*, 1997). Whereas small concentrations of ethyl hexanoate and ethyl octanoate were found during the present study, no ethyl decanoate (*i.e.*, the typical ester present in lambic and gueuze beers (Van Oevelen *et al.*, 1976)) could be found. Next to the acetate esters mentioned above, these ethyl esters are also

flavor-active esters (Pires *et al.*, 2014), contributing to the taste of Belgian red acidic ales and gueuze beers (Spaepen & Verachtert, 1982).

Finally, no *Lactobacillus*, *Candida*, or other fungal species could be isolated from the mature brew samples in the present study, which was in agreement with previous findings (Martens *et al.*, 1997). Possible reasons are that these microorganisms are not able to grow under the experimental cultivation conditions, that they are in a viable but not cultivable state, or that they are no longer viable. Together, the findings on the microbial diversity and metabolite composition of Belgian red acid ales, obtained through 454 pyrosequencing and metabolite target analysis, show many similarities with those previously generated by Martens and coworkers (1997) with classic approaches.

The microbial diversity and metabolite composition of the brew samples from brewery 3 (which produced a red-brown acidic ale) resembled those of brewery 1 (which produced a red acidic ale), while brew samples from brewery 2 (which also produced a red-brown acidic ale) were very different. The microbial communities in brew samples from brewery 3 not only contained *Pediococcus* and *Dekkera*, but also an unclassified fungus and *Lactobacillus* (albeit in a lower concentration). In addition, lower relative abundances of the *Acetobacteraceae* and the *Candida* community members were also found in brew samples from brewery 3, which further contrasted with the findings in brew samples from brewery 1. The higher L-lactic acid and D-lactic acid and the lower acetic acid concentrations in brew samples from brewery 3 compared to those from brewery 1 could be attributed to these differences in community structures. It is not clear if these differences in relative abundances reflect the real structure of the microbial communities of the brew samples, because those abundances based on cultivation and molecular PCR-based identification could be partially attributed to amplification biases (Engelbrektson *et al.*, 2010). Additionally, several fungal OTUs remained unclassified in the brew samples from brewery 3. Different reference databases yielded different taxonomic assignments for these OTUs as a function of completeness and quality, underlining the need for a comprehensive, carefully annotated, fungal reference database (Begerow *et al.*, 2010; Seifert, 2009). Finally, both the microbial diversities and metabolite compositions of the three subsequent brews of brewery 2 were different from each other and from those of breweries 1 and 3. The high lactic acid and acetic acid concentrations of brew sample 2A in combination with a low number of AAB indicated that another community member, such as *Dekkera*, must have been responsible for the acetic acid production. The detection of minor fractions of the reads that were assigned to the genera *Weissella* and *Leuconostoc* in brew

sample 2B is remarkable. These LAB have not yet been isolated from beer, but were recently detected in a barcoded amplicon sequencing study of ACA (Bokulich *et al.*, 2012a).

The outcome of the multivariate analysis partially contrasted with what was known from literature. For instance, the presence of the *Pediococcus* community member was negatively correlated with L-lactic acid, whereas a positive correlation with isobutyl alcohol was found. This probably reflects the usage of lactate by other community members during fermentation. Furthermore, the *Acetobacteraceae* community member was not correlated with acetic acid, besides its positive correlation with isobutyl alcohol and negative correlation with L-lactic acid, contrasting with the metabolic features of this community member. Possibly, the current dataset is too small to unravel valid links between the microbiota and the metabolite compositions of mature Belgian red-brown acidic ales or that community members involved during fermentation and maturation collectively impact the flavor of these beers. Furthermore, the microbiota detected in the finished fermentations and beers might not necessarily represent the active populations during fermentation.

In addition to the high-abundance OTUs discussed above, numerous low-abundance OTUs were discovered during this study. The diverse taxonomical assignments included *Actinobacteria*, *Deinococcus-Thermus*, *Bacteroidetes*, *Beta-* and *Gamma-proteobacteria*, *Kregervanrija*, *Debaryomyces*, *Priceomyces*, *Hyphopichia*, *Wickerhamomyces*, and others. Most of these microorganisms have not been reported in the brewery environment. A similar finding was seen in a study applying barcoded amplicon sequencing to unravel the microbiota of ACA (Bokulich *et al.*, 2012a). One possible explanation is that extensive sequencing of low-diversity microbial communities leads to the accumulation of errors that still remain present after stringent denoising (Dickie, 2010; Huse *et al.*, 2007; Kunin *et al.*, 2010). This leads to a long tail of low-abundant taxa in the species abundance curve, which are likely to be sequencing errors, and stresses the importance of focusing on the most abundant OTUs during amplicon sequencing studies that are performed without extensive biological and technical replicates. Another explanation may be that the physiological status of these microbes prevents their cultivation on the growth media used routinely. In that scenario, the present molecular community profiling method would be the first step to reveal these low-abundant microorganisms involved in Belgian red-brown acidic ale fermentations.

In conclusion, the microbial community diversity and metabolite composition of three Belgian red-brown acidic ales at the end of their maturation phase was


analyzed during the present study. In two of the brew samples of these breweries, the maturation phase was mainly dominated by *Pediococcus* and *Dekkera*. The microbial communities in brewery 2 samples were highly variable and differed from those of breweries 1 and 3. The main metabolites produced during these mixed-culture fermentations were ethanol, L- and D-lactic acid, and acetic acid. Ethyl acetate, isoamyl acetate, ethyl hexanoate, and ethyl octanoate dominated the flavor of the Belgian red-brown acidic ales of the present study.

### 5.1.5 Acknowledgments

The authors acknowledge the financial support of the FWO-Flanders, the Research Council of Ghent University (BOF project) and the Vrije Universiteit Brussel (SRP, IRP, and IOF projects), and the Hercules Foundation. In addition, the authors would like to acknowledge the breweries involved, the BCCM/LMG bacteria collection for providing DNA for MOCK community construction, Charlotte Peeters and Bart Verheyde for bioinformatics support, Leilei Li for sequencing the *groEL* housekeeping gene, and Tom Balzarini for the metabolite target analyses.


### 5.1.6 Supplementary Material

The raw sequence data received from Beckman Coulter were deposited at the Sequence Read Archive (SRA) of the National Center for Biotechnology Information (NCBI) (accession number: SRP035529). The GenBank/EMBL accession number for the Sanger sequences generated during this study are LK024185-LK024190.



Supplementary Material 5.1: Rarefaction analysis of the normalized bacterial (A) and fungal (B) datasets of the three consecutive brew samples (A, B, and C) from breweries 1, 2, and 3


Number of 3% OTUs: i.e., the richness of the OTUs at the 3% level.



*Supplementary Material 5.2: PCoA plots of bacterial (A) and fungal (B) communities from brew samples that were subjected to 454 pyrosequencing*

The positions of the bacterial and fungal communities for each species along the two first principal coordinate axes are illustrated, along with the percentage of variation explained by each axis. The results are based on the Bray–Curtis and weighted UniFrac distances, as indicated on the graphs.






*Supplementary Material 5.3: PCoA plots of bacterial (A) and fungal (B) communities from brew samples that were subjected to 454 pyrosequencing*

The positions of the bacterial and fungal communities for each species along the two first principal coordinate axes are illustrated, along with the percentage of variation explained by each axis. The results are based on the Jaccard and unweighted UniFrac distances, as indicated on the graphs.

		Brewery 1			Brewery 2			Brewery 3		
		A	B	C	A	B	C	A	B	C
BACTERIAL DATASET	OTU 1 <i>Pediococcus</i>	86.51	88.23	83.87	99.44	1.87	23.27	90.27	95.11	92.77
	OTU 2 <i>Acetobacteraceae</i>	11.88	7.96	14.25	0.46	52.15	76.71	0.13	0.11	0.35
	OTU 3 <i>Lactobacillus</i>	0.01	0.01			33.44		3.31	0.04	0.09
	OTU 4 <i>Lactobacillus</i>	0.01	0.01			7.18		0.67	0.04	0.04
	OTU 5 <i>Lactobacillus</i>	1.45	3.71	1.32		0.01	0.02	5.37	4.66	6.70
	OTU 7 <i>Weissella</i>					2.02				0.01
	OTU 8 <i>Leuconostoc</i>					2.02				
FUNGAL DATASET	OTU 1 <i>Dekkera</i>	61.60	85.30	94.08	51.79	74.45	14.02	32.98	82.24	28.21
	OTU 2 unclassified	20.95	1.96	0.64	0.14	0.01	85.07	36.41	16.39	60.25
	OTU 3 <i>Candida</i>	15.00	11.00	4.96				0.01		9.45
	OTU 4 <i>Pichia</i>	0.01			37.45			0.31	0.08	0.03
	OTU 5 <i>Pichia</i>		0.10	0.19	9.93	11.40	0.02	10.42	0.89	0.28
	OTU 6 unclassified	0.01						6.88	0.26	
	OTU 7 <i>Candida</i>					11.08	0.01			
	OTU 8 unclassified							3.92		
	OTU 9 unclassified				0.01			3.19	0.10	
	OTU 10 <i>Kregervanrija</i>		1.03							1.15
	OTU 12 unclassified							1.78		
	OTU 13 unclassified							1.47		
	OTU 14 unclassified	1.98					0.01			
	OTU 15 unclassified							1.44		


Supplementary Material 5.4: Relative abundances (%) of the most abundant (> 0.01% of the reads) bacterial and fungal OTUs used for the construction of the heatmap in Figure 15

Relative abundances of 0.00% are highlighted in grey.




Supplementary Material 5.5: Boxplot of each of the metabolites measured in brew samples from the end of the maturation phase, in which the data points were grouped according to the brewery they originated from

The quantities are expressed in mg/L for all metabolites, except for ethanol [in (vol/vol)%]



*Supplementary Material 5.6: Link between the microbial diversity and metabolite target analysis data set in brew samples from the end of the maturation phase*

Highly positive and highly negative correlations are indicated in green and red, respectively. Correlations closer to zero are indicated in orange.



Supplementary Material 5.7: Boxplot of the metabolite concentrations in the bottled beers, in which the data points were grouped according to the brewery the beers originated from

The quantities are expressed in mg/L for all compounds, except for ethanol in (vol/vol)%.

### 5.1.7 Data Analysis Pipeline

The data analysis pipeline used is based on a paper of Schloss and coworkers ([http://www.mothur.org/wiki/454\\_SOP](http://www.mothur.org/wiki/454_SOP)) (2011a), with several modifications. Schloss and coworkers (2011a) constructed the pipeline based on the analysis of a mock community. The data analysis pipeline involves several steps, including (i) identifying and removing low quality sequences, (ii) trimming of sequences, (iii) denoising flowgrams, (iv) denoising sequences, (v) identifying and removing chimera's, etc.

#### *Identifying and Removing Low-quality Sequences*

Several features are linked with sequences of low quality. Therefore, identifying and removing sequences that contain these features reduces the overall error rate. Examples of features that are linked with low quality reads are the following: sequences with mismatches to the barcode of primer region, sequences that contain ambiguous base calls, sequences that are shorter than 200 bp, sequences that contain homopolymers, sequences that aligned to the incorrect region within the 16S rRNA gene... Similar parameters as applied by Schloss and coworkers (2011a) were used, with the exception of the number of mismatches to the barcode and primer region (which was set to zero) and the number of homopolymers (which was set to 6), for which more stringent values were chosen.

#### *Trimming of Sequences*

The association between error rates and quality scores, as shown by Schloss and coworkers (2011a), indicated that trimming sequences to a break point, where quality score criteria were no longer met, results in reducing the overall error rates. Schloss and coworkers (2011a) proved that a 50 bp sliding window with an average quality score of 35 within the window was the most efficient. Exactly the same approach was used in the present study. Furthermore, the sequences were trimmed to a region where all sequences started and ended in the same alignment positions. This trimming insured that evolutionary consistent regions were being compared. For this purpose, our data was aligned to the SILVA-compatible template alignment, containing 50000 columns. The general approach was (i) to find the closest template for each candidate using kmer (k=8) searching, (ii) to make a pairwise alignment between the candidate and de-gapped template sequences using the Needleman-Wunsch algorithm with a reward of +1 for a match and penalties of -1 and -2

for a mismatch and gap, respectively and (iii) to re-insert gaps to the candidate and template pairwise alignments using the NAST algorithm so that the candidate sequence alignment is compatible with the original template alignment. The aligner does not explicitly take into account the secondary structure of the 16S rRNA gene, but the SILVA-compatible template alignment is based on the secondary structure, resulting in an alignment that will at least be implicitly based on the secondary structure.

### *Denoising Flowgrams*

The Pyronoise algorithm (Quince *et al.*, 2011) was applied, which reduces the sequencing error rate by correcting the original flowgram data using an expectation-maximization algorithm. This algorithm was re-implemented in the mothur software package to take advantage of accelerated clustering algorithms and to make the algorithm more accessible to other researchers. Flowgrams were trimmed to 450 flows because of the reasons explained by Schloss and coworkers (2011a).

### *Denoising Sequences*

Lingering PCR amplification and sequencing errors were removed using the SeqNoise algorithm. This algorithm uses the sequences as inputs and a model that describes rates of substitutions and homopolymeric insertions and deletions.

### *Identifying and Removing Chimera's*

Schloss and coworkers explored several chimera removal strategies, *e.g.*, Chimera Slayer, Uchime, and Perseus. They showed that Uchime and Perseus are preferred above the Chimera Slayer software. We chose to identify and remove chimera's using the Uchime software. As shown by Schloss and coworkers, Uchime is preferred to Perseus because of its lower false discovery rate and because of the previously reported faster execution times.





## 5.2 Comparative Genome Analysis of *Pediococcus damnosus* LMG 28219, a Strain Well-Adapted to the Beer Environment

I. Snauwaert, P. Stragier, L. De Vuyst, P. Vandamme

Submitted to: BCM Genomics, 2014

Author contributions: IS carried out DNA extraction, conducted genome assembly, performed sequence annotation, bioinformatics analyses, and comparative genome analyses. PS performed the OrthoMCL analysis. IS and PV designed and coordinated the study and participated in the analysis of the results. IS, PS, LDV, and PV wrote the manuscript. All authors read and approved the final manuscript.

*Pediococcus damnosus* LMG 28219 is a LAB dominating the maturation phase of Flemish acid beer productions. It proved to be capable of growing in beer, thereby resisting this environment, which is unfavorable for microbial growth. The molecular mechanisms underlying its metabolic capabilities and niche adaptations are currently unknown. In the present study, whole-genome sequencing and comparative genome analysis were used to investigate this strain's mechanisms to reside in the beer niche. The draft genome sequence of *P. damnosus* LMG 28219 harbored 183 contigs, including an intact prophage region and several CDSs involved in plasmid replication. The annotation of 2178 CDSs revealed the presence of many transporters and transcriptional regulators and several genes involved in oxidative stress response, hop resistance, *de novo* folate biosynthesis, and EPS production. Comparative genome analysis of *P. damnosus* LMG 28219 with *Pediococcus claussenii* ATCC BAA-344<sup>T</sup> (beer origin) and *Pediococcus pentosaceus* ATCC 25745 (plant origin) revealed that various hop resistance genes and genes involved in *de novo* folate biosynthesis were unique to the strains isolated from beer. This contrasted with the genes related to osmotic stress responses, which were shared between the strains compared. Furthermore, transcriptional regulators were enriched in the genomes of bacteria capable of growth in beer, suggesting that those cause rapid up- or down-regulation of gene expression. Genome sequence analysis of *P. damnosus* LMG 28219 provided insights into the underlying mechanisms of its adaptation to the beer niche. The results presented will enable analysis of the transcriptome and proteome of *P. damnosus* LMG 28219, which will result in additional knowledge on its metabolic activities.

### 5.2.1 Introduction

Beer is a fermented beverage that is high in ethanol, carbon dioxide and flavorful yeast metabolites, contains hop-derived flavor and antimicrobial components, and is low in pH, oxygen, and residual nutrients (Bokulich & Bamforth, 2013). This environment has selected for unique groups of bacteria specialized in growth in beer, including several species of LAB. Overall, one of the key metabolic actions of LAB is the reduction of pyruvate into lactate to regenerate  $\text{NAD}^+$  by means of lactate dehydrogenase activity. Depending on the beer type produced, lactate production may or may not be desired (Bokulich *et al.*, 2012a; Martens *et al.*, 1997; Van Oevelen *et al.*, 1977; Spitaels *et al.*, 2014b). As most LAB species tolerate high ethanol concentrations and a low pH, hop resistance is the major factor limiting their growth in beer. Hop-derived antimicrobial compounds, known as iso- $\alpha$  acids, cause permeability changes in the bacterial cell wall (Shimwell, 1937), leakage of the cytoplasmic membrane and subsequent inhibition of respiration and protein, DNA, and RNA syntheses (Teuber & Schmalreck, 1973), as well as changes in leucine uptake and proton ionophore activity (Simpson, 1993). Additionally, iso- $\alpha$  acids alter the redox properties of the bacterial cell, causing oxidative stress (Behr & Vogel, 2010). Accordingly, hop resistance is a multifactorial property that implies different mechanisms to counteract the action of iso- $\alpha$  acids. A key factor mediating hop resistance is the ATP-binding cassette multidrug transporter, *HorA*, which extrudes iso- $\alpha$  acids (Sakamoto *et al.*, 2001) and is plasmid-encoded (Sakamoto & Konings, 2003).

LAB species frequently encountered in the beer environment belong to the genera *Pediococcus* and *Lactobacillus*, which are both Gram-positive, facultative anaerobic, chemo-organotrophic bacteria of the family *Lactobacillaceae* of the *Firmicutes* phylum. The species with the highest capacity to grow in beer are *Lactobacillus brevis* and *Pediococcus damnosus*, although the ability of bacteria to grow in beer is a strain- rather than species-specific characteristic (Pittet *et al.*, 2011). Recent studies investigated the mechanisms of *Lactobacillus brevis* for overcoming stresses in beer by means of reverse transcription qPCR and proteomic analyses (Behr *et al.*, 2007; Bergsveinson *et al.*, 2012). Less is known about the adaptations of *P. damnosus* to the beer environment. So far, *P. clausenii* ATCC BAA-344<sup>T</sup> originating from spoiled beer, *P. pentosaceus* ATCC 25745 of plant origin, and *P. pentosaceus* IE-3 originating from a dairy effluent sample, are the only members of the genus *Pediococcus* for which the genome has been sequenced completely

(Midha *et al.*, 2012; Pittet *et al.*, 2012; Makarova *et al.*, 2006). However, draft genome sequences are available for several strains of *Pediococcus acidilactici*, namely strain MA 18/5M originating from animal food, strain DSM 20284 isolated from barley, strain 7\_4 from a human fecal sample, and strain NGRI 0510Q, formerly known as *Pediococcus lolii* (Wieme *et al.*, 2012), isolated from ryegrass silage (<http://genomesonline.org>). In general, *Pediococcus* species possess rather small genomes (approximately 2 MB), encoding a broad repertoire of transporters for efficient carbon and nitrogen acquisition and reflecting a limited range of biosynthetic capabilities. This suggests both extensive gene loss as well as acquisitions via horizontal gene transfer during the evolution of pediococci within their habitats (Makarova *et al.*, 2006). Several strains have plasmids containing genes regulating the fermentation of carbohydrates and encoding different types of resistances.

The strain *P. damnosus* LMG 28219 was isolated in 2013 from a Flemish acid beer at the end of its maturation phase. Flemish acid beers are produced by a mixed-culture fermentation and represent culturally important products, for which microbial activities play critical roles in beer production and quality formation (Martens *et al.*, 1997; Spitaels *et al.*, 2014b). In the present study, the draft genome sequence of *P. damnosus* LMG 28219 is presented and analyzed to obtain insights into its genome-based metabolic features. A better understanding of the molecular mechanisms underlying its metabolic capabilities enabled detailed insights into the mechanisms of adaptation of this strain to the beer environment. Furthermore, comparison of *P. damnosus* LMG 28219 with other sequenced members of the genus *Pediococcus* addressed the potentially unique properties of this strain and other strains adapted to the beer environment.

## 5.2.2 Materials and Methods

### Bacterial Strain and Growth Conditions

The strain *P. damnosus* LMG 28219 was used for genome sequencing. It was identified through sequence analysis of the *pheS* gene, as described previously (De Bruyne *et al.*, 2008b). To obtain cell pellets, *P. damnosus* LMG 28219 was propagated in MRS broth (Oxoid, Basingstoke, Hampshire, United Kingdom) at 28°C for 3 days, followed by microcentrifugation (11000 rpm, 15 min, 4°C).

## DNA Extraction and Illumina Sequencing

Total DNA was extracted using a procedure applied by Gevers and colleagues (2001), with several modifications. (i) The thawed pellet was washed in 1 ml TES buffer (6.7% sucrose, 50 mM Tris-HCl, pH 8.0, 1 mM EDTA) and resuspended in 275  $\mu$ L STET buffer (8.0% sucrose, 5.0% Triton X-100, 50 mM Tris-HCl, pH 8.0, 50 mM EDTA). (ii) Ninety  $\mu$ L lysozyme-mutanolysin-proteinase K solution (TES buffer containing 1667 U/mL mutanolysine, 33 mg/mL lysozyme and 2.78 mg/mL proteinase K) was added and the suspension was incubated at 37°C for 1 h. (iii) Prior to extraction with a phenol/chloroform/isoamylalcohol (49.5:49.5:1.0) solution, proteins were precipitated by adding 250  $\mu$ L ammonium acetate. (iv) Five  $\mu$ L RNase (2 mg/L) was added to the DNA solution, which was incubated at 37°C for 60 min and stored at -20°C. The integrity, concentration, and purity of the DNA isolated was evaluated using 1.0% (wt/vol) agarose gels, stained in ethidium bromide, and by spectrophotometric measurements at 234, 260, and 280 nm. A fluorescent stain-based kit (Qubit® dsDNA Broad range assay kit; Life Technologies, Carlsbad, CA, USA) was used for an accurate determination of the DNA concentration. Because of the low DNA concentration obtained, the extraction procedure was performed in triplicate and the resulting DNAs were pooled, after which an additional purification [by adding 12.5  $\mu$ L sodium acetate (3 M, pH 4.8) and 200  $\mu$ L 95% ice-cold ethanol to 100  $\mu$ L DNA product] and quantification step was performed.

## Illumina Sequencing and Genome Assembly

Library preparation and genome sequencing was performed by BaseClear BV (Leiden, The Netherlands). A paired-end DNA library with a mean insert length size between 230 and 360 bp was sequenced, with average reads of 100 bp, on an Illumina HiSeq2500 apparatus (Illumina Inc., San Diego, CA, USA). An initial *de novo* assembly was performed in CLC Genomics Workbench v6.5.1 (CLC Inc., Aarhus, Denmark), using quality-trimmed (based on a threshold of  $Q = 30$ ) and paired reads. All contigs shorter than 500 bp were discarded. Contigs were ordered automatically with MAUVE (Darling *et al.*, 2004) by comparing the draft genome sequence of *P. damnosus* LMG 28219 with the complete chromosomal DNA of *P. clausenii* ATCC BAA-344<sup>T</sup>. Contigs that did not show similarities towards the *P. clausenii* ATCC BAA-344<sup>T</sup> chromosomal DNA were blasted against a plasmid and phage database, using the PATRIC (*i.e.*, Pathosystems Resource Integration Center) (Wattam *et al.*, 2014) and PHAST (*i.e.*, PHAge Search Tool) (You *et al.*, 2011) websites, respectively. In addition, the average G+C content of the contigs was calculated using an in house developed Python script.

## Genome Annotation

Functional annotation and metabolic reconstruction were performed with (i) the RAST server (Aziz *et al.*, 2008), using GLIMMER (Delcher *et al.*, 1999) for gene calling and allowing frameshift corrections, backfilling of gaps, and automatic fixing of errors; (ii) the IMG-Expert review (IMG-ER) annotation pipeline, using GenePRIMP for gene prediction (Mavromatis *et al.*, 2009); and (iii) the NCBI's PGAAP, which uses GeneMark and GeneMark.HMM for gene calling (Lukashin & Borodovsky, 1998). The automated gene prediction and annotation was followed by manual curation of the CDSs of interest using BLASTp (Altschul *et al.*, 1997), UniProt (<http://www.uniprot.org/>), and InterProScan (Zdobnov & Apweiler, 2001).

## Comparative Genomics

Comparative genomics of the draft genome sequence of *P. damnosus* LMG 28219 was performed using the available complete genome sequences of *P. claussenii* ATCC BAA-344<sup>T</sup> and *P. pentosaceus* ATCC 25745. The ANIs between the draft genome (*P. damnosus* LMG 28219) and both reference genomes were calculated using the *in silico* DDH method implemented in the Jspecies software (Rosselló-Mora, 2006), using BLAST, as proposed by Goris and colleagues (2007). The OrthoMCL tool was used with default settings for ortholog finding in the genomes of *P. damnosus* LMG 28219, *P. claussenii* ATCC BAA-344<sup>T</sup>, and *P. pentosaceus* ATCC 25745 (<http://orthomcl.org/>). The Kyoto encyclopedia of genes and genomes (KEGG) database was used for the reconstruction of the folate biosynthesis metabolic map (<http://www.genome.jp/kegg/>).

## 5.2.3 Results and Discussion

### General Architecture and Annotation of the *Pediococcus damnosus* LMG 28219 Draft Genome

Paired-end sequencing of the *P. damnosus* LMG 28219 genomic DNA yielded 3,137,316 reads with a total number of 2,231,216 bp that were assembled into 183 contigs (N50 of 24659), consisting of 69 large (> 10,000 nucleotides) and 114 small

(< 10,000 nucleotides) contigs. This genome hence represents an intermediate size among the LAB (Midha *et al.*, 2012; Makarova *et al.*, 2006). An overview of the G+C content and the length of the contigs is presented in Supplementary Material 5.8. The G+C content of the complete draft genome averaged 38.2 mol%. A total of 91 contigs could be mapped onto the *P. clausenii* ATCC BAA-344<sup>T</sup> chromosomal DNA and were ordered accordingly (Table 8). Two clustered regularly interspaced short palindromic repeat (CRISPR) arrays were found on contigs 71 and 150, whereas three CRISPR-associated CDSs (AH70\_09625, AH70\_09630, and AH70\_09635) were found on contig 71. One intact prophage region (G+C content, 39.3%; region length, 39.4 kb) containing 50 CDSs was predicted and identified as the *Lactobacillus* phage Sha1, which was originally isolated from kimchi (Yoon *et al.*, 2011). This phage has an isomeric head and a long tail and is classified as a member of the large family of *Siphoviridae*. The genes of phage Sha1 are organized into five functional clusters: replication/regulation/modification, packaging, structure/morphogenesis, lysis, and lysogeny. Most of the phage-related CDSs were found on contig 7, which was predicted to be part of the chromosomal DNA.

A BLAST search of the remaining non-chromosomally encoded contigs against a plasmid-specific database revealed many similarities towards plasmids of different bacterial species (see below). Many of these contigs (as listed in Table 8) were similar to the plasmids of *P. clausenii* ATCC BAA-344<sup>T</sup>, with the exception of plasmids pPECL-1 and pPECL-2, onto which no contigs mapped. These two plasmids are small and cryptic, whereas the other six plasmids of *P. clausenii* ATCC BAA-344<sup>T</sup> range from 16 to 36 kb and contribute to roughly 7% of the strain's coding capacity (Pittet *et al.*, 2013). Four, six, ten, four, eight, and three contigs mapped onto plasmids pPECL-3, pPECL-4, pPECL-5, pPECL-6, pPECL-7, and pPECL-8 of *P. clausenii* ATCC BAA-344<sup>T</sup>, respectively. A total of 19 contigs did not show high similarities towards the plasmids of *P. clausenii* ATCC BAA-344<sup>T</sup>. Contigs 16 and 57 were similar to plasmids pBM400 and WSH-002\_p1 of *Bacillus megaterium* strains QM B1551 and WSH-002, respectively, whereas others were similar to plasmids of *Lactobacillus brevis* strains 925A, KB290, and ATCC 367 (contigs 123, 134, and 140, respectively), *Lactobacillus plantarum* WCFS1 (contig 37), *Lactobacillus buchneri* strains CD034 and NRRL B-30929 (contigs 100 and 122, respectively), or other *Lactobacillus* species (all remaining contigs). In addition, several CDSs involved in plasmid replication were found, which encoded replication initiation proteins (AH70\_10315 on contig 8, AH70\_03600 on contig 21, AH70\_00575 on contig 107, AH70\_02015 on contig 138, and AH70\_03140 on contig 182).

Table 8: Overview of the *P. damnosus* LMG 28219 draft genome.

Draft genome of <i>P. damnosus</i> LMG 28219	Mean G+C content (%)	Mean size consensus (bp)	Best BLAST hit
82, 98, 145, 116, 77, 119, 47, 53, 5, 89, 56, 76, 166, 84, 81, 136, 102, 141, 72, 30, 52, 129, 74, 54, 19, 110, 147, 49, 26, 85, 137, 61, 146, 177, 7, 68, 131, 104, 87, 32, 17, 80, 66, 165, 2, 20, 101, 96, 4, 59, 45, 164, 118, 64, 176, 69, 90, 24, 127, 10, 46, 139, 114, 60, 124, 23, 83, 51, 151, 169, 106, 11, 121, 78, 55, 43, 130, 48, 65, 168, 27, 62, 60, 79, 6, 152, 120, 73, 35, 99, 29, 71*	38.3	20814	chromosomal DNA (NC_016605)
21, 39, 41, 183	36.9	4040	pPECL-3 (NC_016636)
12, 21, 94, 95, 112, 135	38.5	3419	pPECL-4 (NC_016607)
1, 21, 39, 40, 95, 97, 107, 132, 135, 170	38.6	5198	pPECL-5 (NC_016608)
21, 107, 135, 178	37.0	4924	pPECL-6 (NC_017017)
8, 21, 39, 41, 86, 94, 182, 183	38.1	3926	pPECL-7 (NC_017018)
70, 14, 97*	42.5	9516	pPECL-8 (NC_017019)
16, 57	49.0	2538	<i>Bacillus megaterium</i> QM B1551 (pBM400, NC_004604 ) and WSH-002 (WSH-002_p1, CP003018)
123, 134, 140	37.2	4034	<i>Lb. brevis</i> 925A (pLB925A04, NC_012551), KB290 (pKB290-4, AP012171), and ATCC 367 (plasmid 1, NC_008498)
37	44.6	1242	<i>Lb. plantarum</i> WCFS1 (pWCFS103, NC_006377)
100, 122	33.9	5879	<i>Lb. buchneri</i> CD034 (pCD034-3, CP003044) and NRRL B-30929 (pLBUC01, NC_015420)
18, 22, 93, 108, 109, 113, 115, 143, 158, 162, 163	40.6	2797	Several <i>Lactobacillus</i> spp.

Contigs with the highest hit scores are highlighted in bold. \*Contig numbers are mentioned in ordered fashion. p: plasmid, *Lb.*: *Lactobacillus*.

Gene finding and annotation of the *P. damnosus* LMG 28219 draft genome with the IMG-ER software resulted in a total gene count of 2266, among which 96.12% were CDSs. A total of 79.08% of the CDSs could be assigned to a protein, among which 23.39% and 12.80% were predicted to be enzymes and transporters, respectively. This agrees with the property of LAB to encode a broad repertoire of transporters for efficient nutrient uptake and reflects their limited biosynthetic capabilities (Makarova *et al.*, 2006). Additionally, 2.60% of the genes encoded signal peptides, whereas 26.04% encoded transmembrane proteins. Furthermore, a total of 56 tRNA and three 5S, 16S, and 23S rRNA genes were predicted. As expected, most CDSs were involved in carbohydrate, protein, DNA, and RNA metabolism and in cell wall and capsule construction. These findings agreed with those of Makarova and colleagues (2006), who analyzed 12 genomes belonging to the order *Lactobacillales* and defined a core of *Lactobacillales*-specific clusters of orthologous groups (LaCOGs). The functional distribution of the conserved core of 567 LaCOGs showed that the majority encodes components of the information processing systems (translation, transcription, and replication), which are likely to perform essential functions. With the exception of 9 LaCOGs (*i.e.*, COGs 0230, 0419, 0454, 0457, 0470, 0762, 2815, 2855, and 4608), all core LaCOGs were found in the draft genome of *P. damnosus* LMG 28219. Genome closure could possibly lead to the identification of these missing core functions, although an update of the core LaCOGs is mandatory, because only 12 *Lactobacillales* species out of more than 500 species of this order were included; in addition, only one of these 12 species belonged to the genus *Pediococcus*.

Next to these core functions, the draft genome sequence of *P. damnosus* LMG 28219 comprised 50 predicted CDSs that were associated with virulence, disease, and defense, including three multidrug resistance efflux pumps. Also, 42 CDSs related to stress response were detected, seven of which were involved in oxidative stress response. Four of the non-core LaCOGs present in the *P. damnosus* LMG 28219 draft genome comprised 12 or more CDSs in a COG, namely arabinose efflux permeases (COG2814), transcriptional regulators (COG1309), predicted transcriptional regulators (COG0789), and transposases and their inactivated derivatives (COG2826). Arabinose efflux permeases belong to the major facilitator superfamily, a large and diverse group of secondary transporters that includes uniporters, symporters, and antiporters (Reddy *et al.*, 2012). A total of 12 and 10 CDSs of COG2814 were annotated as arabinose efflux permeases and drug resistance transporters from the EmrB/QacA subfamily, respectively. In addition, one CDS was predicted to be a drug resistance transporter of the Bcr/CflA subfamily and one was predicted to be a transporter belonging to the sugar transporter family. The COG1309 members belonged to the TetR family of transcriptional repressors, which



are involved in transcriptional control of multidrug efflux pumps, pathways for the biosynthesis of antibiotics, responses to osmotic stress and toxic chemicals, control of catabolic pathways, differentiation processes, and pathogenicity. Analysis of the CDSs of COG0789 did not reveal insights into their function. Finally, the CDSs in COG2826 were all transposases belonging to the insertion sequence IS30 family. The presence of IS elements may lead to genomic instability due to its role in gene loss and gene gain (Darmon & Leach, 2014).

### Comparative Genomics

The draft genome sequence of *P. damnosus* LMG 28219 was compared with the complete genome sequences of *P. pentosaceus* ATCC 25745 and *P. claussenii* ATCC BAA-344<sup>T</sup>. The number of CDSs predicted in the draft genome of *P. damnosus* LMG 28219 was higher compared to those in the reference genomes (Table 9). In addition, the number of genes in *P. claussenii* ATCC BAA-344<sup>T</sup> exceeded that of *P. pentosaceus* ATCC 25745. This could be explained by the need to acquire additional functions during evolution to be able to survive in the hostile beer environment, compared to the plant environment. Possibly, *P. pentosaceus* ATCC 25745 can acquire more nutrients from its environment compared to *P. damnosus* LMG 28219 and *P. claussenii* ATCC BAA-344<sup>T</sup>. The number of rRNA operons in the genome of *P. damnosus* LMG 28219 was lower compared to that in *P. pentosaceus* ATCC 25745 and *P. claussenii* ATCC BAA-344<sup>T</sup>, which may reflect differences in the ecological competitiveness (Makarova *et al.*, 2006; Klappenbach *et al.*, 2000; Di Mattia *et al.*, 2002). In addition, the ANIs of *P. damnosus* LMG 28219 versus *P. pentosaceus* ATCC 25745 and *P. claussenii* ATCC BAA-344<sup>T</sup> were 69.81% and 69.28%, respectively, whereas the ANIs between *P. pentosaceus* ATCC 25745 and *P. claussenii* ATCC BAA-344<sup>T</sup> were 71.53%. These results agreed with previous findings, indicating that *P. claussenii* and *P. pentosaceus* are evolutionarily closer related to each other compared to *P. damnosus* (De Bruyne *et al.*, 2008b).

The predicted proteome of *P. damnosus* LMG 28219 was assigned into orthologous clusters, along with the proteomes of *P. claussenii* ATCC BAA-344<sup>T</sup> and *P. pentosaceus* ATCC 25745 to predict unique and/or shared characteristics between these LAB species. A total of 1062 putative orthologous proteins were shared between *P. pentosaceus* ATCC 25745 (of plant origin), *P. claussenii* ATCC BAA-344<sup>T</sup>, and *P. damnosus* LMG 28219 (both of beer origin), whereas 25, 30, and 30 orthologs were unique to *P. pentosaceus* ATCC 25745, *P. claussenii* ATCC BAA-344<sup>T</sup>, and *P. damnosus* LMG 28219, respectively (Figure 18; Supplementary

Table 9: Comparison of genome characteristics of *P. damnosus* LMG 28219 with *P. pentosaceus* ATCC 25745 and *P. clausenii* ATCC BAA-344<sup>T</sup>

	<i>P. pentosaceus</i> ATCC 25745	<i>P. clausenii</i> ATCC BAA-344 <sup>T</sup>	<i>P. damnosus</i> LMG 28219
Accession number*	CP000422	CP003137*	JANK000000000
Origin	plant	beer	beer
Size	1,832,387	1,966,362	2,231,216
CDSs	1755	1892	2178
rRNA genes	5	4	3
tRNA genes	55	57	56
GC%	37.4	37.0	38.2

\* Only the Genbank record of the chromosomal DNA is given

Material 5.9). Genes that are shared between *P. damnosus* LMG 28219 and *P. clausenii* ATCC BAA-344<sup>T</sup>, but not by *P. pentosaceus* ATCC 25745, possibly promote survival in beer. For instance, the presence of genes involved in hop resistance, osmotic stress response, EPS production, and the presence or absence of complete or partial metabolic pathways may lead to the strain's ability to grow in beer. Furthermore, the functions enriched in the genomes of bacteria capable of growing in beer are likely to contribute to their specific phenotypic traits.

### Genes Involved in Hop Resistance

Research on marker genes associated with hop resistance suggested that these genes are typically acquired via horizontal gene transfer (Suzuki *et al.*, 2004; Iijima *et al.*, 2007; Hayashi *et al.*, 2001). Although several putative hop resistance genes have been described, only the presence of *horA* definitively correlates with LAB growth in beer, with the presence of *horA* together with *hitA* and/or *horC* allowing fast bacterial growth (Bergsveinson *et al.*, 2012). A recent transcriptome study of *P. clausenii* ATCC BAA-344<sup>T</sup> showed increased transcript levels of several genes during growth in beer compared to growth in MRS broth (Pittet *et al.*, 2013). This was especially the case for genes encoded on plasmids pPECL-3, pPECL-5, and pPECL-8. Contrasting to pPECL-3 and pPECL-5, many CDSs on pPECL-8 showed strong homology towards contigs 70, 14, and 97 in the draft genome of *P. damnosus* LMG 28219 (Figure 19). Plasmid pPECL-8 harbored the previously mentioned *horA* gene, which is involved in hop resistance. Surprisingly, several orthologs of the *P. clausenii* ATCC BAA 344<sup>T</sup> *horA* gene were found both in the draft genome of *P. damnosus* LMG 28219 (AH70\_02075 on contig





Figure 18: Visualization of the OrthoMCL output comparing the number of unique and/or shared orthologs of *P. clausenii* ATCC BAA-344<sup>T</sup>, *P. pentosaceus* ATCC 25745, and *P. damnosus* LMG 28219

14 and AH70\_01035 on contig 114) and in the complete genome of *P. pentosaceus* ATCC 25745 (YP\_805121), which all encoded ABC-type multidrug transporters. In depth analysis of these CDSs revealed that only AH70\_02075 showed high (99%) amino acid sequence homology towards the *horA* gene of *P. clausenii* ATCC BAA 344<sup>T</sup>; the remaining CDSs showed less than 70% homology. Possibly, these *horA* orthologs are descendants of a common ancestor that evolved towards a different functionality, with the *horA* gene in *P. clausenii* ATCC BAA-344<sup>T</sup> and AH70\_02075 in *P. damnosus* LMG 28219 being specialized in the extrusion of iso- $\alpha$  acids.

Interestingly, several studies revealed that the *horA* gene is consistently surrounded by the same set of genes in bacteria growing in beer, suggesting that these entire regions were acquired via horizontal gene transfer (Pittet *et al.*, 2013; Iijima *et al.*, 2007) and could possibly play a role in their adaptation to the beer niche. Most of these *horA*-surrounding genes are involved in phospholipid or cell wall biosynthesis. Orthologs of PECL\_1950, PECL\_1952, and PECL\_1954 flanking the *horA* gene were found in the genome of *P. damnosus* LMG 28219 (AH70\_09615 on contig 70 and AH70\_02080 and AH70\_02070 on contig 14, respectively) but not in the *P. pentosaceus* ATCC 25745 genome sequence. AH70\_09615 and AH70\_02080 were annotated as an acyl-phosphate glycerol 3-phosphate and a glycosyl transferase,

respectively. Both PECL\_1950 and PECL\_1952 are members of the lysophospholipid acyltransferase superfamily, which contains acyltransferases of *de novo* and remodeling pathways of glycerophospholipid biosynthesis. Finally, AH70\_02070 contained a family 8 glycosyl transferase domain that catalyzes the transfer of sugar moieties from activated donor molecules to specific acceptor molecules, forming glycosidic bonds.

pPECL-8 (*P. claussenii* ATCC BAA-344<sup>T</sup>)



Contigs 70, 14, and 97 of *P. damnosus* LMG 28219

Figure 19: Comparison of plasmid pPECL-8 from *P. claussenii* ATCC BAA 344<sup>T</sup> and contigs 70, 14, and 90 of *P. damnosus* LMG 28219

The Artemis comparison tool (Carver *et al.*, 2005) alignment was based on a BLASTn comparison using default settings. The red lines indicate that the aligned regions had the same orientation, whereas the blue lines indicate that the aligned regions were inversely oriented. The *horA* gene and its surrounding genes are indicated as grey blocks [from left to right: AH70\_091615 (glycosyl transferase), AH70\_09610 (glycosyl transferase), AH70\_02080 (acyl-phosphate glycerol 3-phosphate), HorA: AH70\_02075 (multidrug transporter), and AH70\_02070 (glycosyl transferase)].

Next to the *horA* gene, the *hitA* gene encodes a divalent cation transporter and confers hop resistance by importing manganese and thereby counteracting proton gradient dissipation (Hayashi *et al.*, 2001). Orthologs of the *hitA* gene of *Lb. brevis* L5784 (AB035808) were found in the genomes of *P. damnosus* LMG 28219 (AH70\_05540), *P. claussenii* ATCC BAA 344<sup>T</sup> (AEV94625.1), and *P. pentosaceus* ATCC 25745 (YP\_805164). The reason for the presence of the *hitA* gene in *P. pentosaceus* ATCC 25745 of plant origin remains unclear. Furthermore, HorC has been suggested to be a proton motive force-dependent multidrug transporter, whose expression is under control of the HorB transcriptional regulator (Iijima *et al.*, 2006). Orthologs of the *horC* gene of *Lactobacillus backii* LMG 23555 (BAF56899.1) were found on contig 18 (AH70\_03090) in the *P. damnosus* LMG 28219 and *P. claussenii* ATCC BAA 344<sup>T</sup> genome sequences (AEV95756.1), but not in that of *P. pentosaceus* ATCC 25745. Finally, *horB*, *bsrA*, and *bsrB* homologs were not found in the draft genome sequence of *P. damnosus* LMG 28219.

### Folate Biosynthesis

The beer isolates *P. damnosus* LMG 28219 and *P. claussenii* ATCC BAA-344<sup>T</sup> contained a set of genes involved in folate biosynthesis that were absent in the genome of *P. pentosaceus* ATCC 25745 (of plant origin). These included a 5,10-methylenetetrahydrofolate reductase (EC 1.5.1.20, AH70\_04245 on contig 29), a dihydropteroate synthase (EC 2.5.1.15, AH70\_05560 on contig 4), a 2-amino-4-hydroxy-6-hydroxymethyldihydropteridine pyrophosphokinase (EC 2.7.6.3, AH70\_05580 on contig 4), a GTP cyclohydrolase I type 1 (EC 3.5.4.16, AH70\_05575 on contig 4), and a dihydroneopterin aldolase (EC 4.1.2.25, AH70\_05585 on contig 4).

Because folate-dependent formylation of the initiator tRNA is a hallmark of bacterial translation and because bacteria cannot import formylmethionyl-tRNA, folate is essential for bacterial growth (de Crecy-Lagard *et al.*, 2007). Most bacteria make folate *de novo*, starting from GTP and chorismate (de Crecy-Lagard *et al.*, 2007). The first enzyme in *de novo* folate biosynthesis is GTP cyclohydrolase I that catalyzes a complex reaction, in which the five-membered imidazole ring of GTP is opened and a six-membered dihydropyrazine ring is formed (Figure 20). The resulting 7,8-dihydroneopterin triphosphate is then converted into the corresponding monophosphate by a specific pyrophosphatase. A pyrophosphatase was found in the genome of *P. claussenii* ATCC BAA-344<sup>T</sup> but was absent in the draft genome sequence of *P. damnosus* LMG 28219. BLASTp of the pyrophosphatase gene against the predicted CDS amino acid sequences of *P. damnosus* LMG 28219 did

not reveal the presence of a homologous gene. Possibly, the gene performing this function in *P. damnosus* LMG 28219 has not been previously identified or is missing due to non-orthologous gene replacement. Alternatively, some missing information in the draft genome sequence of *P. damnosus* LMG 28219 may account for this gap in the folate biosynthesis pathway. Dihydroneopterin aldolase subsequently releases glycoaldehyde to produce 6-hydroxymethyl-7,8-dihydropterin, which is then pyrophosphorylated by hydroxymethyldihydropterin pyrophosphokinase. 6-Hydroxymethyl-7,8-dihydropterin pyrophosphate and 4-aminobenzoic acid (pABA) moieties are condensed by dihydropteroate synthase and this results in the production of dihydropteroate. The enzymes involved in the conversion of dihydropteroate into tetrahydrofolate-polyglutamate are shared by the three genomes analyzed. These differences in folate biosynthesis potential between the *P. damnosus* LMG 28219, *P. clausenii* ATCC BAA-344<sup>T</sup>, and *P. pentosaceus* ATCC 25745 genomes may be explained by the environmental acquisition of folate by the latter strain. Plants produce folate *de novo*, so possibly *P. pentosaceus* ATCC 25745 harbors folate transporters to import folate produced by its host and lost genes involved in folate biosynthesis during evolution.




Figure 20: Folate biosynthesis pathway reconstruction

The enzymes found in both *P. dammosus* LMG 28219 and *P. clausenii* ATCC BAA 344<sup>T</sup> are highlighted in bold, with the exception of EC 3.1.3.1 that was only found in the genome of *P. clausenii* ATCC BAA 344<sup>T</sup> (highlighted in bold and italic). The remaining enzymes were shared by *P. dammosus* LMG 28219, *P. clausenii* ATCC BAA 344<sup>T</sup>, and *P. pentosaceus* ATCC 25745. The mechanism of the reaction highlighted with a dashed arrow is uncertain and needs further investigation. The figure was constructed using the ChemBioDraw software v. 13.0 (Perkin Elmer Inc., Waltham, MA, USA). ATP: adenosine triphosphate, ADP: adenosine diphosphate, AMP: adenosine monophosphate, NADP(H): nicotinamide adenine dinucleotide phosphate, P(P)i: (pyro)phosphate, EC: enzyme commission, GTP: guanoside triphosphate.

### EPS Production

Laboratory experiments indicated that *P. damnosus* LMG 28219 produces EPS. The ability of a strain to produce EPS is not directly correlated to its ability to reside in beer but probably has importance in biofilm formation (Pittet *et al.*, 2011), thereby enabling persistence in the brewery environment. The EPS produced by *P. claussenii* ATCC BAA 344<sup>T</sup> is a high-molecular-mass  $\beta$ -glucan produced by the action of a transmembrane glycosyl transferase (*gtf*) gene. This gene is located on plasmid pPECL-7, which is not essential for growth in beer (Pittet *et al.*, 2013). The fibrillar polymer consists of a trisaccharide repeating unit with a  $\beta$ -1,3-linked glucose backbone and branches made up of single  $\beta$ -1,2-linked D-glucopyranosyl residues. Walling and colleagues (2005) reported a glucosyl transferase gene (*dps*) in *P. damnosus* IOEB8801, originating from wine, that produces a linear backbone of 3- $\beta$ -D-glucose-1 moieties. Surprisingly, no homologies towards the *gtf* and *dps* genes were found in the draft genome sequence of *P. damnosus* LMG 28219. Yet, contig 56 harbored a CDS (AH70\_07835), containing a glycosyl transferase (group 2) domain, which may be involved in EPS production. OrthoMCL analysis indicated that this CDS was unique to *P. damnosus* LMG 28219. Other CDS candidates involved in EPS production by *P. damnosus* LMG 28219 were AH70\_01405, AH70\_01410, and AH70\_01415, which were all predicted to be present on contig 122. These proteins did not show amino acid sequence homologies of more than 70% towards the proteomes of *P. claussenii* ATCC BAA 344<sup>T</sup> and *P. pentosaceus* ATCC 25745. AH70\_01405 is an EPS biosynthesis protein, consisting of an AAA domain containing a P-loop, and showed similarities towards the capsular EPS family protein (EHO53752) found in the genome of *Lactobacillus kisonensis* F0435, originating from the human oral cavity (Chen *et al.*, 2010). AH70\_01410 is a lipopolysaccharide biosynthesis domain that is involved in the biosynthesis of EPS. This protein showed more than 70% protein sequence homology towards the chain length determinant protein (EEI18212) of *Lb. buchneri* ATCC 11577, isolated from the human oral cavity (Tilden & Svec, 1952). Finally, AH70\_01415 harbored a cell envelope-related transcriptional attenuator domain, which describes a domain of unknown function that is found in the predicted extracellular domain of a number of putative membrane-bound proteins (Zdobnov & Apweiler, 2001). One of those is CpsA, a putative regulatory protein involved in EPS biosynthesis (Zdobnov & Apweiler, 2001). AH70\_01415 showed more than 70% protein sequence homology towards a cell envelope-like function transcriptional attenuator common domain protein (EEI18211) of *Lb. buchneri* ATCC 11577 and a biofilm regulatory protein of *Lb. kisonensis* F0435 (see above). Besides the presence of proteins potentially involved in EPS production in *P. damnosus* LMG 28219 (as discussed above), the mechanism of EPS production in *P. damnosus* LMG 28219 remains unclear.



### *Genes Involved in Oxidative Stress Response*

Pittet and coworkers (2013) found a set of highly transcribed genes in *P. clausenii* ATCC BAA 344<sup>T</sup> during growth in beer as a response to the oxidative stress imposed by hops. These genes are manganese transport proteins, methionine sulfoxide reductases MrsA and MsrB as well as other metal transport and homeostasis proteins. Orthologs of these genes were found in the genomes of *P. damnosus* LMG 28219 and *P. pentosaceus* ATCC 25745, indicating that the presence of these genes is not unique to strains capable of growing in beer. These results indicate that growth in beer is a multifactorial reaction of the cell towards a challenging environment, not only involving the presence of specific genes but also the up-and down-regulation of specific sets of genes.

### *Functions Enriched in the Genomes of Bacteria Originating from Beer*

Next to functions unique to bacteria originating from beer, enriched functions could also provide insight into the mechanisms of niche adaptation. The four most abundant non-core LaCOGs (COG 2814, COG1309, COG0789, and COG2826, discussed above) in the *P. damnosus* LMG 28219 draft genome were also present in the *P. clausenii* ATCC BAA-344<sup>T</sup> and *P. pentosaceus* ATCC 25745 genomes, but were enriched only in the genomes of the bacterial strains originating from beer (i.e., *P. damnosus* LMG 28219 and *P. clausenii* ATCC BAA-344<sup>T</sup>) (Figure 21). Other enriched functions were COG1846 (transcriptional regulators), COG0596 (predicted hydrolases or acyltransferases of the alpha/beta hydrolase superfamily), and COG0745 (response regulators consisting of a CheY-like receiver domain and a winged-helix DNA-binding domain). The enrichment of the above-mentioned functions in bacteria growing in beer was further substantiated when analyzing additional genome sequences of the genus *Pediococcus* and *Lactobacillus*. Many COGs that were enriched in bacteria originating from beer are involved in transcriptional regulation, indicating that genes can be up- or down-regulated in hop-stressed cells. Currently, it is not clear in which processes these regulators are involved but this may be revealed by transcriptome analyses.

Figure 21: Overview of the functions enriched in the genomes of LAB of beer origin compared to those of LAB not originating from beer

The number of CDSs in the different COGs are listed. The color gradient ranges from white to black with increasing number of CDS and applies per row. IMG-ER was used for the determination of the number of CDSs. COG: cluster of orthologous groups, *Lb. Lactobacillus*, IS: insertion sequence.

## 5.2.4 Conclusions

The draft genome of *P. damnosus* LMG 28219 and its comparative analysis with other pediococci provided insights into the adaptation of this strain to the beer environment. These adaptations included the presence of the *horA* gene and its surrounding genes, genes involved in *de novo* folate biosynthesis, genes involved in the production of EPS, and the enrichment of functions related to transcriptional regulation. Furthermore, the results presented in this study will enable future transcriptome analysis of *P. damnosus* LMG 28219, which can provide additional insights into its metabolic activities.

## 5.2.5 Acknowledgments

The authors acknowledge the financial support of FWO-Flanders, BOF project and the Vrije Universiteit Brussel (SRP, IRP, and IOF projects), and of the Hercules Foundation. In addition, the authors would like to acknowledge the brewery involved and Charlotte Peeters and Bart Verheyde for the bioinformatics support.

## 5.2.6 Supplementary Material

The raw sequence data received from BaseClear BV were deposited at the SRA of GenBank (accession number SRP035530). This Whole-genome shotgun project was deposited at DDBJ/EMBL/GenBank under the accession number JANK000000000 after automatic annotation by PGAAP and manual curation of the genes of interest.

*Supplementary Material 5.8: Overview of the G+C content and the length of the contigs of the draft genome of P. damnosus LMG 28219*

contig no.	GC-content (%)	length (bp)	contig no.	GC-content (%)	length (bp)	contig no.	GC-content (%)	length (bp)
1	41.83	502	62	39.12	25631	123	38.10	1756
2	35.96	29429	63	38.10	1614	124	37.86	19366
3	35.66	3738	64	36.64	52573	125	42.39	2958
4	39.18	17052	65	38.67	22379	126	37.48	1174
5	34.15	9243	66	37.62	30560	127	34.09	20107
6	37.64	16203	67	38.29	1319	128	39.60	9231
7	39.31	53329	68	38.15	53951	129	37.49	5865
8	36.84	11309	69	36.79	15596	130	38.24	2691
9	41.49	1569	70	44.45	1982	131	39.49	5242
10	39.66	24545	71	37.03	18224	132	38.02	2967
11	37.39	27167	72	36.32	18506	133	43.34	953
12	37.00	5940	73	38.73	7361	134	34.51	6286
13	44.28	1154	74	38.62	35374	135	34.29	630
14	41.30	6274	75	45.65	5490	136	37.48	22993
15	43.70	1906	76	38.60	10142	137	36.68	3539
16	49.05	1790	77	39.44	21192	138	38.83	703
17	36.98	15880	78	44.06	6518	139	39.28	6973
18	42.10	14520	79	42.35	6140	140	39.10	4059
19	37.06	39520	80	37.70	29394	141	32.78	6114
20	37.61	15298	81	39.17	43485	142	34.46	711
21	38.42	10835	82	38.44	13240	143	34.61	1598
22	39.32	5570	83	38.08	17031	144	37.44	2094
23	37.66	28945	84	39.63	10334	145	41.14	6624
24	35.63	10409	85	37.86	14162	146	34.62	11446
25	37.24	1450	86	41.89	2134	147	38.32	6765
26	37.53	22807	87	36.99	2933	148	38.77	1251
27	39.68	41367	88	41.39	720	149	33.61	2565
28	38.52	3580	89	33.63	5376	150	39.22	2675
29	37.66	112846	90	38.38	7835	151	40.74	7364
30	37.97	23931	91	40.83	5787	152	39.43	8752
31	36.80	8354	92	43.40	735	153	36.07	1594
32	36.98	67552	93	43.89	565	154	34.98	1921
33	39.12	1337	94	46.11	1028	155	35.21	676
34	37.68	6696	95	37.95	1149	156	34.56	2584
35	39.11	42230	96	35.55	7300	157	41.35	1468
36	47.74	1906	97	41.72	20292	158	41.37	556
37	44.61	1242	98	39.53	9354	159	30.99	755
38	41.52	9357	99	37.68	9719	160	37.77	5809
39	38.53	1547	100	34.96	1533	161	46.71	1824
40	37.76	1213	101	37.85	21944	162	39.71	1244
41	36.10	2906	102	40.15	24659	163	39.02	3350
42	39.01	905	103	37.56	13572	164	35.96	2912
43	36.94	25254	104	37.97	31232	165	38.27	2710
44	46.78	6624	105	38.67	2801	166	35.70	2678
45	37.45	23239	106	40.88	2275	167	36.36	1276
46	39.37	68863	107	36.22	7228	168	39.08	16295
47	40.25	20521	108	44.34	857	169	38.69	4125
48	36.41	11759	109	41.12	642	170	41.54	5611
49	37.67	36252	110	39.39	12345	171	32.33	532
50	37.34	31306	111	40.97	3937	172	30.56	1705
51	37.11	16670	112	36.98	933	173	39.62	785
52	39.53	55967	113	41.76	771	174	36.17	1247
53	38.50	59279	114	38.19	40025	175	36.95	609
54	38.44	23452	115	39.38	1097	176	49.68	1872
55	36.90	18764	116	39.03	16480	177	43.89	1399
56	38.94	35768	117	37.61	1550	178	38.88	1003
57	48.97	3286	118	39.27	4077	179	45.93	553
58	36.80	22440	119	38.40	22575	180	33.47	983
59	36.99	22335	120	37.74	12548	181	41.15	593
60	40.23	17760	121	38.45	3555	182	32.04	774
61	37.60	8570	122	32.87	10225	183	34.59	873

*Supplementary Material 5.9: Shared and unique orthologous proteins of P. damnosus LMG 28219, P. clausenii ATCC BAA 344<sup>T</sup>, and P. pentosaceus ATCC 25745*

clau: *P. clausenii* ATCC BAA 344<sup>T</sup>, ldam: *P. damnosus* LMG 28219, pent: *P. pentosaceus* ATCC 25745

is0000: clau|AEV94415.1 clau|AEV95932.1 clau|AEV95653.1 ldam|AH70\_02430 ldam|AH70\_03715 ldam|AH70\_03720 ldam|AH70\_03725 ldam|AH70\_07075 ldam|AH70\_01065 pent|YP\_805047 pent|YP\_803934  
is0001: clau|AEV95460.1 clau|AEV95547.1 clau|AEV95762.1 ldam|AH70\_05025 ldam|AH70\_06205 ldam|AH70\_09260 pent|YP\_804136 pent|YP\_804474 pent|YP\_804906 pent|YP\_804593  
is0002: clau|AEV96110.1 clau|AEV96124.1 clau|AEV96156.1 clau|AFD55023.1 ldam|AH70\_03040 ldam|AH70\_05620 ldam|AH70\_11090 ldam|AH70\_11270  
is0003: clau|AEV94465.1 clau|AEV96016.1 ldam|AH70\_01170 ldam|AH70\_04480 ldam|AH70\_08070 pent|YP\_803658 pent|YP\_803822  
is0004: clau|AEV96148.1 clau|AEV96190.2 ldam|AH70\_00605 ldam|AH70\_11200 ldam|AH70\_03075 ldam|AH70\_03865 pent|YP\_805170  
is0005: ldam|AH70\_01040 ldam|AH70\_02645 ldam|AH70\_10065 ldam|AH70\_02650 ldam|AH70\_07425 ldam|AH70\_08475 ldam|AH70\_09300  
is0006: clau|AEV94344.1 clau|AEV95200.1 ldam|AH70\_04185 ldam|AH70\_05235 pent|YP\_803558 pent|YP\_804457  
is0007: clau|AEV94407.1 clau|AEV95586.1 pent|YP\_803794 pent|YP\_803942 ldam|AH70\_00840 ldam|AH70\_08055  
is0008: clau|AEV94515.1 clau|AEV95780.1 clau|AEV96066.1 ldam|AH70\_01320 pent|YP\_803931 ldam|AH70\_06130  
is0009: clau|AEV95463.1 clau|AEV95699.1 ldam|AH70\_00670 ldam|AH70\_09265 pent|YP\_803905 pent|YP\_804133  
is0010: ldam|AH70\_01495 ldam|AH70\_08760 ldam|AH70\_06900 ldam|AH70\_05735 ldam|AH70\_02020 ldam|AH70\_10145  
is0011: clau|AEV94632.1 ldam|AH70\_01940 ldam|AH70\_10475 pent|YP\_804995 clau|AEV95979.1 pent|YP\_805114  
is0012: clau|AEV94454.1 clau|AEV96074.1 clau|AEV95794.1 pent|YP\_804423 ldam|AH70\_06030  
is0013: clau|AEV94821.1 clau|AEV94913.1 pent|YP\_803615 pent|YP\_803966 ldam|AH70\_09875  
is0014: clau|AEV96109.1 clau|AEV96157.1 clau|AEV96123.1 clau|AFD55024.1 ldam|AH70\_03620  
is0015: ldam|AH70\_02400 ldam|AH70\_04355 clau|AEV96063.1 pent|YP\_804927 pent|YP\_805296  
is0016: ldam|AH70\_03055 ldam|AH70\_07765 ldam|AH70\_07795 clau|AEV94710.1 pent|YP\_804054  
is0017: ldam|AH70\_08370 ldam|AH70\_09835 pent|YP\_803719 pent|YP\_805210 clau|AEV96005.1  
is0018: pent|YP\_803703 pent|YP\_804555 pent|YP\_804045 pent|YP\_804079 pent|YP\_805062  
is0019: ldam|AEV94860.1 ldam|AH70\_07675 clau|AEV95588.1 ldam|AH70\_01525 pent|YP\_803940  
is0020: clau|AEV94602.1 clau|AEV96113.1 ldam|AH70\_01240 pent|YP\_804037  
is0021: clau|AEV94627.1 clau|AEV94669.1 ldam|AH70\_04530 pent|YP\_803636  
is0022: clau|AEV94628.1 clau|AEV95664.1 ldam|AH70\_04535 pent|YP\_803637  
is0023: clau|AEV94795.1 clau|AFD55028.1 ldam|AH70\_01045 ldam|AH70\_03875  
is0024: clau|AEV95797.1 clau|AEV96182.1 ldam|AH70\_08980 pent|YP\_805108  
is0025: clau|AEV95888.1 clau|AEV95889.1 pent|YP\_805266 ldam|AH70\_04700  
is0026: clau|AEV96083.1 clau|AEV96093.1 clau|AEV96188.1 ldam|AH70\_10315  
is0027: clau|AEV96119.1 clau|AEV96162.1 ldam|AH70\_03600 ldam|AH70\_00575  
is0028: ldam|AH70\_00820 ldam|AH70\_01285 clau|AEV96115.1 pent|YP\_804032  
is0029: ldam|AH70\_01030 ldam|AH70\_02720 clau|AEV94534.1 pent|YP\_805122  
is0030: ldam|AH70\_01035 ldam|AH70\_02075 clau|AEV96194.1 pent|YP\_805121  
is0031: ldam|AH70\_08835 ldam|AH70\_10820 ldam|AH70\_03360 ldam|AH70\_06115  
is0032: ldam|AH70\_01270 ldam|AH70\_02155 ldam|AH70\_03595 clau|AEV96118.1  
is0033: ldam|AH70\_01335 ldam|AH70\_06980 clau|AEV95990.1 pent|YP\_805277  
is0034: ldam|AH70\_01420 ldam|AH70\_01835 ldam|AH70\_03395 clau|AEV96185.1  
is0035: ldam|AH70\_01660 ldam|AH70\_01665 clau|AEV95481.1 pent|YP\_804113  
is0036: ldam|AH70\_01810 ldam|AH70\_03840 clau|AEV95485.1 pent|YP\_804104  
is0037: ldam|AH70\_01850 ldam|AH70\_03095 clau|AEV96173.1 pent|YP\_804033  
is0038: ldam|AH70\_02050 ldam|AH70\_06475 ldam|AH70\_11060 pent|YP\_803954  
is0039: ldam|AH70\_02480 ldam|AH70\_08740 clau|AEV96170.1 pent|YP\_805144  
is0040: ldam|AH70\_02515 ldam|AH70\_10990 ldam|AH70\_01120 ldam|AH70\_05655  
is0041: ldam|AH70\_02670 ldam|AH70\_10425 clau|AEV95240.1 pent|YP\_804410  
is0042: ldam|AH70\_04015 ldam|AH70\_06460 clau|AEV95437.1 pent|YP\_804167  
is0043: ldam|AH70\_05435 ldam|AH70\_07100 clau|AEV94712.1 pent|YP\_803867  
is0044: ldam|AH70\_06240 ldam|AH70\_11190 clau|AEV96052.1 pent|YP\_804728  
is0045: ldam|AH70\_06530 ldam|AH70\_10445 clau|AEV94387.1 pent|YP\_803603  
is0046: ldam|AH70\_07095 ldam|AH70\_08780 ldam|AH70\_10125 clau|AEV96025.1  
is0047: ldam|AH70\_07210 ldam|AH70\_10440 ldam|AH70\_07245 clau|AEV95904.1  
is0048: ldam|AH70\_07235 ldam|AH70\_09690 clau|AEV94930.1 pent|YP\_805282  
is0049: ldam|AH70\_07775 ldam|AH70\_10230 clau|AEV95913.1 pent|YP\_804465  
is0050: ldam|AH70\_08310 ldam|AH70\_11215 clau|AEV96199.1 pent|YP\_804726  
is0051: ldam|AH70\_09460 ldam|AH70\_10290 clau|AEV94894.1 pent|YP\_804504  
is0052: ldam|AH70\_10005 ldam|AH70\_10995 clau|AEV95471.1 pent|YP\_804126  
is0053: pent|YP\_803729 pent|YP\_805064 clau|AEV96015.1 ldam|AH70\_04485  
is0054: pent|YP\_803842 pent|YP\_803919 clau|AEV94745.1 ldam|AH70\_07525  
is0055: pent|YP\_803850 pent|YP\_805265 clau|AEV94498.1 ldam|AH70\_07445  
is0056: pent|YP\_803953 pent|YP\_805169 clau|AEV94766.1 ldam|AH70\_07355  
is0057: pent|YP\_804144 pent|YP\_804735 clau|AEV95025.1 ldam|AH70\_05545  
is0058: pent|YP\_804251 pent|YP\_804532 clau|AEV95150.1 ldam|AH70\_09565  
is0059: pent|YP\_804280 pent|YP\_804506 clau|AEV94889.1 ldam|AH70\_09485  
is0060: pent|YP\_804402 pent|YP\_805227 clau|AEV94555.1 ldam|AH70\_07970  
is0061: ldam|AH70\_06575 ldam|AH70\_10765 ldam|AH70\_06865 ldam|AH70\_06580  
is0062: clau|AEV94358.1 ldam|AH70\_10660 pent|YP\_803588 pent|YP\_805105  
is0063: clau|AEV94359.1 ldam|AH70\_05550 pent|YP\_805104 pent|YP\_803824  
is0064: clau|AEV94383.1 pent|YP\_803653 clau|AEV96058.1 ldam|AH70\_06960  
is0065: clau|AEV94389.1 pent|YP\_804544 clau|AEV95765.1 ldam|AH70\_09650

is0066: clau|AEV94392.1 ldam|AH70\_09580 clau|AEV96114.1 pent|YP\_804038  
is0067: clau|AEV94438.1 pent|YP\_804361 clau|AEV94674.1 ldam|AH70\_10550  
is0068: clau|AEV94440.1 pent|YP\_804363 clau|AEV94675.1 ldam|AH70\_10555  
is0069: clau|AEV94441.1 pent|YP\_804364 clau|AEV94676.1 ldam|AH70\_10560  
is0070: clau|AEV94442.1 pent|YP\_804365 clau|AEV94677.1 ldam|AH70\_10565  
is0071: clau|AEV94484.1 ldam|AH70\_04720 pent|YP\_803828 pent|YP\_803639  
is0072: clau|AEV94512.1 ldam|AH70\_06815 ldam|AH70\_06955 pent|YP\_805058  
is0073: clau|AEV94561.1 ldam|AH70\_07395 clau|AEV95958.1 pent|YP\_803596  
is0074: clau|AEV94568.1 ldam|AH70\_03440 clau|AEV95456.1 pent|YP\_804140  
is0075: clau|AEV94647.1 ldam|AH70\_06860 pent|YP\_803577 pent|YP\_804981  
is0076: clau|AEV94708.1 ldam|AH70\_04665 pent|YP\_805110 pent|YP\_804623  
is0077: clau|AEV94755.1 ldam|AH70\_01855 clau|AEV95802.1 pent|YP\_804106  
is0078: clau|AEV94807.1 ldam|AH70\_04870 pent|YP\_804948 pent|YP\_803750  
is0079: clau|AEV94876.1 ldam|AH70\_03315 clau|AEV96022.1 pent|YP\_805244  
is0080: clau|AEV95036.1 pent|YP\_804722 clau|AEV95668.1 ldam|AH70\_09680  
is0081: clau|AEV95338.1 ldam|AH70\_11130 pent|YP\_804312 ldam|AH70\_11135  
is0082: clau|AEV95468.1 ldam|AH70\_11015 pent|YP\_803674 pent|YP\_805067  
is0083: clau|AEV95486.1 ldam|AH70\_05300 pent|YP\_805066 pent|YP\_805071  
is0084: clau|AEV95798.1 ldam|AH70\_08975 clau|AEV96183.1 pent|YP\_805109  
is0085: clau|AEV95831.1 ldam|AH70\_01710 pent|YP\_803601 pent|YP\_805148  
is0086: clau|AEV95835.1 ldam|AH70\_00940 clau|AEV95855.1 pent|YP\_805175  
is0087: clau|AEV95860.1 ldam|AH70\_02675 clau|AEV96096.1 pent|YP\_803702  
is0088: clau|AEV95884.1 ldam|AH70\_03500 pent|YP\_804540 pent|YP\_804051  
is0089: clau|AEV96192.1 ldam|AH70\_09610 pent|YP\_803759 ldam|AH70\_08745  
is0090: clau|AEV94734.1 clau|AEV94735.1 pent|YP\_805068  
is0091: clau|AEV95602.1 clau|AEV95610.1 clau|AEV95603.1  
is0092: clau|AEV95615.1 clau|AEV95621.1 clau|AEV95617.1  
is0093: clau|AEV95754.1 clau|AEV95941.1 ldam|AH70\_03640  
is0094: clau|AEV96092.1 clau|AEV96189.1 ldam|AH70\_05630  
is0095: clau|AEV96099.1 clau|AEV96179.1 ldam|AH70\_03145  
is0096: clau|AEV96105.1 clau|AEV96112.1 ldam|AH70\_08970  
is0097: clau|AEV96121.1 clau|AFD55026.1 ldam|AH70\_01860  
is0098: clau|AEV96140.1 clau|AFD55022.1 ldam|AH70\_01815  
is0099: clau|AEV96142.1 clau|AEV96206.1 ldam|AH70\_11255  
is0100: ldam|AH70\_00345 ldam|AH70\_11230 clau|AEV96203.1  
is0101: ldam|AH70\_00835 ldam|AH70\_03900 ldam|AH70\_03855  
is0102: ldam|AH70\_00865 ldam|AH70\_10190 pent|YP\_804449  
is0103: ldam|AH70\_00985 ldam|AH70\_01290 ldam|AH70\_07910  
is0104: ldam|AH70\_01260 ldam|AH70\_01515 clau|AEV96031.1  
is0105: ldam|AH70\_01305 ldam|AH70\_02490 ldam|AH70\_01745  
is0106: ldam|AH70\_01920 ldam|AH70\_05480 ldam|AH70\_02555  
is0107: ldam|AH70\_02340 ldam|AH70\_08735 ldam|AH70\_02485  
is0108: ldam|AH70\_03165 ldam|AH70\_04585 ldam|AH70\_04610  
is0109: ldam|AH70\_03265 ldam|AH70\_03310 clau|AEV94877.1  
is0110: ldam|AH70\_04255 ldam|AH70\_07600 clau|AEV95845.1  
is0111: ldam|AH70\_04670 ldam|AH70\_04920 clau|AEV94852.1  
is0112: ldam|AH70\_05350 ldam|AH70\_09665 pent|YP\_803604  
is0113: ldam|AH70\_05470 ldam|AH70\_06015 ldam|AH70\_05615  
is0114: ldam|AH70\_05725 ldam|AH70\_08765 ldam|AH70\_07160  
is0115: ldam|AH70\_07085 ldam|AH70\_08755 clau|AEV95658.1  
is0116: ldam|AH70\_07270 ldam|AH70\_07840 ldam|AH70\_07845  
is0117: ldam|AH70\_08250 ldam|AH70\_10705 pent|YP\_804836  
is0118: pent|YP\_803711 pent|YP\_804553 ldam|AH70\_01325  
is0119: pent|YP\_804248 pent|YP\_804536 ldam|AH70\_02630  
is0120: pent|YP\_804259 pent|YP\_804521 ldam|AH70\_09535  
is0121: clau|AEV94340.1 ldam|AH70\_04205 pent|YP\_803554  
is0122: clau|AEV94341.1 ldam|AH70\_04200 pent|YP\_803555  
is0123: clau|AEV94342.1 ldam|AH70\_04195 pent|YP\_803556  
is0124: clau|AEV94343.1 ldam|AH70\_04190 pent|YP\_803557  
is0125: clau|AEV94345.1 ldam|AH70\_04180 pent|YP\_803559  
is0126: clau|AEV94346.1 ldam|AH70\_10645 pent|YP\_803560  
is0127: clau|AEV94347.1 ldam|AH70\_10650 pent|YP\_803561  
is0128: clau|AEV94348.1 ldam|AH70\_10655 pent|YP\_803562  
is0129: clau|AEV94349.1 ldam|AH70\_10675 pent|YP\_803563  
is0130: clau|AEV94350.1 ldam|AH70\_10680 pent|YP\_803564  
is0131: clau|AEV94351.1 ldam|AH70\_10685 pent|YP\_803565  
is0132: clau|AEV94352.1 ldam|AH70\_10690 pent|YP\_803566  
is0133: clau|AEV94354.1 ldam|AH70\_01115 pent|YP\_803567  
is0134: clau|AEV94357.1 ldam|AH70\_01220 pent|YP\_803569  
is0135: clau|AEV94361.1 ldam|AH70\_01695 pent|YP\_803826  
is0136: clau|AEV94368.1 ldam|AH70\_09525 pent|YP\_804859  
is0137: clau|AEV94370.1 ldam|AH70\_03755 pent|YP\_805224  
is0138: clau|AEV94371.1 ldam|AH70\_03760 pent|YP\_805223  
is0139: clau|AEV94372.1 ldam|AH70\_03765 pent|YP\_805222  
is0140: clau|AEV94373.1 ldam|AH70\_03770 pent|YP\_805221  
is0141: clau|AEV94374.1 ldam|AH70\_03775 pent|YP\_805220  
is0142: clau|AEV94375.1 ldam|AH70\_03785 pent|YP\_805218  
is0143: clau|AEV94378.1 ldam|AH70\_01455 pent|YP\_803692  
is0144: clau|AEV94379.1 ldam|AH70\_04440 pent|YP\_803651  
is0145: clau|AEV94380.1 ldam|AH70\_04445 pent|YP\_803652  
is0146: clau|AEV94382.1 ldam|AH70\_00855 pent|YP\_804860  
is0147: clau|AEV94384.1 ldam|AH70\_04460 pent|YP\_803654  
is0148: clau|AEV94385.1 ldam|AH70\_04470 pent|YP\_805171

is0149: clau|AEV94386.1 ldam|AH70\_04390 pent|YP\_803776  
is0150: clau|AEV94390.1 ldam|AH70\_10715 pent|YP\_804116  
is0151: clau|AEV94391.1 ldam|AH70\_10795 pent|YP\_803778  
is0152: clau|AEV94393.1 ldam|AH70\_11120 pent|YP\_805188  
is0153: clau|AEV94394.1 ldam|AH70\_11280 pent|YP\_803779  
is0154: clau|AEV94395.1 ldam|AH70\_11285 pent|YP\_803780  
is0155: clau|AEV94399.1 ldam|AH70\_11290 pent|YP\_803781  
is0156: clau|AEV94400.1 ldam|AH70\_11295 pent|YP\_803782  
is0157: clau|AEV94401.1 ldam|AH70\_11300 pent|YP\_803783  
is0158: clau|AEV94402.1 ldam|AH70\_11305 pent|YP\_803784  
is0159: clau|AEV94408.1 ldam|AH70\_02170 pent|YP\_803795  
is0160: clau|AEV94409.1 ldam|AH70\_07265 pent|YP\_804421  
is0161: clau|AEV94412.1 ldam|AH70\_02180 pent|YP\_803796  
is0162: clau|AEV94413.1 ldam|AH70\_02185 pent|YP\_803797  
is0163: clau|AEV94414.1 ldam|AH70\_02190 pent|YP\_803798  
is0164: clau|AEV94416.1 ldam|AH70\_10070 pent|YP\_803801  
is0165: clau|AEV94418.1 ldam|AH70\_10095 pent|YP\_803804  
is0166: clau|AEV94419.1 ldam|AH70\_10100 pent|YP\_803805  
is0167: clau|AEV94420.1 ldam|AH70\_10110 pent|YP\_803806  
is0168: clau|AEV94421.1 ldam|AH70\_10115 pent|YP\_803807  
is0169: clau|AEV94423.1 ldam|AH70\_10155 pent|YP\_803809  
is0170: clau|AEV94424.1 ldam|AH70\_10160 pent|YP\_803810  
is0171: clau|AEV94425.1 ldam|AH70\_10180 pent|YP\_803811  
is0172: clau|AEV94426.1 ldam|AH70\_01255 pent|YP\_803812  
is0173: clau|AEV94427.1 ldam|AH70\_01245 pent|YP\_803813  
is0174: clau|AEV94435.1 ldam|AH70\_03470 pent|YP\_804358  
is0175: clau|AEV94448.1 ldam|AH70\_00910 pent|YP\_803609  
is0176: clau|AEV94450.1 ldam|AH70\_04620 pent|YP\_805208  
is0177: clau|AEV94451.1 ldam|AH70\_01235 pent|YP\_803816  
is0178: clau|AEV94452.1 ldam|AH70\_01230 pent|YP\_803817  
is0179: clau|AEV94453.1 ldam|AH70\_01225 pent|YP\_803818  
is0180: clau|AEV94455.1 ldam|AH70\_04110 pent|YP\_804838  
is0181: clau|AEV94462.1 ldam|AH70\_01190 pent|YP\_803819  
is0182: clau|AEV94463.1 ldam|AH70\_01180 pent|YP\_803820  
is0183: clau|AEV94464.1 ldam|AH70\_01175 pent|YP\_803821  
is0184: clau|AEV94466.1 ldam|AH70\_01165 pent|YP\_803823  
is0185: clau|AEV94469.1 ldam|AH70\_06480 pent|YP\_803679  
is0186: clau|AEV94470.1 ldam|AH70\_06485 pent|YP\_803680  
is0187: clau|AEV94471.1 ldam|AH70\_06490 pent|YP\_803681  
is0188: clau|AEV94472.1 ldam|AH70\_06495 pent|YP\_803682  
is0189: clau|AEV94473.1 ldam|AH70\_06500 pent|YP\_803683  
is0190: clau|AEV94474.1 ldam|AH70\_06505 pent|YP\_803684  
is0191: clau|AEV94475.1 ldam|AH70\_06510 pent|YP\_803685  
is0192: clau|AEV94477.1 ldam|AH70\_06515 pent|YP\_803686  
is0193: clau|AEV94478.1 ldam|AH70\_06520 pent|YP\_803687  
is0194: clau|AEV94483.1 ldam|AH70\_04900 pent|YP\_803827  
is0195: clau|AEV94485.1 ldam|AH70\_07620 pent|YP\_803829  
is0196: clau|AEV94486.1 ldam|AH70\_07615 pent|YP\_803830  
is0197: clau|AEV94487.1 ldam|AH70\_07590 pent|YP\_803831  
is0198: clau|AEV94488.1 ldam|AH70\_07585 pent|YP\_803832  
is0199: clau|AEV94489.1 ldam|AH70\_07580 pent|YP\_803834  
is0200: clau|AEV94493.1 ldam|AH70\_07555 pent|YP\_803838  
is0201: clau|AEV94494.1 ldam|AH70\_07560 pent|YP\_803837  
is0202: clau|AEV94495.1 pent|YP\_803836 ldam|AH70\_07570  
is0203: clau|AEV94497.1 ldam|AH70\_07450 pent|YP\_803849  
is0204: clau|AEV94499.1 ldam|AH70\_10575 pent|YP\_803848  
is0205: clau|AEV94506.1 ldam|AH70\_07220 pent|YP\_805264  
is0206: clau|AEV94509.1 ldam|AH70\_07480 pent|YP\_803845  
is0207: clau|AEV94511.1 ldam|AH70\_06820 pent|YP\_805059  
is0208: clau|AEV94513.1 ldam|AH70\_06810 pent|YP\_805057  
is0209: clau|AEV94514.1 ldam|AH70\_10960 pent|YP\_805056  
is0210: clau|AEV94516.1 ldam|AH70\_10895 pent|YP\_805053  
is0211: clau|AEV94517.1 ldam|AH70\_10900 pent|YP\_805052  
is0212: clau|AEV94518.1 ldam|AH70\_10905 pent|YP\_805051  
is0213: clau|AEV94519.1 ldam|AH70\_10910 pent|YP\_805050  
is0214: clau|AEV94522.1 ldam|AH70\_01095 pent|YP\_805049  
is0215: clau|AEV94525.1 ldam|AH70\_01075 pent|YP\_805048  
is0216: clau|AEV94526.1 ldam|AH70\_08220 pent|YP\_805046  
is0217: clau|AEV94527.1 ldam|AH70\_09645 pent|YP\_805045  
is0218: clau|AEV94528.1 ldam|AH70\_07790 pent|YP\_805044  
is0219: clau|AEV94529.1 ldam|AH70\_07155 pent|YP\_805043  
is0220: clau|AEV94531.1 ldam|AH70\_07800 pent|YP\_805041  
is0221: clau|AEV94532.1 ldam|AH70\_06010 pent|YP\_803765  
is0222: clau|AEV94536.1 pent|YP\_805039 ldam|AH70\_07855  
is0223: clau|AEV94537.1 ldam|AH70\_07865 pent|YP\_805037  
is0224: clau|AEV94538.1 ldam|AH70\_07870 pent|YP\_805036  
is0225: clau|AEV94539.1 ldam|AH70\_07875 pent|YP\_805035  
is0226: clau|AEV94540.1 ldam|AH70\_07880 pent|YP\_805034  
is0227: clau|AEV94541.1 ldam|AH70\_07885 pent|YP\_805033  
is0228: clau|AEV94542.1 ldam|AH70\_07890 pent|YP\_805032  
is0229: clau|AEV94543.1 ldam|AH70\_07895 pent|YP\_805031  
is0230: clau|AEV94544.1 ldam|AH70\_07900 pent|YP\_805030  
is0231: clau|AEV94545.1 ldam|AH70\_07905 pent|YP\_805029

is0232: clau|AEV94546.1 ldam|AH70\_07915 pent|YP\_805028  
is0233: clau|AEV94547.1 ldam|AH70\_07920 pent|YP\_805027  
is0234: clau|AEV94548.1 ldam|AH70\_07925 pent|YP\_805026  
is0235: clau|AEV94549.1 ldam|AH70\_07930 pent|YP\_805025  
is0236: clau|AEV94550.1 ldam|AH70\_07935 pent|YP\_805024  
is0237: clau|AEV94551.1 ldam|AH70\_07940 pent|YP\_805023  
is0238: clau|AEV94552.1 ldam|AH70\_07945 pent|YP\_805022  
is0239: clau|AEV94553.1 ldam|AH70\_07950 pent|YP\_805021  
is0240: clau|AEV94554.1 ldam|AH70\_07955 pent|YP\_805020  
is0241: clau|AEV94556.1 ldam|AH70\_07980 pent|YP\_805019  
is0242: clau|AEV94557.1 ldam|AH70\_07985 pent|YP\_805018  
is0243: clau|AEV94558.1 ldam|AH70\_07990 pent|YP\_805017  
is0244: clau|AEV94562.1 ldam|AH70\_07390 pent|YP\_804022  
is0245: clau|AEV94563.1 ldam|AH70\_07385 pent|YP\_804023  
is0246: clau|AEV94564.1 ldam|AH70\_07380 pent|YP\_804024  
is0247: clau|AEV94565.1 ldam|AH70\_07375 pent|YP\_804025  
is0248: clau|AEV94566.1 ldam|AH70\_07370 pent|YP\_804026  
is0249: clau|AEV94588.1 ldam|AH70\_07995 pent|YP\_805016  
is0250: clau|AEV94589.1 ldam|AH70\_08000 pent|YP\_805015  
is0251: clau|AEV94590.1 ldam|AH70\_08005 pent|YP\_805014  
is0252: clau|AEV94591.1 ldam|AH70\_10055 pent|YP\_805013  
is0253: clau|AEV94592.1 ldam|AH70\_10050 pent|YP\_805012  
is0254: clau|AEV94593.1 ldam|AH70\_10045 pent|YP\_805011  
is0255: clau|AEV94594.1 ldam|AH70\_10040 pent|YP\_805010  
is0256: clau|AEV94595.1 ldam|AH70\_10035 pent|YP\_805009  
is0257: clau|AEV94596.1 ldam|AH70\_10030 pent|YP\_805008  
is0258: clau|AEV94597.1 ldam|AH70\_10025 pent|YP\_805007  
is0259: clau|AEV94598.1 ldam|AH70\_10020 pent|YP\_805006  
is0260: clau|AEV94599.1 ldam|AH70\_10015 pent|YP\_805005  
is0261: clau|AEV94600.1 ldam|AH70\_10840 pent|YP\_805004  
is0262: clau|AEV94605.1 ldam|AH70\_03345 pent|YP\_805082  
is0263: clau|AEV94609.1 ldam|AH70\_01490 pent|YP\_805141  
is0264: clau|AEV94614.1 ldam|AH70\_02540 pent|YP\_803582  
is0265: clau|AEV94621.1 ldam|AH70\_10800 pent|YP\_805001  
is0266: clau|AEV94622.1 ldam|AH70\_10450 pent|YP\_805000  
is0267: clau|AEV94623.1 ldam|AH70\_10455 pent|YP\_804999  
is0268: clau|AEV94625.1 ldam|AH70\_05540 pent|YP\_805164  
is0269: clau|AEV94629.1 ldam|AH70\_10460 pent|YP\_804998  
is0270: clau|AEV94630.1 ldam|AH70\_10465 pent|YP\_804997  
is0271: clau|AEV94631.1 ldam|AH70\_10470 pent|YP\_804996  
is0272: clau|AEV94641.1 ldam|AH70\_10485 pent|YP\_804993  
is0273: clau|AEV94642.1 ldam|AH70\_10490 pent|YP\_804992  
is0274: clau|AEV94651.1 ldam|AH70\_10510 pent|YP\_804991  
is0275: clau|AEV94652.1 ldam|AH70\_10515 pent|YP\_804990  
is0276: clau|AEV94653.1 ldam|AH70\_10520 pent|YP\_804989  
is0277: clau|AEV94654.1 ldam|AH70\_10525 pent|YP\_804988  
is0278: clau|AEV94655.1 ldam|AH70\_10505 pent|YP\_804987  
is0279: clau|AEV94657.1 ldam|AH70\_10540 pent|YP\_804986  
is0280: clau|AEV94660.1 ldam|AH70\_03190 pent|YP\_803634  
is0281: clau|AEV94673.1 ldam|AH70\_10545 pent|YP\_804984  
is0282: clau|AEV94678.1 ldam|AH70\_10585 pent|YP\_804983  
is0283: clau|AEV94680.1 ldam|AH70\_10590 pent|YP\_804980  
is0284: clau|AEV94681.1 ldam|AH70\_10595 pent|YP\_804979  
is0285: clau|AEV94682.1 ldam|AH70\_10600 pent|YP\_804978  
is0286: clau|AEV94683.1 ldam|AH70\_10605 pent|YP\_804977  
is0287: clau|AEV94684.1 ldam|AH70\_10610 pent|YP\_804976  
is0288: clau|AEV94685.1 ldam|AH70\_10615 pent|YP\_804975  
is0289: clau|AEV94686.1 ldam|AH70\_10620 pent|YP\_804974  
is0290: clau|AEV94688.1 ldam|AH70\_09585 pent|YP\_803876  
is0291: clau|AEV94690.1 ldam|AH70\_09330 pent|YP\_803878  
is0292: clau|AEV94693.1 ldam|AH70\_09295 pent|YP\_803879  
is0293: clau|AEV94695.1 ldam|AH70\_09290 pent|YP\_803880  
is0294: clau|AEV94696.1 ldam|AH70\_01950 pent|YP\_803883  
is0295: clau|AEV94697.1 ldam|AH70\_01955 pent|YP\_803882  
is0296: clau|AEV94698.1 ldam|AH70\_01970 pent|YP\_803881  
is0297: clau|AEV94700.1 ldam|AH70\_01945 pent|YP\_803884  
is0298: clau|AEV94701.1 ldam|AH70\_01910 pent|YP\_803885  
is0299: clau|AEV94704.1 ldam|AH70\_01705 pent|YP\_805226  
is0300: clau|AEV94711.1 ldam|AH70\_01480 pent|YP\_805225  
is0301: clau|AEV94719.1 ldam|AH70\_01110 pent|YP\_803769  
is0302: clau|AEV94723.1 ldam|AH70\_09640 pent|YP\_805077  
is0303: clau|AEV94728.1 ldam|AH70\_05310 pent|YP\_805076  
is0304: clau|AEV94729.1 ldam|AH70\_05305 pent|YP\_805075  
is0305: clau|AEV94730.1 ldam|AH70\_09710 pent|YP\_805074  
is0306: clau|AEV94731.1 ldam|AH70\_08350 pent|YP\_805073  
is0307: clau|AEV94732.1 ldam|AH70\_02125 pent|YP\_805070  
is0308: clau|AEV94733.1 ldam|AH70\_02130 pent|YP\_805069  
is0309: clau|AEV94741.1 ldam|AH70\_07545 pent|YP\_803839  
is0310: clau|AEV94742.1 ldam|AH70\_07540 pent|YP\_803840  
is0311: clau|AEV94743.1 ldam|AH70\_07535 pent|YP\_803841  
is0312: clau|AEV94746.1 ldam|AH70\_07510 pent|YP\_803843  
is0313: clau|AEV94747.1 ldam|AH70\_07505 pent|YP\_803844  
is0314: clau|AEV94750.1 ldam|AH70\_10725 pent|YP\_803863

is0315: clau|AEV94752.1 ldam|AH70\_06440 pent|YP\_803868  
is0316: clau|AEV94756.1 ldam|AH70\_06415 pent|YP\_803873  
is0317: clau|AEV94757.1 ldam|AH70\_04265 pent|YP\_803671  
is0318: clau|AEV94758.1 ldam|AH70\_04260 pent|YP\_803670  
is0319: clau|AEV94759.1 ldam|AH70\_02330 pent|YP\_805212  
is0320: clau|AEV94764.1 ldam|AH70\_09605 pent|YP\_803874  
is0321: clau|AEV94765.1 ldam|AH70\_09600 pent|YP\_803875  
is0322: clau|AEV94768.1 ldam|AH70\_10625 pent|YP\_804973  
is0323: clau|AEV94769.1 ldam|AH70\_10630 pent|YP\_804972  
is0324: clau|AEV94770.1 ldam|AH70\_10635 pent|YP\_804971  
is0325: clau|AEV94771.1 ldam|AH70\_10640 pent|YP\_804970  
is0326: clau|AEV94772.1 ldam|AH70\_04750 pent|YP\_804968  
is0327: clau|AEV94773.1 ldam|AH70\_04755 pent|YP\_804967  
is0328: clau|AEV94774.1 ldam|AH70\_04760 pent|YP\_804966  
is0329: clau|AEV94775.1 ldam|AH70\_04765 pent|YP\_804965  
is0330: clau|AEV94776.1 ldam|AH70\_04770 pent|YP\_804964  
is0331: clau|AEV94777.1 ldam|AH70\_04775 pent|YP\_804963  
is0332: clau|AEV94779.1 ldam|AH70\_04170 pent|YP\_805138  
is0333: clau|AEV94781.1 ldam|AH70\_07185 pent|YP\_803851  
is0334: clau|AEV94782.1 ldam|AH70\_07180 pent|YP\_803852  
is0335: clau|AEV94783.1 ldam|AH70\_07175 pent|YP\_803853  
is0336: clau|AEV94784.1 ldam|AH70\_07170 pent|YP\_803854  
is0337: clau|AEV94786.1 ldam|AH70\_06525 pent|YP\_805147  
is0338: clau|AEV94791.1 ldam|AH70\_04795 pent|YP\_804962  
is0339: clau|AEV94793.1 ldam|AH70\_04810 pent|YP\_804960  
is0340: clau|AEV94797.1 ldam|AH70\_04820 pent|YP\_804958  
is0341: clau|AEV94798.1 ldam|AH70\_04825 pent|YP\_804957  
is0342: clau|AEV94799.1 ldam|AH70\_04830 pent|YP\_804956  
is0343: clau|AEV94800.1 ldam|AH70\_04835 pent|YP\_804955  
is0344: clau|AEV94801.1 ldam|AH70\_04840 pent|YP\_804954  
is0345: clau|AEV94803.1 ldam|AH70\_04850 pent|YP\_804952  
is0346: clau|AEV94804.1 ldam|AH70\_04855 pent|YP\_804951  
is0347: clau|AEV94805.1 ldam|AH70\_04860 pent|YP\_804950  
is0348: clau|AEV94806.1 ldam|AH70\_04865 pent|YP\_804949  
is0349: clau|AEV94808.1 ldam|AH70\_04875 pent|YP\_804947  
is0350: clau|AEV94809.1 ldam|AH70\_04880 pent|YP\_804946  
is0351: clau|AEV94810.1 ldam|AH70\_04885 pent|YP\_804945  
is0352: clau|AEV94811.1 ldam|AH70\_04890 pent|YP\_804944  
is0353: clau|AEV94812.1 ldam|AH70\_01730 pent|YP\_803957  
is0354: clau|AEV94813.1 ldam|AH70\_01735 pent|YP\_803958  
is0355: clau|AEV94814.1 ldam|AH70\_01740 pent|YP\_803959  
is0356: clau|AEV94815.1 ldam|AH70\_09845 pent|YP\_803960  
is0357: clau|AEV94816.1 ldam|AH70\_09850 pent|YP\_803961  
is0358: clau|AEV94817.1 ldam|AH70\_09855 pent|YP\_803962  
is0359: clau|AEV94818.1 ldam|AH70\_09860 pent|YP\_803963  
is0360: clau|AEV94819.1 ldam|AH70\_09865 pent|YP\_803964  
is0361: clau|AEV94820.1 ldam|AH70\_09870 pent|YP\_803965  
is0362: clau|AEV94822.1 ldam|AH70\_09880 pent|YP\_803968  
is0363: clau|AEV94823.1 ldam|AH70\_09885 pent|YP\_803969  
is0364: clau|AEV94824.1 ldam|AH70\_09890 pent|YP\_803970  
is0365: clau|AEV94825.1 ldam|AH70\_09895 pent|YP\_803971  
is0366: clau|AEV94826.1 ldam|AH70\_09900 pent|YP\_803972  
is0367: clau|AEV94827.1 ldam|AH70\_09905 pent|YP\_803973  
is0368: clau|AEV94828.1 ldam|AH70\_09910 pent|YP\_803974  
is0369: clau|AEV94829.1 ldam|AH70\_09915 pent|YP\_803975  
is0370: clau|AEV94830.1 ldam|AH70\_09920 pent|YP\_803976  
is0371: clau|AEV94831.1 ldam|AH70\_09925 pent|YP\_803977  
is0372: clau|AEV94833.1 ldam|AH70\_09935 pent|YP\_803979  
is0373: clau|AEV94834.1 ldam|AH70\_09940 pent|YP\_803980  
is0374: clau|AEV94835.1 ldam|AH70\_09945 pent|YP\_803981  
is0375: clau|AEV94836.1 ldam|AH70\_09950 pent|YP\_803982  
is0376: clau|AEV94837.1 ldam|AH70\_09955 pent|YP\_803983  
is0377: clau|AEV94839.1 ldam|AH70\_09960 pent|YP\_803984  
is0378: clau|AEV94840.1 ldam|AH70\_09970 pent|YP\_803986  
is0379: clau|AEV94841.1 ldam|AH70\_04645 pent|YP\_803987  
is0380: clau|AEV94842.1 ldam|AH70\_09985 pent|YP\_803988  
is0381: clau|AEV94843.1 ldam|AH70\_09990 pent|YP\_803989  
is0382: clau|AEV94844.1 ldam|AH70\_09995 pent|YP\_803990  
is0383: clau|AEV94845.1 ldam|AH70\_07740 pent|YP\_803993  
is0384: clau|AEV94846.1 ldam|AH70\_07730 pent|YP\_803994  
is0385: clau|AEV94847.1 ldam|AH70\_07725 pent|YP\_803995  
is0386: clau|AEV94848.1 ldam|AH70\_07720 pent|YP\_803996  
is0387: clau|AEV94849.1 ldam|AH70\_07715 pent|YP\_803997  
is0388: clau|AEV94850.1 ldam|AH70\_07710 pent|YP\_803998  
is0389: clau|AEV94851.1 ldam|AH70\_07705 pent|YP\_803999  
is0390: clau|AEV94853.1 ldam|AH70\_07690 pent|YP\_804000  
is0391: clau|AEV94854.1 ldam|AH70\_07685 pent|YP\_804001  
is0392: clau|AEV94855.1 ldam|AH70\_07680 pent|YP\_804002  
is0393: clau|AEV94861.1 ldam|AH70\_07670 pent|YP\_804003  
is0394: clau|AEV94862.1 ldam|AH70\_07665 pent|YP\_804004  
is0395: clau|AEV94863.1 ldam|AH70\_07660 pent|YP\_804005  
is0396: clau|AEV94864.1 ldam|AH70\_07655 pent|YP\_804006  
is0397: clau|AEV94865.1 ldam|AH70\_07650 pent|YP\_804007

is0398: clau|AEV94866.1 ldam|AH70\_07645 pent|YP\_804008  
is0399: clau|AEV94867.1 ldam|AH70\_07640 pent|YP\_804009  
is0400: clau|AEV94868.1 ldam|AH70\_07635 pent|YP\_804010  
is0401: clau|AEV94869.1 ldam|AH70\_07630 pent|YP\_804011  
is0402: clau|AEV94870.1 ldam|AH70\_03380 pent|YP\_804012  
is0403: clau|AEV94871.1 ldam|AH70\_03375 pent|YP\_804013  
is0404: clau|AEV94872.1 ldam|AH70\_03370 pent|YP\_804014  
is0405: clau|AEV94873.1 ldam|AH70\_03365 pent|YP\_804015  
is0406: clau|AEV94874.1 ldam|AH70\_03355 pent|YP\_804016  
is0407: clau|AEV94882.1 ldam|AH70\_03305 pent|YP\_804073  
is0408: clau|AEV94895.1 ldam|AH70\_09455 pent|YP\_804503  
is0409: clau|AEV94897.1 ldam|AH70\_09445 pent|YP\_804501  
is0410: clau|AEV94898.1 ldam|AH70\_09435 pent|YP\_804499  
is0411: clau|AEV94899.1 ldam|AH70\_09430 pent|YP\_804498  
is0412: clau|AEV94907.1 ldam|AH70\_03250 pent|YP\_804835  
is0413: clau|AEV94908.1 ldam|AH70\_03245 pent|YP\_804834  
is0414: clau|AEV94911.1 ldam|AH70\_03240 pent|YP\_804833  
is0415: clau|AEV94912.1 ldam|AH70\_03235 pent|YP\_804832  
is0416: clau|AEV94923.1 ldam|AH70\_03230 pent|YP\_804831  
is0417: clau|AEV94924.1 ldam|AH70\_07405 pent|YP\_804020  
is0418: clau|AEV94925.1 ldam|AH70\_07400 pent|YP\_804021  
is0419: clau|AEV94926.1 ldam|AH70\_08595 pent|YP\_803643  
is0420: clau|AEV94928.1 ldam|AH70\_01805 pent|YP\_804102  
is0421: clau|AEV94934.1 ldam|AH70\_05290 pent|YP\_803763  
is0422: clau|AEV94937.1 ldam|AH70\_03200 pent|YP\_804822  
is0423: clau|AEV94938.1 ldam|AH70\_03160 pent|YP\_804821  
is0424: clau|AEV94939.1 ldam|AH70\_03155 pent|YP\_804820  
is0425: clau|AEV94940.1 ldam|AH70\_00760 pent|YP\_804819  
is0426: clau|AEV94941.1 ldam|AH70\_00765 pent|YP\_804818  
is0427: clau|AEV94942.1 ldam|AH70\_00770 pent|YP\_804817  
is0428: clau|AEV94943.1 ldam|AH70\_01885 pent|YP\_803767  
is0429: clau|AEV94944.1 ldam|AH70\_01880 pent|YP\_803768  
is0430: clau|AEV94945.1 ldam|AH70\_00775 pent|YP\_804816  
is0431: clau|AEV94946.1 ldam|AH70\_00780 pent|YP\_804815  
is0432: clau|AEV94947.1 ldam|AH70\_00785 pent|YP\_804814  
is0433: clau|AEV94948.1 ldam|AH70\_00790 pent|YP\_804813  
is0434: clau|AEV94949.1 ldam|AH70\_00795 pent|YP\_804812  
is0435: clau|AEV94950.1 ldam|AH70\_00800 pent|YP\_804811  
is0436: clau|AEV94951.1 ldam|AH70\_00805 pent|YP\_804810  
is0437: clau|AEV94952.1 ldam|AH70\_00810 pent|YP\_804809  
is0438: clau|AEV94953.1 ldam|AH70\_02280 pent|YP\_804806  
is0439: clau|AEV94954.1 ldam|AH70\_02285 pent|YP\_804805  
is0440: clau|AEV94955.1 ldam|AH70\_02290 pent|YP\_804804  
is0441: clau|AEV94956.1 ldam|AH70\_02295 pent|YP\_804803  
is0442: clau|AEV94957.1 ldam|AH70\_02300 pent|YP\_804802  
is0443: clau|AEV94958.1 ldam|AH70\_02305 pent|YP\_804801  
is0444: clau|AEV94959.1 ldam|AH70\_02310 pent|YP\_804800  
is0445: clau|AEV94960.1 ldam|AH70\_02315 pent|YP\_804799  
is0446: clau|AEV94961.1 ldam|AH70\_06790 pent|YP\_804798  
is0447: clau|AEV94963.1 ldam|AH70\_06775 pent|YP\_804796  
is0448: clau|AEV94964.1 ldam|AH70\_06770 pent|YP\_804795  
is0449: clau|AEV94965.1 ldam|AH70\_06765 pent|YP\_804794  
is0450: clau|AEV94966.1 ldam|AH70\_06760 pent|YP\_804793  
is0451: clau|AEV94967.1 ldam|AH70\_06755 pent|YP\_804792  
is0452: clau|AEV94969.1 ldam|AH70\_06735 pent|YP\_804791  
is0453: clau|AEV94970.1 ldam|AH70\_06730 pent|YP\_804790  
is0454: clau|AEV94971.1 ldam|AH70\_06725 pent|YP\_804789  
is0455: clau|AEV94972.1 ldam|AH70\_06720 pent|YP\_804788  
is0456: clau|AEV94974.1 ldam|AH70\_06715 pent|YP\_804786  
is0457: clau|AEV94975.1 ldam|AH70\_06710 pent|YP\_805229  
is0458: clau|AEV94976.1 ldam|AH70\_06705 pent|YP\_804785  
is0459: clau|AEV94977.1 ldam|AH70\_06695 pent|YP\_804784  
is0460: clau|AEV94978.1 ldam|AH70\_06690 pent|YP\_804783  
is0461: clau|AEV94979.1 ldam|AH70\_06680 pent|YP\_804781  
is0462: clau|AEV94980.1 ldam|AH70\_06665 pent|YP\_804780  
is0463: clau|AEV94981.1 ldam|AH70\_06660 pent|YP\_804779  
is0464: clau|AEV94982.1 ldam|AH70\_06655 pent|YP\_804778  
is0465: clau|AEV94983.1 ldam|AH70\_06650 pent|YP\_804777  
is0466: clau|AEV94984.1 ldam|AH70\_06645 pent|YP\_804776  
is0467: clau|AEV94985.1 ldam|AH70\_06640 pent|YP\_804775  
is0468: clau|AEV94986.1 ldam|AH70\_06635 pent|YP\_804774  
is0469: clau|AEV94987.1 ldam|AH70\_06630 pent|YP\_804773  
is0470: clau|AEV94988.1 ldam|AH70\_06625 pent|YP\_804772  
is0471: clau|AEV94989.1 ldam|AH70\_06620 pent|YP\_804771  
is0472: clau|AEV94990.1 ldam|AH70\_06615 pent|YP\_804770  
is0473: clau|AEV94991.1 ldam|AH70\_03920 pent|YP\_804769  
is0474: clau|AEV94992.1 ldam|AH70\_03925 pent|YP\_804768  
is0475: clau|AEV94993.1 ldam|AH70\_03930 pent|YP\_804767  
is0476: clau|AEV94995.1 ldam|AH70\_03935 pent|YP\_804765  
is0477: clau|AEV94996.1 ldam|AH70\_03940 pent|YP\_804764  
is0478: clau|AEV94997.1 ldam|AH70\_03945 pent|YP\_804763  
is0479: clau|AEV94999.1 ldam|AH70\_03955 pent|YP\_804761  
is0480: clau|AEV95000.1 ldam|AH70\_03960 pent|YP\_804760

is0481: clau|AEV95001.1 ldam|AH70\_03965 pent|YP\_804759  
is0482: clau|AEV95002.1 ldam|AH70\_03970 pent|YP\_804758  
is0483: clau|AEV95003.1 ldam|AH70\_03975 pent|YP\_804757  
is0484: clau|AEV95004.1 ldam|AH70\_03980 pent|YP\_804756  
is0485: clau|AEV95005.1 ldam|AH70\_03985 pent|YP\_804755  
is0486: clau|AEV95006.1 ldam|AH70\_03990 pent|YP\_804754  
is0487: clau|AEV95008.1 ldam|AH70\_03995 pent|YP\_804752  
is0488: clau|AEV95009.1 ldam|AH70\_04000 pent|YP\_804751  
is0489: clau|AEV95010.1 ldam|AH70\_04005 pent|YP\_804750  
is0490: clau|AEV95011.1 ldam|AH70\_04010 pent|YP\_804749  
is0491: clau|AEV95012.1 ldam|AH70\_10850 pent|YP\_804748  
is0492: clau|AEV95013.1 ldam|AH70\_10855 pent|YP\_804747  
is0493: clau|AEV95014.1 ldam|AH70\_10860 pent|YP\_804746  
is0494: clau|AEV95015.1 ldam|AH70\_10865 pent|YP\_804745  
is0495: clau|AEV95016.1 ldam|AH70\_10870 pent|YP\_804744  
is0496: clau|AEV95017.1 ldam|AH70\_10875 pent|YP\_804743  
is0497: clau|AEV95018.1 ldam|AH70\_10880 pent|YP\_804742  
is0498: clau|AEV95019.1 ldam|AH70\_10885 pent|YP\_804741  
is0499: clau|AEV95020.1 ldam|AH70\_02010 pent|YP\_804740  
is0500: clau|AEV95021.1 ldam|AH70\_02005 pent|YP\_804739  
is0501: clau|AEV95022.1 ldam|AH70\_02000 pent|YP\_804738  
is0502: clau|AEV95024.1 ldam|AH70\_01990 pent|YP\_804736  
is0503: clau|AEV95026.1 ldam|AH70\_08375 pent|YP\_804734  
is0504: clau|AEV95027.1 ldam|AH70\_08380 pent|YP\_804733  
is0505: clau|AEV95031.1 ldam|AH70\_03215 pent|YP\_804826  
is0506: clau|AEV95032.1 ldam|AH70\_10175 pent|YP\_803590  
is0507: clau|AEV95033.1 ldam|AH70\_10720 pent|YP\_804727  
is0508: clau|AEV95035.1 ldam|AH70\_02200 pent|YP\_804723  
is0509: clau|AEV95037.1 ldam|AH70\_02205 pent|YP\_804721  
is0510: clau|AEV95038.1 ldam|AH70\_02210 pent|YP\_804720  
is0511: clau|AEV95039.1 ldam|AH70\_02215 pent|YP\_804719  
is0512: clau|AEV95040.1 ldam|AH70\_02220 pent|YP\_804718  
is0513: clau|AEV95042.1 ldam|AH70\_02230 pent|YP\_804716  
is0514: clau|AEV95044.1 ldam|AH70\_02240 pent|YP\_804714  
is0515: clau|AEV95045.1 ldam|AH70\_02245 pent|YP\_804713  
is0516: clau|AEV95046.1 ldam|AH70\_02250 pent|YP\_804712  
is0517: clau|AEV95047.1 ldam|AH70\_02255 pent|YP\_804711  
is0518: clau|AEV95048.1 ldam|AH70\_02260 pent|YP\_804710  
is0519: clau|AEV95049.1 ldam|AH70\_02265 pent|YP\_804709  
is0520: clau|AEV95050.1 ldam|AH70\_02270 pent|YP\_804708  
is0521: clau|AEV95051.1 ldam|AH70\_02275 pent|YP\_804707  
is0522: clau|AEV95054.1 ldam|AH70\_08985 pent|YP\_804705  
is0523: clau|AEV95055.1 ldam|AH70\_08990 pent|YP\_804704  
is0524: clau|AEV95056.1 ldam|AH70\_08995 pent|YP\_804703  
is0525: clau|AEV95057.1 ldam|AH70\_09000 pent|YP\_804702  
is0526: clau|AEV95060.1 ldam|AH70\_09015 pent|YP\_804700  
is0527: clau|AEV95061.1 ldam|AH70\_09020 pent|YP\_804699  
is0528: clau|AEV95062.1 ldam|AH70\_09025 pent|YP\_804698  
is0529: clau|AEV95063.1 ldam|AH70\_09030 pent|YP\_804697  
is0530: clau|AEV95064.1 ldam|AH70\_09035 pent|YP\_804696  
is0531: clau|AEV95065.1 ldam|AH70\_09315 pent|YP\_804730  
is0532: clau|AEV95066.1 ldam|AH70\_09310 pent|YP\_804729  
is0533: clau|AEV95067.1 ldam|AH70\_09040 pent|YP\_804695  
is0534: clau|AEV95068.1 ldam|AH70\_09045 pent|YP\_804694  
is0535: clau|AEV95069.1 ldam|AH70\_09050 pent|YP\_804693  
is0536: clau|AEV95070.1 ldam|AH70\_09055 pent|YP\_804692  
is0537: clau|AEV95071.1 ldam|AH70\_09065 pent|YP\_804691  
is0538: clau|AEV95072.1 ldam|AH70\_09070 pent|YP\_804690  
is0539: clau|AEV95073.1 ldam|AH70\_09080 pent|YP\_804688  
is0540: clau|AEV95074.1 ldam|AH70\_09085 pent|YP\_804687  
is0541: clau|AEV95075.1 ldam|AH70\_09090 pent|YP\_804686  
is0542: clau|AEV95076.1 ldam|AH70\_09095 pent|YP\_804685  
is0543: clau|AEV95077.1 ldam|AH70\_09100 pent|YP\_804684  
is0544: clau|AEV95078.1 ldam|AH70\_09105 pent|YP\_804683  
is0545: clau|AEV95079.1 ldam|AH70\_09110 pent|YP\_804682  
is0546: clau|AEV95080.1 ldam|AH70\_09115 pent|YP\_804681  
is0547: clau|AEV95081.1 ldam|AH70\_09120 pent|YP\_804680  
is0548: clau|AEV95082.1 ldam|AH70\_09125 pent|YP\_804679  
is0549: clau|AEV95083.1 ldam|AH70\_09135 pent|YP\_804678  
is0550: clau|AEV95084.1 ldam|AH70\_09140 pent|YP\_804677  
is0551: clau|AEV95085.1 ldam|AH70\_09145 pent|YP\_804676  
is0552: clau|AEV95086.1 ldam|AH70\_09150 pent|YP\_804675  
is0553: clau|AEV95088.1 ldam|AH70\_09160 pent|YP\_804673  
is0554: clau|AEV95089.1 ldam|AH70\_09165 pent|YP\_804672  
is0555: clau|AEV95090.1 ldam|AH70\_09170 pent|YP\_804671  
is0556: clau|AEV95091.1 ldam|AH70\_09175 pent|YP\_804670  
is0557: clau|AEV95092.1 ldam|AH70\_09180 pent|YP\_804669  
is0558: clau|AEV95093.1 ldam|AH70\_09185 pent|YP\_804668  
is0559: clau|AEV95094.1 ldam|AH70\_09190 pent|YP\_804667  
is0560: clau|AEV95095.1 ldam|AH70\_09195 pent|YP\_804666  
is0561: clau|AEV95096.1 ldam|AH70\_09200 pent|YP\_804665  
is0562: clau|AEV95097.1 ldam|AH70\_09205 pent|YP\_804664  
is0563: clau|AEV95098.1 ldam|AH70\_09210 pent|YP\_804663



is0647:	clau	AEV95212.1	ldam	AH70_08850	pent	YP_804446
is0648:	clau	AEV95214.1	ldam	AH70_08855	pent	YP_804444
is0649:	clau	AEV95215.1	ldam	AH70_05000	pent	YP_804598
is0650:	clau	AEV95216.1	ldam	AH70_04995	pent	YP_804599
is0651:	clau	AEV95219.1	ldam	AH70_04980	pent	YP_804602
is0652:	clau	AEV95220.1	ldam	AH70_04975	pent	YP_804603
is0653:	clau	AEV95221.1	ldam	AH70_04970	pent	YP_804604
is0654:	clau	AEV95222.1	ldam	AH70_04965	pent	YP_804605
is0655:	clau	AEV95223.1	ldam	AH70_04960	pent	YP_804606
is0656:	clau	AEV95224.1	ldam	AH70_04955	pent	YP_804607
is0657:	clau	AEV95226.1	ldam	AH70_02765	pent	YP_804610
is0658:	clau	AEV95227.1	ldam	AH70_02770	pent	YP_804611
is0659:	clau	AEV95228.1	ldam	AH70_02780	pent	YP_804612
is0660:	clau	AEV95229.1	ldam	AH70_02785	pent	YP_804613
is0661:	clau	AEV95230.1	ldam	AH70_02790	pent	YP_804614
is0662:	clau	AEV95231.1	ldam	AH70_02795	pent	YP_804615
is0663:	clau	AEV95232.1	ldam	AH70_02800	pent	YP_804616
is0664:	clau	AEV95234.1	ldam	AH70_02820	pent	YP_804619
is0665:	clau	AEV95235.1	ldam	AH70_02825	pent	YP_804620
is0666:	clau	AEV95236.1	ldam	AH70_10435	pent	YP_804621
is0667:	clau	AEV95237.1	ldam	AH70_10430	pent	YP_804622
is0668:	clau	AEV95241.1	ldam	AH70_10420	pent	YP_804398
is0669:	clau	AEV95242.1	ldam	AH70_10415	pent	YP_804397
is0670:	clau	AEV95243.1	ldam	AH70_10410	pent	YP_804396
is0671:	clau	AEV95244.1	ldam	AH70_10405	pent	YP_804395
is0672:	clau	AEV95245.1	ldam	AH70_10400	pent	YP_804394
is0673:	clau	AEV95246.1	ldam	AH70_10395	pent	YP_804393
is0674:	clau	AEV95247.1	ldam	AH70_10390	pent	YP_804392
is0675:	clau	AEV95248.1	ldam	AH70_10385	pent	YP_804391
is0676:	clau	AEV95249.1	ldam	AH70_10380	pent	YP_804390
is0677:	clau	AEV95250.1	ldam	AH70_10375	pent	YP_804389
is0678:	clau	AEV95251.1	ldam	AH70_10370	pent	YP_804388
is0679:	clau	AEV95252.1	ldam	AH70_10365	pent	YP_804387
is0680:	clau	AEV95253.1	ldam	AH70_10360	pent	YP_804386
is0681:	clau	AEV95254.1	ldam	AH70_10355	pent	YP_804385
is0682:	clau	AEV95255.1	ldam	AH70_10350	pent	YP_804384
is0683:	clau	AEV95256.1	ldam	AH70_10345	pent	YP_804383
is0684:	clau	AEV95257.1	ldam	AH70_10340	pent	YP_804382
is0685:	clau	AEV95258.1	ldam	AH70_10335	pent	YP_804381
is0686:	clau	AEV95259.1	ldam	AH70_10330	pent	YP_804380
is0687:	clau	AEV95260.1	ldam	AH70_10325	pent	YP_804379
is0688:	clau	AEV95261.1	ldam	AH70_10320	pent	YP_804378
is0689:	clau	AEV95262.1	ldam	AH70_05660	pent	YP_804377
is0690:	clau	AEV95264.1	ldam	AH70_05670	pent	YP_804375
is0691:	clau	AEV95266.1	ldam	AH70_05690	pent	YP_804373
is0692:	clau	AEV95267.1	ldam	AH70_05695	pent	YP_804372
is0693:	clau	AEV95268.1	ldam	AH70_05700	pent	YP_804371
is0694:	clau	AEV95269.1	ldam	AH70_05710	pent	YP_804160
is0695:	clau	AEV95270.1	ldam	AH70_05715	pent	YP_804370
is0696:	clau	AEV95271.1	ldam	AH70_05720	pent	YP_804369
is0697:	clau	AEV95273.1	ldam	AH70_08860	pent	YP_804443
is0698:	clau	AEV95274.1	ldam	AH70_08865	pent	YP_804442
is0699:	clau	AEV95275.1	ldam	AH70_08870	pent	YP_804441
is0700:	clau	AEV95276.1	ldam	AH70_08875	pent	YP_804440
is0701:	clau	AEV95277.1	ldam	AH7		

is0730: clau|AEV95306.1 ldam|AH70\_03555 pent|YP\_804344  
is0731: clau|AEV95307.1 ldam|AH70\_03550 pent|YP\_804343  
is0732: clau|AEV95308.1 ldam|AH70\_03545 pent|YP\_804342  
is0733: clau|AEV95309.1 ldam|AH70\_03540 pent|YP\_804341  
is0734: clau|AEV95310.1 ldam|AH70\_03535 pent|YP\_804340  
is0735: clau|AEV95311.1 ldam|AH70\_03530 pent|YP\_804339  
is0736: clau|AEV95312.1 ldam|AH70\_03525 pent|YP\_804338  
is0737: clau|AEV95313.1 ldam|AH70\_00250 pent|YP\_804337  
is0738: clau|AEV95314.1 ldam|AH70\_00245 pent|YP\_804336  
is0739: clau|AEV95315.1 ldam|AH70\_00240 pent|YP\_804335  
is0740: clau|AEV95316.1 ldam|AH70\_00235 pent|YP\_804334  
is0741: clau|AEV95317.1 ldam|AH70\_00230 pent|YP\_804333  
is0742: clau|AEV95318.1 ldam|AH70\_00225 pent|YP\_804332  
is0743: clau|AEV95319.1 ldam|AH70\_00220 pent|YP\_804331  
is0744: clau|AEV95320.1 ldam|AH70\_00215 pent|YP\_804330  
is0745: clau|AEV95321.1 ldam|AH70\_00210 pent|YP\_804329  
is0746: clau|AEV95322.1 ldam|AH70\_00205 pent|YP\_804328  
is0747: clau|AEV95323.1 ldam|AH70\_00200 pent|YP\_804327  
is0748: clau|AEV95324.1 ldam|AH70\_00195 pent|YP\_804326  
is0749: clau|AEV95326.1 ldam|AH70\_00165 pent|YP\_804324  
is0750: clau|AEV95327.1 ldam|AH70\_00160 pent|YP\_804323  
is0751: clau|AEV95328.1 ldam|AH70\_00155 pent|YP\_804322  
is0752: clau|AEV95329.1 ldam|AH70\_00150 pent|YP\_804321  
is0753: clau|AEV95330.1 ldam|AH70\_00145 pent|YP\_804320  
is0754: clau|AEV95331.1 ldam|AH70\_00140 pent|YP\_804319  
is0755: clau|AEV95332.1 ldam|AH70\_11165 pent|YP\_804318  
is0756: clau|AEV95333.1 ldam|AH70\_11160 pent|YP\_804317  
is0757: clau|AEV95334.1 ldam|AH70\_11155 pent|YP\_804316  
is0758: clau|AEV95335.1 ldam|AH70\_11150 pent|YP\_804315  
is0759: clau|AEV95336.1 ldam|AH70\_11145 pent|YP\_804314  
is0760: clau|AEV95337.1 ldam|AH70\_11140 pent|YP\_804313  
is0761: clau|AEV95339.1 ldam|AH70\_03285 pent|YP\_804048  
is0762: clau|AEV95346.1 ldam|AH70\_11110 pent|YP\_804311  
is0763: clau|AEV95353.1 ldam|AH70\_08215 pent|YP\_804245  
is0764: clau|AEV95354.1 ldam|AH70\_08210 pent|YP\_804244  
is0765: clau|AEV95355.1 ldam|AH70\_08205 pent|YP\_804243  
is0766: clau|AEV95356.1 ldam|AH70\_08200 pent|YP\_804242  
is0767: clau|AEV95357.1 ldam|AH70\_08195 pent|YP\_804241  
is0768: clau|AEV95358.1 ldam|AH70\_08190 pent|YP\_804240  
is0769: clau|AEV95359.1 ldam|AH70\_08185 pent|YP\_804239  
is0770: clau|AEV95360.1 ldam|AH70\_08180 pent|YP\_804238  
is0771: clau|AEV95361.1 ldam|AH70\_08175 pent|YP\_804237  
is0772: clau|AEV95363.1 ldam|AH70\_08170 pent|YP\_804236  
is0773: clau|AEV95364.1 ldam|AH70\_08165 pent|YP\_804235  
is0774: clau|AEV95365.1 ldam|AH70\_08160 pent|YP\_804234  
is0775: clau|AEV95366.1 ldam|AH70\_08155 pent|YP\_804233  
is0776: clau|AEV95367.1 ldam|AH70\_08150 pent|YP\_804232  
is0777: clau|AEV95368.1 ldam|AH70\_08145 pent|YP\_804230  
is0778: clau|AEV95369.1 ldam|AH70\_08140 pent|YP\_804229  
is0779: clau|AEV95370.1 ldam|AH70\_08135 pent|YP\_804228  
is0780: clau|AEV95371.1 ldam|AH70\_08130 pent|YP\_804227  
is0781: clau|AEV95372.1 ldam|AH70\_08125 pent|YP\_804226  
is0782: clau|AEV95373.1 ldam|AH70\_08120 pent|YP\_804225  
is0783: clau|AEV95374.1 ldam|AH70\_08115 pent|YP\_804224  
is0784: clau|AEV95376.1 ldam|AH70\_05960 pent|YP\_804222  
is0785: clau|AEV95377.1 ldam|AH70\_05955 pent|YP\_804221  
is0786: clau|AEV95378.1 ldam|AH70\_05950 pent|YP\_804220  
is0787: clau|AEV95379.1 ldam|AH70\_05945 pent|YP\_804219  
is0788: clau|AEV95380.1 ldam|AH70\_05940 pent|YP\_804218  
is0789: clau|AEV95381.1 ldam|AH70\_05935 pent|YP\_804217  
is0790: clau|AEV95382.1 ldam|AH70\_05930 pent|YP\_804216  
is0791: clau|AEV95383.1 ldam|AH70\_05925 pent|YP\_804215  
is0792: clau|AEV95384.1 ldam|AH70\_05920 pent|YP\_804214  
is0793: clau|AEV95385.1 ldam|AH70\_05915 pent|YP\_804213  
is0794: clau|AEV95386.1 ldam|AH70\_05910 pent|YP\_804212  
is0795: clau|AEV95387.1 ldam|AH70\_05905 pent|YP\_804211  
is0796: clau|AEV95388.1 ldam|AH70\_05900 pent|YP\_804210  
is0797: clau|AEV95389.1 ldam|AH70\_05895 pent|YP\_804209  
is0798: clau|AEV95390.1 ldam|AH70\_05890 pent|YP\_804208  
is0799: clau|AEV95391.1 ldam|AH70\_05885 pent|YP\_804207  
is0800: clau|AEV95392.1 ldam|AH70\_05880 pent|YP\_804206  
is0801: clau|AEV95393.1 ldam|AH70\_05875 pent|YP\_804205  
is0802: clau|AEV95395.1 ldam|AH70\_05870 pent|YP\_804204  
is0803: clau|AEV95396.1 ldam|AH70\_05865 pent|YP\_804203  
is0804: clau|AEV95397.1 ldam|AH70\_05860 pent|YP\_804202  
is0805: clau|AEV95398.1 ldam|AH70\_05855 pent|YP\_804197  
is0806: clau|AEV95399.1 ldam|AH70\_05850 pent|YP\_804196  
is0807: clau|AEV95400.1 ldam|AH70\_05845 pent|YP\_804195  
is0808: clau|AEV95401.1 ldam|AH70\_05835 pent|YP\_804194  
is0809: clau|AEV95402.1 ldam|AH70\_05830 pent|YP\_804193  
is0810: clau|AEV95403.1 ldam|AH70\_05825 pent|YP\_804192  
is0811: clau|AEV95404.1 ldam|AH70\_02590 pent|YP\_804191  
is0812: clau|AEV95406.1 ldam|AH70\_08730 pent|YP\_804190

is0813: clau|AEV95407.1 ldam|AH70\_08725 pent|YP\_804189  
is0814: clau|AEV95408.1 ldam|AH70\_08720 pent|YP\_804188  
is0815: clau|AEV95409.1 ldam|AH70\_08715 pent|YP\_804187  
is0816: clau|AEV95410.1 ldam|AH70\_08710 pent|YP\_804186  
is0817: clau|AEV95411.1 ldam|AH70\_08705 pent|YP\_804185  
is0818: clau|AEV95412.1 ldam|AH70\_08700 pent|YP\_804184  
is0819: clau|AEV95413.1 ldam|AH70\_08695 pent|YP\_804183  
is0820: clau|AEV95414.1 ldam|AH70\_08690 pent|YP\_804182  
is0821: clau|AEV95415.1 ldam|AH70\_08685 pent|YP\_804181  
is0822: clau|AEV95416.1 ldam|AH70\_08680 pent|YP\_804180  
is0823: clau|AEV95417.1 ldam|AH70\_08675 pent|YP\_804179  
is0824: clau|AEV95418.1 ldam|AH70\_08670 pent|YP\_804178  
is0825: clau|AEV95419.1 ldam|AH70\_08665 pent|YP\_804177  
is0826: clau|AEV95420.1 ldam|AH70\_08660 pent|YP\_804176  
is0827: clau|AEV95421.1 ldam|AH70\_08655 pent|YP\_804175  
is0828: clau|AEV95422.1 ldam|AH70\_08650 pent|YP\_804174  
is0829: clau|AEV95423.1 ldam|AH70\_08645 pent|YP\_804173  
is0830: clau|AEV95424.1 ldam|AH70\_08630 pent|YP\_804172  
is0831: clau|AEV95428.1 ldam|AH70\_08620 pent|YP\_804166  
is0832: clau|AEV95429.1 ldam|AH70\_08615 pent|YP\_804165  
is0833: clau|AEV95430.1 ldam|AH70\_08610 pent|YP\_804164  
is0834: clau|AEV95431.1 ldam|AH70\_08605 pent|YP\_804163  
is0835: clau|AEV95438.1 ldam|AH70\_08570 pent|YP\_804159  
is0836: clau|AEV95439.1 ldam|AH70\_08565 pent|YP\_804158  
is0837: clau|AEV95441.1 ldam|AH70\_08560 pent|YP\_804156  
is0838: clau|AEV95442.1 ldam|AH70\_08590 pent|YP\_804155  
is0839: clau|AEV95443.1 ldam|AH70\_08550 pent|YP\_804154  
is0840: clau|AEV95444.1 ldam|AH70\_08540 pent|YP\_804152  
is0841: clau|AEV95445.1 ldam|AH70\_08535 pent|YP\_804151  
is0842: clau|AEV95446.1 ldam|AH70\_08525 pent|YP\_804150  
is0843: clau|AEV95447.1 ldam|AH70\_08520 pent|YP\_804149  
is0844: clau|AEV95448.1 ldam|AH70\_08515 pent|YP\_804148  
is0845: clau|AEV95449.1 ldam|AH70\_08510 pent|YP\_804147  
is0846: clau|AEV95451.1 ldam|AH70\_08495 pent|YP\_804145  
is0847: clau|AEV95453.1 ldam|AH70\_09230 pent|YP\_804143  
is0848: clau|AEV95454.1 ldam|AH70\_09235 pent|YP\_804142  
is0849: clau|AEV95455.1 ldam|AH70\_09240 pent|YP\_804141  
is0850: clau|AEV95457.1 ldam|AH70\_09245 pent|YP\_804139  
is0851: clau|AEV95458.1 ldam|AH70\_09250 pent|YP\_804138  
is0852: clau|AEV95459.1 ldam|AH70\_09255 pent|YP\_804137  
is0853: clau|AEV95461.1 ldam|AH70\_05390 pent|YP\_804135  
is0854: clau|AEV95462.1 ldam|AH70\_05395 pent|YP\_804134  
is0855: clau|AEV95464.1 ldam|AH70\_09270 pent|YP\_804132  
is0856: clau|AEV95465.1 ldam|AH70\_11030 pent|YP\_804131  
is0857: clau|AEV95466.1 ldam|AH70\_11025 pent|YP\_804130  
is0858: clau|AEV95467.1 ldam|AH70\_11020 pent|YP\_804129  
is0859: clau|AEV95469.1 ldam|AH70\_11010 pent|YP\_804128  
is0860: clau|AEV95470.1 ldam|AH70\_11005 pent|YP\_804127  
is0861: clau|AEV95473.1 ldam|AH70\_02420 pent|YP\_804123  
is0862: clau|AEV95474.1 ldam|AH70\_02435 pent|YP\_804122  
is0863: clau|AEV95475.1 ldam|AH70\_01650 pent|YP\_804121  
is0864: clau|AEV95476.1 ldam|AH70\_00025 pent|YP\_804120  
is0865: clau|AEV95477.1 ldam|AH70\_00030 pent|YP\_804119  
is0866: clau|AEV95478.1 ldam|AH70\_00035 pent|YP\_804118  
is0867: clau|AEV95479.1 ldam|AH70\_00040 pent|YP\_804117  
is0868: clau|AEV95480.1 ldam|AH70\_00045 pent|YP\_804114  
is0869: clau|AEV95484.1 ldam|AH70\_03845 pent|YP\_804105  
is0870: clau|AEV95489.1 ldam|AH70\_01590 pent|YP\_804098  
is0871: clau|AEV95490.1 ldam|AH70\_01585 pent|YP\_804097  
is0872: clau|AEV95491.1 ldam|AH70\_01580 pent|YP\_804096  
is0873: clau|AEV95492.1 ldam|AH70\_01575 pent|YP\_804095  
is0874: clau|AEV95494.1 ldam|AH70\_00065 pent|YP\_804846  
is0875: clau|AEV95495.1 ldam|AH70\_00070 pent|YP\_804847  
is0876: clau|AEV95496.1 ldam|AH70\_00075 pent|YP\_804848  
is0877: clau|AEV95497.1 ldam|AH70\_00080 pent|YP\_804849  
is0878: clau|AEV95498.1 ldam|AH70\_00085 pent|YP\_804850  
is0879: clau|AEV95499.1 ldam|AH70\_00090 pent|YP\_804851  
is0880: clau|AEV95500.1 ldam|AH70\_00095 pent|YP\_804852  
is0881: clau|AEV95501.1 ldam|AH70\_00100 pent|YP\_804853  
is0882: clau|AEV95502.1 ldam|AH70\_00105 pent|YP\_804854  
is0883: clau|AEV95503.1 ldam|AH70\_00110 pent|YP\_804855  
is0884: clau|AEV95504.1 ldam|AH70\_00115 pent|YP\_804856  
is0885: clau|AEV95505.1 ldam|AH70\_00120 pent|YP\_804857  
is0886: clau|AEV95506.1 ldam|AH70\_06410 pent|YP\_804865  
is0887: clau|AEV95507.1 ldam|AH70\_06405 pent|YP\_804866  
is0888: clau|AEV95508.1 ldam|AH70\_06400 pent|YP\_804867  
is0889: clau|AEV95509.1 ldam|AH70\_06395 pent|YP\_804868  
is0890: clau|AEV95510.1 ldam|AH70\_06390 pent|YP\_804869  
is0891: clau|AEV95511.1 ldam|AH70\_06385 pent|YP\_804870  
is0892: clau|AEV95512.1 ldam|AH70\_06380 pent|YP\_804871  
is0893: clau|AEV95513.1 ldam|AH70\_06375 pent|YP\_804872  
is0894: clau|AEV95514.1 ldam|AH70\_06370 pent|YP\_804873  
is0895: clau|AEV95515.1 ldam|AH70\_06365 pent|YP\_804874

is0896: clau|AEV95516.1 ldam|AH70\_06355 pent|YP\_804876  
is0897: clau|AEV95517.1 ldam|AH70\_06350 pent|YP\_804877  
is0898: clau|AEV95518.1 ldam|AH70\_06345 pent|YP\_804878  
is0899: clau|AEV95519.1 ldam|AH70\_06340 pent|YP\_804879  
is0900: clau|AEV95520.1 ldam|AH70\_06335 pent|YP\_804880  
is0901: clau|AEV95521.1 ldam|AH70\_06330 pent|YP\_804881  
is0902: clau|AEV95522.1 ldam|AH70\_06325 pent|YP\_804882  
is0903: clau|AEV95523.1 ldam|AH70\_06320 pent|YP\_804883  
is0904: clau|AEV95524.1 ldam|AH70\_06315 pent|YP\_804884  
is0905: clau|AEV95526.1 ldam|AH70\_06310 pent|YP\_804886  
is0906: clau|AEV95527.1 ldam|AH70\_06305 pent|YP\_804887  
is0907: clau|AEV95528.1 ldam|AH70\_06300 pent|YP\_804888  
is0908: clau|AEV95529.1 ldam|AH70\_06295 pent|YP\_804889  
is0909: clau|AEV95530.1 ldam|AH70\_06290 pent|YP\_804890  
is0910: clau|AEV95531.1 ldam|AH70\_06285 pent|YP\_804891  
is0911: clau|AEV95532.1 ldam|AH70\_06280 pent|YP\_804892  
is0912: clau|AEV95533.1 ldam|AH70\_06275 pent|YP\_804893  
is0913: clau|AEV95534.1 ldam|AH70\_06270 pent|YP\_804894  
is0914: clau|AEV95535.1 ldam|AH70\_06265 pent|YP\_804895  
is0915: clau|AEV95536.1 ldam|AH70\_06260 pent|YP\_804896  
is0916: clau|AEV95537.1 ldam|AH70\_06255 pent|YP\_804897  
is0917: clau|AEV95538.1 ldam|AH70\_06250 pent|YP\_804898  
is0918: clau|AEV95539.1 ldam|AH70\_06245 pent|YP\_804899  
is0919: clau|AEV95541.1 ldam|AH70\_06235 pent|YP\_804900  
is0920: clau|AEV95542.1 ldam|AH70\_06230 pent|YP\_804901  
is0921: clau|AEV95543.1 ldam|AH70\_06225 pent|YP\_804902  
is0922: clau|AEV95545.1 ldam|AH70\_06215 pent|YP\_804904  
is0923: clau|AEV95546.1 ldam|AH70\_06210 pent|YP\_804905  
is0924: clau|AEV95548.1 ldam|AH70\_06200 pent|YP\_804907  
is0925: clau|AEV95549.1 ldam|AH70\_06150 pent|YP\_804909  
is0926: clau|AEV95550.1 ldam|AH70\_06145 pent|YP\_804910  
is0927: clau|AEV95551.1 ldam|AH70\_06140 pent|YP\_804911  
is0928: clau|AEV95554.1 ldam|AH70\_06120 pent|YP\_804914  
is0929: clau|AEV95555.1 ldam|AH70\_06110 pent|YP\_804916  
is0930: clau|AEV95556.1 ldam|AH70\_06105 pent|YP\_804917  
is0931: clau|AEV95557.1 ldam|AH70\_06100 pent|YP\_804918  
is0932: clau|AEV95560.1 ldam|AH70\_06070 pent|YP\_804919  
is0933: clau|AEV95561.1 ldam|AH70\_06065 pent|YP\_804920  
is0934: clau|AEV95562.1 ldam|AH70\_06060 pent|YP\_804921  
is0935: clau|AEV95563.1 ldam|AH70\_06050 pent|YP\_804922  
is0936: clau|AEV95565.1 ldam|AH70\_06035 pent|YP\_804923  
is0937: clau|AEV95568.1 ldam|AH70\_06000 pent|YP\_804902  
is0938: clau|AEV95569.1 ldam|AH70\_09355 pent|YP\_804310  
is0939: clau|AEV95574.1 ldam|AH70\_02045 pent|YP\_804928  
is0940: clau|AEV95575.1 ldam|AH70\_04385 pent|YP\_804087  
is0941: clau|AEV95581.1 ldam|AH70\_00850 pent|YP\_803950  
is0942: clau|AEV95583.1 ldam|AH70\_00845 pent|YP\_803945  
is0943: clau|AEV95587.1 ldam|AH70\_01530 pent|YP\_803941  
is0944: clau|AEV95590.1 ldam|AH70\_07700 pent|YP\_805086  
is0945: clau|AEV95633.1 pent|YP\_805168 ldam|AH70\_01090  
is0946: clau|AEV95649.1 ldam|AH70\_03745 pent|YP\_803938  
is0947: clau|AEV95650.1 ldam|AH70\_03740 pent|YP\_803937  
is0948: clau|AEV95651.1 ldam|AH70\_03735 pent|YP\_803936  
is0949: clau|AEV95652.1 ldam|AH70\_03730 pent|YP\_803935  
is0950: clau|AEV95654.1 ldam|AH70\_03710 pent|YP\_803933  
is0951: clau|AEV95659.1 ldam|AH70\_01520 pent|YP\_803613  
is0952: clau|AEV95660.1 ldam|AH70\_04465 pent|YP\_803699  
is0953: clau|AEV95661.1 ldam|AH70\_01205 pent|YP\_805238  
is0954: clau|AEV95662.1 ldam|AH70\_01200 pent|YP\_805239  
is0955: clau|AEV95663.1 ldam|AH70\_01195 pent|YP\_805240  
is0956: clau|AEV95670.1 ldam|AH70\_07035 pent|YP\_803930  
is0957: clau|AEV95671.1 ldam|AH70\_07030 pent|YP\_803929  
is0958: clau|AEV95672.1 ldam|AH70\_07025 pent|YP\_803928  
is0959: clau|AEV95673.1 ldam|AH70\_07020 pent|YP\_803927  
is0960: clau|AEV95674.1 ldam|AH70\_07015 pent|YP\_803926  
is0961: clau|AEV95676.1 ldam|AH70\_07005 pent|YP\_803924  
is0962: clau|AEV95677.1 ldam|AH70\_07000 pent|YP\_803923  
is0963: clau|AEV95678.1 ldam|AH70\_06995 pent|YP\_803922  
is0964: clau|AEV95680.1 ldam|AH70\_02385 pent|YP\_803920  
is0965: clau|AEV95681.1 ldam|AH70\_02380 pent|YP\_803918  
is0966: clau|AEV95682.1 ldam|AH70\_02375 pent|YP\_805099  
is0967: clau|AEV95683.1 ldam|AH70\_02740 pent|YP\_803917  
is0968: clau|AEV95684.1 ldam|AH70\_01960 pent|YP\_803916  
is0969: clau|AEV95686.1 ldam|AH70\_02735 pent|YP\_803915  
is0970: clau|AEV95687.1 ldam|AH70\_02730 pent|YP\_803914  
is0971: clau|AEV95691.1 ldam|AH70\_00625 pent|YP\_803913  
is0972: clau|AEV95692.1 ldam|AH70\_00630 pent|YP\_803912  
is0973: clau|AEV95693.1 ldam|AH70\_00635 pent|YP\_803911  
is0974: clau|AEV95694.1 ldam|AH70\_00640 pent|YP\_803910  
is0975: clau|AEV95695.1 ldam|AH70\_07065 pent|YP\_803909  
is0976: clau|AEV95696.1 ldam|AH70\_00655 pent|YP\_803908  
is0977: clau|AEV95698.1 ldam|AH70\_00665 pent|YP\_803906  
is0978: clau|AEV95700.1 ldam|AH70\_00675 pent|YP\_803904

is0979: clau|AEV95701.1 ldam|AH70\_00690 pent|YP\_803903  
is0980: clau|AEV95702.1 ldam|AH70\_00695 pent|YP\_803902  
is0981: clau|AEV95704.1 ldam|AH70\_00700 pent|YP\_803901  
is0982: clau|AEV95705.1 ldam|AH70\_00705 pent|YP\_803900  
is0983: clau|AEV95706.1 ldam|AH70\_00710 pent|YP\_803899  
is0984: clau|AEV95709.1 ldam|AH70\_00725 pent|YP\_803896  
is0985: clau|AEV95710.1 ldam|AH70\_00730 pent|YP\_803895  
is0986: clau|AEV95711.1 ldam|AH70\_04165 pent|YP\_803756  
is0987: clau|AEV95712.1 ldam|AH70\_00740 pent|YP\_803893  
is0988: clau|AEV95716.1 ldam|AH70\_01355 pent|YP\_803891  
is0989: clau|AEV95717.1 ldam|AH70\_01360 pent|YP\_803890  
is0990: clau|AEV95718.1 ldam|AH70\_01875 pent|YP\_803889  
is0991: clau|AEV95720.1 ldam|AH70\_01895 pent|YP\_803888  
is0992: clau|AEV95721.1 ldam|AH70\_01900 pent|YP\_803887  
is0993: clau|AEV95723.1 ldam|AH70\_10580 pent|YP\_803766  
is0994: clau|AEV95724.1 ldam|AH70\_07785 pent|YP\_803764  
is0995: clau|AEV95725.1 ldam|AH70\_03330 pent|YP\_804170  
is0996: clau|AEV95726.1 ldam|AH70\_10090 pent|YP\_805079  
is0997: clau|AEV95728.1 ldam|AH70\_10060 pent|YP\_805081  
is0998: clau|AEV95730.1 ldam|AH70\_05980 pent|YP\_803754  
is0999: clau|AEV95731.1 ldam|AH70\_04250 pent|YP\_805084  
is1000: clau|AEV95732.1 ldam|AH70\_07695 pent|YP\_805085  
is1001: clau|AEV95734.1 ldam|AH70\_09010 pent|YP\_805096  
is1002: clau|AEV95735.1 ldam|AH70\_08095 pent|YP\_803698  
is1003: clau|AEV95736.1 ldam|AH70\_01675 pent|YP\_803585  
is1004: clau|AEV95739.1 ldam|AH70\_05355 pent|YP\_803688  
is1005: clau|AEV95742.1 ldam|AH70\_05460 pent|YP\_804049  
is1006: clau|AEV95750.1 ldam|AH70\_04350 pent|YP\_805101  
is1007: clau|AEV95752.1 ldam|AH70\_05555 pent|YP\_804941  
is1008: clau|AEV95759.1 ldam|AH70\_08245 pent|YP\_803574  
is1009: clau|AEV95761.1 ldam|AH70\_08235 pent|YP\_803576  
is1010: clau|AEV95766.1 ldam|AH70\_09750 pent|YP\_805139  
is1011: clau|AEV95767.1 ldam|AH70\_00540 pent|YP\_805128  
is1012: clau|AEV95770.1 ldam|AH70\_04650 pent|YP\_804839  
is1013: clau|AEV95778.1 ldam|AH70\_00020 pent|YP\_804089  
is1014: clau|AEV95785.1 ldam|AH70\_08230 pent|YP\_803663  
is1015: clau|AEV95786.1 ldam|AH70\_06555 pent|YP\_803696  
is1016: clau|AEV95788.1 ldam|AH70\_07825 pent|YP\_804405  
is1017: clau|AEV95792.1 ldam|AH70\_05265 pent|YP\_805131  
is1018: clau|AEV95804.1 ldam|AH70\_06750 pent|YP\_803761  
is1019: clau|AEV95808.1 ldam|AH70\_05705 pent|YP\_804453  
is1020: clau|AEV95820.1 ldam|AH70\_01555 pent|YP\_803657  
is1021: clau|AEV95826.1 ldam|AH70\_03340 pent|YP\_804093  
is1022: clau|AEV95829.1 ldam|AH70\_06540 pent|YP\_805145  
is1023: clau|AEV95830.1 ldam|AH70\_06535 pent|YP\_805146  
is1024: clau|AEV95833.1 pent|YP\_803946 ldam|AH70\_05450  
is1025: clau|AEV95846.1 ldam|AH70\_04930 pent|YP\_805142  
is1026: clau|AEV95853.1 ldam|AH70\_04150 pent|YP\_805173  
is1027: clau|AEV95854.1 ldam|AH70\_04145 pent|YP\_805174  
is1028: clau|AEV95861.1 ldam|AH70\_04120 pent|YP\_805176  
is1029: clau|AEV95862.1 ldam|AH70\_04115 pent|YP\_805177  
is1030: clau|AEV95863.1 ldam|AH70\_04105 pent|YP\_805178  
is1031: clau|AEV95864.1 ldam|AH70\_04100 pent|YP\_805179  
is1032: clau|AEV95865.1 ldam|AH70\_04095 pent|YP\_805180  
is1033: clau|AEV95866.1 ldam|AH70\_10780 pent|YP\_805181  
is1034: clau|AEV95867.1 ldam|AH70\_04080 pent|YP\_805182  
is1035: clau|AEV95869.1 ldam|AH70\_04075 pent|YP\_805183  
is1036: clau|AEV95870.1 ldam|AH70\_04070 pent|YP\_805184  
is1037: clau|AEV95872.1 ldam|AH70\_06800 pent|YP\_805072  
is1038: clau|AEV95873.1 ldam|AH70\_04065 pent|YP\_805185  
is1039: clau|AEV95874.1 ldam|AH70\_04060 pent|YP\_805186  
is1040: clau|AEV95875.1 ldam|AH70\_04055 pent|YP\_805187  
is1041: clau|AEV95881.1 ldam|AH70\_03750 pent|YP\_803744  
is1042: clau|AEV95883.1 ldam|AH70\_04050 pent|YP\_804539  
is1043: clau|AEV95885.1 ldam|AH70\_04040 pent|YP\_804541  
is1044: clau|AEV95890.1 ldam|AH70\_04690 pent|YP\_805267  
is1045: clau|AEV95891.1 ldam|AH70\_04685 pent|YP\_805268  
is1046: clau|AEV95897.1 ldam|AH70\_08450 pent|YP\_805189  
is1047: clau|AEV95898.1 ldam|AH70\_08420 pent|YP\_805190  
is1048: clau|AEV95899.1 ldam|AH70\_08415 pent|YP\_805191  
is1049: clau|AEV95900.1 ldam|AH70\_08345 pent|YP\_805193  
is1050: clau|AEV95901.1 ldam|AH70\_08340 pent|YP\_805194  
is1051: clau|AEV95902.1 ldam|AH70\_08325 pent|YP\_805195  
is1052: clau|AEV95912.1 ldam|AH70\_10220 pent|YP\_804464  
is1053: clau|AEV95929.1 ldam|AH70\_01155 pent|YP\_803775  
is1054: clau|AEV95930.1 ldam|AH70\_01160 pent|YP\_803774  
is1055: clau|AEV95931.1 ldam|AH70\_01150 pent|YP\_803773  
is1056: clau|AEV95935.1 ldam|AH70\_05425 pent|YP\_805201  
is1057: clau|AEV95936.1 ldam|AH70\_01965 pent|YP\_805202  
is1058: clau|AEV95937.1 ldam|AH70\_04790 pent|YP\_805204  
is1059: clau|AEV95944.1 ldam|AH70\_06745 pent|YP\_805206  
is1060: clau|AEV95945.1 ldam|AH70\_06740 pent|YP\_805207  
is1061: clau|AEV95961.1 ldam|AH70\_05320 pent|YP\_803772

is1062: clau|AEV95973.1 ldam|AH70\_09325 pent|YP\_805111  
is1063: clau|AEV95974.1 ldam|AH70\_05340 pent|YP\_803645  
is1064: clau|AEV95975.1 ldam|AH70\_05335 pent|YP\_803644  
is1065: clau|AEV95976.1 ldam|AH70\_05445 pent|YP\_805112  
is1066: clau|AEV95982.1 ldam|AH70\_00190 pent|YP\_805115  
is1067: clau|AEV95983.1 ldam|AH70\_00185 pent|YP\_805116  
is1068: clau|AEV95984.1 ldam|AH70\_00180 pent|YP\_805117  
is1069: clau|AEV95985.1 ldam|AH70\_00175 pent|YP\_805118  
is1070: clau|AEV95989.1 ldam|AH70\_05465 pent|YP\_805119  
is1071: clau|AEV95999.1 ldam|AH70\_05405 pent|YP\_805120  
is1072: clau|AEV96004.1 ldam|AH70\_09810 pent|YP\_805209  
is1073: clau|AEV96010.1 ldam|AH70\_02475 pent|YP\_805213  
is1074: clau|AEV96011.1 ldam|AH70\_02470 pent|YP\_805214  
is1075: clau|AEV96012.1 ldam|AH70\_05975 pent|YP\_804074  
is1076: clau|AEV96014.1 ldam|AH70\_04495 pent|YP\_803659  
is1077: clau|AEV96021.1 ldam|AH70\_01055 pent|YP\_805243  
is1078: clau|AEV96033.1 ldam|AH70\_05330 pent|YP\_805102  
is1079: clau|AEV96034.1 ldam|AH70\_05325 pent|YP\_805103  
is1080: clau|AEV96037.1 ldam|AH70\_11350 pent|YP\_805292  
is1081: clau|AEV96038.1 ldam|AH70\_11345 pent|YP\_805255  
is1082: clau|AEV96039.1 ldam|AH70\_11340 pent|YP\_805256  
is1083: clau|AEV96040.1 ldam|AH70\_11335 pent|YP\_805257  
is1084: clau|AEV96041.1 ldam|AH70\_11330 pent|YP\_805258  
is1085: clau|AEV96042.1 ldam|AH70\_11325 pent|YP\_805259  
is1086: clau|AEV96043.1 ldam|AH70\_11320 pent|YP\_805260  
is1087: clau|AEV96044.1 ldam|AH70\_01025 pent|YP\_803955  
is1088: clau|AEV96054.1 ldam|AH70\_04400 pent|YP\_803595  
is1089: clau|AEV96055.1 ldam|AH70\_04395 pent|YP\_803594  
is1090: clau|AEV96057.1 ldam|AH70\_06700 pent|YP\_803593  
is1091: clau|AEV96059.1 ldam|AH70\_04380 pent|YP\_803591  
is1092: clau|AEV96060.1 ldam|AH70\_04375 pent|YP\_805293  
is1093: clau|AEV96061.1 ldam|AH70\_04370 pent|YP\_805294  
is1094: clau|AEV96062.1 ldam|AH70\_04365 pent|YP\_805295  
is1095: clau|AEV96064.1 ldam|AH70\_04325 pent|YP\_805297  
is1096: clau|AEV96065.1 ldam|AH70\_04320 pent|YP\_805298  
is1097: clau|AEV96067.1 ldam|AH70\_04315 pent|YP\_805299  
is1098: clau|AEV96068.1 ldam|AH70\_04310 pent|YP\_805300  
is1099: clau|AEV96070.1 ldam|AH70\_04295 pent|YP\_805301  
is1100: clau|AEV96072.1 ldam|AH70\_04280 pent|YP\_805302  
is1101: clau|AEV96075.1 ldam|AH70\_04230 pent|YP\_805304  
is1102: clau|AEV96076.1 ldam|AH70\_04225 pent|YP\_805305  
is1103: clau|AEV96077.1 ldam|AH70\_04215 pent|YP\_805306  
is1104: clau|AEV96078.1 ldam|AH70\_04210 pent|YP\_805307  
is1105: clau|AEV96085.1 ldam|AH70\_00375 pent|YP\_804862  
is1106: clau|AEV96176.1 ldam|AH70\_10915 pent|YP\_804516  
is1107: clau|AEV96200.1 ldam|AH70\_11220 pent|YP\_804725  
is1108: clau|AEV96201.1 ldam|AH70\_08320 pent|YP\_804724  
is1109: clau|AEV94411.1 clau|AEV95841.1  
is1110: clau|AEV94501.1 clau|AEV96001.1  
is1111: clau|AEV94572.1 clau|AEV94573.1  
is1112: clau|AEV94583.1 clau|AEV94584.1  
is1113: clau|AEV94635.1 clau|AEV94902.1  
is1114: clau|AEV94702.1 clau|AEV94703.1  
is1115: clau|AEV94722.1 clau|AEV94767.1  
is1116: clau|AEV94754.1 clau|AEV95595.1  
is1117: clau|AEV94788.1 clau|AEV95837.1  
is1118: clau|AEV94789.1 clau|AEV95836.1  
is1119: clau|AEV94794.1 clau|AFD55029.1  
is1120: clau|AEV94838.1 clau|AEV95209.1  
is1121: clau|AEV95140.1 clau|AEV95152.1  
is1122: clau|AEV95144.1 clau|AEV95148.1  
is1123: clau|AEV95570.1 clau|AEV96181.1  
is1124: clau|AEV95598.1 clau|AEV95630.1  
is1125: clau|AEV95613.1 clau|AEV96168.1  
is1126: clau|AEV95614.1 clau|AEV95616.1  
is1127: clau|AEV95703.1 clau|AEV95729.1  
is1128: clau|AEV95755.1 clau|AEV95942.1  
is1129: clau|AEV96084.1 clau|AEV96094.1  
is1130: clau|AEV96103.1 clau|AFD55018.1  
is1131: clau|AEV96106.1 clau|AEV96161.1  
is1132: clau|AEV96107.1 clau|AEV96160.1  
is1133: clau|AEV96116.1 clau|AFD55021.1  
is1134: clau|AEV96151.1 clau|AEV96166.1  
is1135: ldam|AH70\_00005 ldam|AH70\_01020  
is1136: ldam|AH70\_00875 ldam|AH70\_05645  
is1137: ldam|AH70\_00895 ldam|AH70\_00905  
is1138: ldam|AH70\_01050 ldam|AH70\_03880  
is1139: ldam|AH70\_01105 ldam|AH70\_06805  
is1140: ldam|AH70\_01310 ldam|AH70\_06595  
is1141: ldam|AH70\_01445 ldam|AH70\_01450  
is1142: ldam|AH70\_02065 ldam|AH70\_10305  
is1143: ldam|AH70\_02085 ldam|AH70\_03670  
is1144: ldam|AH70\_02325 ldam|AH70\_05255

is1145: ldam|AH70\_02865 ldam|AH70\_08110  
is1146: ldam|AH70\_02870 ldam|AH70\_07835  
is1147: ldam|AH70\_03800 ldam|AH70\_03810  
is1148: ldam|AH70\_04305 ldam|AH70\_06840  
is1149: ldam|AH70\_04410 ldam|AH70\_04420  
is1150: ldam|AH70\_04505 ldam|AH70\_04510  
is1151: ldam|AH70\_05740 ldam|AH70\_07090  
is1152: ldam|AH70\_08385 ldam|AH70\_08390  
is1153: ldam|AH70\_08460 ldam|AH70\_08820  
is1154: pent|YP\_803667 pent|YP\_804546  
is1155: pent|YP\_803705 pent|YP\_803731  
is1156: pent|YP\_803726 pent|YP\_803742  
is1157: pent|YP\_803727 pent|YP\_803743  
is1158: pent|YP\_804046 pent|YP\_803746  
is1159: pent|YP\_804060 pent|YP\_805233  
is1160: pent|YP\_804231 pent|YP\_804969  
is1161: pent|YP\_804250 pent|YP\_804533  
is1162: pent|YP\_804260 pent|YP\_804520  
is1163: pent|YP\_804261 pent|YP\_804519  
is1164: pent|YP\_804266 pent|YP\_804513  
is1165: pent|YP\_804268 pent|YP\_804512  
is1166: pent|YP\_804269 pent|YP\_804511  
is1167: pent|YP\_804270 pent|YP\_804508  
is1168: pent|YP\_804408 pent|YP\_804528  
is1169: pent|YP\_804417 pent|YP\_805211  
is1170: pent|YP\_804828 pent|YP\_804829  
is1171: pent|YP\_805090 pent|YP\_805091  
is1172: pent|YP\_804255 pent|YP\_804527  
is1173: pent|YP\_804257 pent|YP\_804525  
is1174: pent|YP\_804258 pent|YP\_804522  
is1175: pent|YP\_804265 pent|YP\_804514  
is1176: pent|YP\_804406 pent|YP\_805251  
is1177: pent|YP\_804858 pent|YP\_805165  
is1178: ldam|AH70\_01275 ldam|AH70\_02150  
is1179: clau|AEV95619.1 clau|AEV95623.1  
is1180: clau|AEV96171.1 clau|AEV96172.1  
is1181: clau|AEV94339.1 pent|YP\_805308  
is1182: clau|AEV94353.1 ldam|AH70\_10695  
is1183: clau|AEV94360.1 ldam|AH70\_00285  
is1184: clau|AEV94376.1 pent|YP\_805217  
is1185: clau|AEV94388.1 pent|YP\_803777  
is1186: clau|AEV94396.1 ldam|AH70\_09655  
is1187: clau|AEV94404.1 pent|YP\_803785  
is1188: clau|AEV94405.1 pent|YP\_803786  
is1189: clau|AEV94417.1 pent|YP\_803803  
is1190: clau|AEV94422.1 pent|YP\_803808  
is1191: clau|AEV94431.1 ldam|AH70\_00880  
is1192: clau|AEV94432.1 pent|YP\_804355  
is1193: clau|AEV94433.1 pent|YP\_804356  
is1194: clau|AEV94434.1 pent|YP\_804357  
is1195: clau|AEV94436.1 pent|YP\_804359  
is1196: clau|AEV94437.1 pent|YP\_804360  
is1197: clau|AEV94439.1 pent|YP\_804362  
is1198: clau|AEV94443.1 pent|YP\_804366  
is1199: clau|AEV94445.1 pent|YP\_805100  
is1200: clau|AEV94447.1 pent|YP\_803608  
is1201: clau|AEV94449.1 ldam|AH70\_09685  
is1202: clau|AEV94456.1 pent|YP\_803859  
is1203: clau|AEV94457.1 pent|YP\_803860  
is1204: clau|AEV94458.1 pent|YP\_803861  
is1205: clau|AEV94459.1 pent|YP\_803862  
is1206: clau|AEV94460.1 pent|YP\_803857  
is1207: clau|AEV94461.1 pent|YP\_803669  
is1208: clau|AEV94468.1 ldam|AH70\_08440  
is1209: clau|AEV94479.1 pent|YP\_803675  
is1210: clau|AEV94480.1 pent|YP\_803676  
is1211: clau|AEV94481.1 pent|YP\_803677  
is1212: clau|AEV94490.1 ldam|AH70\_02660  
is1213: clau|AEV94491.1 ldam|AH70\_02665  
is1214: pent|YP\_804076 ldam|AH70\_01000  
is1215: clau|AEV94492.1 ldam|AH70\_10140  
is1216: clau|AEV94496.1 ldam|AH70\_10500  
is1217: clau|AEV94500.1 pent|YP\_804161  
is1218: clau|AEV94502.1 pent|YP\_805241  
is1219: clau|AEV94504.1 pent|YP\_805149  
is1220: clau|AEV94507.1 pent|YP\_803678  
is1221: clau|AEV94510.1 pent|YP\_804124  
is1222: clau|AEV94521.1 ldam|AH70\_01100  
is1223: clau|AEV94530.1 pent|YP\_805042  
is1224: clau|AEV94533.1 pent|YP\_803640  
is1225: clau|AEV94535.1 ldam|AH70\_05285  
is1226: clau|AEV94567.1 ldam|AH70\_07365  
is1227: clau|AEV94581.1 ldam|AH70\_03460

is1228: clau|AEV94587.1 ldam|AH70\_02845  
is1229: clau|AEV94601.1 pent|YP\_805002  
is1230: clau|AEV94610.1 pent|YP\_803583  
is1231: clau|AEV94611.1 pent|YP\_803584  
is1232: clau|AEV94612.1 ldam|AH70\_02620  
is1233: clau|AEV94615.1 pent|YP\_803581  
is1234: clau|AEV94616.1 ldam|AH70\_06985  
is1235: clau|AEV94619.1 ldam|AH70\_04245  
is1236: clau|AEV94624.1 ldam|AH70\_05535  
is1237: clau|AEV94633.1 ldam|AH70\_05345  
is1238: clau|AEV94644.1 ldam|AH70\_04020  
is1239: clau|AEV94645.1 ldam|AH70\_04025  
is1240: clau|AEV94646.1 pent|YP\_804982  
is1241: clau|AEV94650.1 pent|YP\_804985  
is1242: clau|AEV94658.1 ldam|AH70\_06565  
is1243: clau|AEV94661.1 ldam|AH70\_10165  
is1244: clau|AEV94672.1 ldam|AH70\_04910  
is1245: clau|AEV94687.1 pent|YP\_804162  
is1246: clau|AEV94689.1 pent|YP\_803877  
is1247: clau|AEV94692.1 ldam|AH70\_09305  
is1248: clau|AEV94694.1 ldam|AH70\_01985  
is1249: clau|AEV94699.1 ldam|AH70\_01975  
is1250: clau|AEV94705.1 ldam|AH70\_01700  
is1251: clau|AEV94706.1 pent|YP\_804566  
is1252: clau|AEV94707.1 pent|YP\_804565  
is1253: clau|AEV94713.1 pent|YP\_803771  
is1254: clau|AEV94716.1 ldam|AH70\_07550  
is1255: clau|AEV94724.1 ldam|AH70\_08270  
is1256: clau|AEV94725.1 ldam|AH70\_00330  
is1257: clau|AEV94726.1 ldam|AH70\_00325  
is1258: clau|AEV94727.1 ldam|AH70\_00320  
is1259: clau|AEV94736.1 ldam|AH70\_09745  
is1260: clau|AEV94737.1 ldam|AH70\_09740  
is1261: clau|AEV94738.1 ldam|AH70\_09725  
is1262: clau|AEV94739.1 ldam|AH70\_09720  
is1263: clau|AEV94740.1 ldam|AH70\_09715  
is1264: clau|AEV94744.1 ldam|AH70\_07530  
is1265: clau|AEV94748.1 ldam|AH70\_07500  
is1266: clau|AEV94749.1 ldam|AH70\_07495  
is1267: clau|AEV94751.1 pent|YP\_803864  
is1268: clau|AEV94780.1 ldam|AH70\_07165  
is1269: clau|AEV94785.1 pent|YP\_803855  
is1270: clau|AEV94787.1 ldam|AH70\_10010  
is1271: clau|AEV94792.1 pent|YP\_804961  
is1272: pent|YP\_804057 ldam|AH70\_02855  
is1273: clau|AEV94796.1 pent|YP\_804959  
is1274: clau|AEV94802.1 pent|YP\_804953  
is1275: clau|AEV94856.1 pent|YP\_804841  
is1276: clau|AEV94857.1 pent|YP\_804842  
is1277: clau|AEV94858.1 pent|YP\_804843  
is1278: clau|AEV94859.1 pent|YP\_804844  
is1279: clau|AEV94879.1 pent|YP\_805253  
is1280: clau|AEV94880.1 pent|YP\_805252  
is1281: clau|AEV94881.1 ldam|AH70\_06855  
is1282: clau|AEV94883.1 pent|YP\_804072  
is1283: clau|AEV94884.1 pent|YP\_804071  
is1284: clau|AEV94886.1 pent|YP\_804070  
is1285: clau|AEV94887.1 ldam|AH70\_10270  
is1286: clau|AEV94888.1 ldam|AH70\_10275  
is1287: clau|AEV94893.1 pent|YP\_804505  
is1288: clau|AEV94900.1 ldam|AH70\_05815  
is1289: clau|AEV94904.1 ldam|AH70\_08355  
is1290: clau|AEV94905.1 ldam|AH70\_08360  
is1291: clau|AEV94906.1 pent|YP\_804837  
is1292: clau|AEV94909.1 pent|YP\_803815  
is1293: clau|AEV94914.1 pent|YP\_803616  
is1294: clau|AEV94915.1 pent|YP\_803617  
is1295: clau|AEV94916.1 pent|YP\_803618  
is1296: clau|AEV94917.1 pent|YP\_803619  
is1297: clau|AEV94918.1 pent|YP\_803620  
is1298: clau|AEV94919.1 pent|YP\_803621  
is1299: clau|AEV94931.1 pent|YP\_803598  
is1300: clau|AEV94932.1 pent|YP\_803599  
is1301: clau|AEV94935.1 pent|YP\_803762  
is1302: clau|AEV94936.1 pent|YP\_804823  
is1303: clau|AEV94962.1 pent|YP\_804797  
is1304: clau|AEV94968.1 ldam|AH70\_07360  
is1305: clau|AEV94973.1 pent|YP\_804787  
is1306: clau|AEV94994.1 pent|YP\_804766  
is1307: clau|AEV94998.1 pent|YP\_804762  
is1308: clau|AEV95007.1 pent|YP\_804753  
is1309: clau|AEV95023.1 pent|YP\_804737  
is1310: clau|AEV95028.1 pent|YP\_804732

is1311: clau|AEV95029.1 pent|YP\_804731  
is1312: clau|AEV95052.1 ldam|AH70\_02360  
is1313: clau|AEV95087.1 pent|YP\_804674  
is1314: clau|AEV95113.1 pent|YP\_804649  
is1315: clau|AEV95115.1 ldam|AH70\_00430  
is1316: clau|AEV95155.1 ldam|AH70\_09340  
is1317: clau|AEV95156.1 pent|YP\_804597  
is1318: clau|AEV95176.1 ldam|AH70\_05110  
is1319: clau|AEV95196.1 pent|YP\_804461  
is1320: clau|AEV95207.1 pent|YP\_804448  
is1321: clau|AEV95208.1 ldam|AH70\_01060  
is1322: clau|AEV95210.1 ldam|AH70\_05765  
is1323: clau|AEV95213.1 pent|YP\_804445  
is1324: clau|AEV95217.1 pent|YP\_804600  
is1325: clau|AEV95218.1 pent|YP\_804601  
is1326: clau|AEV95225.1 pent|YP\_804608  
is1327: clau|AEV95233.1 pent|YP\_804617  
is1328: clau|AEV95263.1 pent|YP\_804376  
is1329: clau|AEV95265.1 pent|YP\_804374  
is1330: clau|AEV95325.1 pent|YP\_804325  
is1331: clau|AEV95340.1 pent|YP\_803673  
is1332: clau|AEV95344.1 pent|YP\_804588  
is1333: clau|AEV95347.1 ldam|AH70\_05560  
is1334: clau|AEV95348.1 ldam|AH70\_05565  
is1335: clau|AEV95349.1 ldam|AH70\_05570  
is1336: clau|AEV95350.1 ldam|AH70\_05575  
is1337: clau|AEV95352.1 ldam|AH70\_05585  
is1338: clau|AEV95375.1 pent|YP\_804223  
is1339: clau|AEV95394.1 pent|YP\_804034  
is1340: clau|AEV95405.1 ldam|AH70\_00275  
is1341: clau|AEV95427.1 ldam|AH70\_08555  
is1342: clau|AEV95434.1 ldam|AH70\_01140  
is1343: clau|AEV95450.1 pent|YP\_804146  
is1344: clau|AEV95472.1 pent|YP\_804125  
is1345: clau|AEV95493.1 pent|YP\_804094  
is1346: clau|AEV95525.1 pent|YP\_804885  
is1347: clau|AEV95544.1 ldam|AH70\_06220  
is1348: clau|AEV95552.1 pent|YP\_804912  
is1349: clau|AEV95553.1 pent|YP\_804913  
is1350: clau|AEV95558.1 ldam|AH70\_06095  
is1351: clau|AEV95559.1 ldam|AH70\_06085  
is1352: clau|AEV95564.1 ldam|AH70\_06045  
is1353: clau|AEV95566.1 pent|YP\_804924  
is1354: clau|AEV95567.1 pent|YP\_804925  
is1355: clau|AEV95571.1 pent|YP\_804926  
is1356: clau|AEV95573.1 ldam|AH70\_10955  
is1357: clau|AEV95584.1 pent|YP\_803944  
is1358: clau|AEV95585.1 pent|YP\_803943  
is1359: clau|AEV95637.1 ldam|AH70\_04360  
is1360: clau|AEV95667.1 pent|YP\_805161  
is1361: clau|AEV95669.1 pent|YP\_805163  
is1362: ldam|AH70\_03515 pent|YP\_804053  
is1363: clau|AEV95679.1 pent|YP\_803921  
is1364: clau|AEV95685.1 ldam|AH70\_05385  
is1365: clau|AEV95689.1 ldam|AH70\_00560  
is1366: clau|AEV95690.1 ldam|AH70\_00565  
is1367: clau|AEV95697.1 pent|YP\_803907  
is1368: clau|AEV95707.1 pent|YP\_803898  
is1369: clau|AEV95713.1 pent|YP\_803570  
is1370: clau|AEV95714.1 pent|YP\_803572  
is1371: clau|AEV95715.1 pent|YP\_803892  
is1372: clau|AEV95719.1 ldam|AH70\_10195  
is1373: clau|AEV95722.1 ldam|AH70\_01905  
is1374: clau|AEV95727.1 pent|YP\_805080  
is1375: clau|AEV95737.1 ldam|AH70\_05755  
is1376: clau|AEV95741.1 ldam|AH70\_01350  
is1377: clau|AEV95743.1 pent|YP\_804031  
is1378: clau|AEV95744.1 ldam|AH70\_04340  
is1379: clau|AEV95745.1 ldam|AH70\_04335  
is1380: clau|AEV95746.1 ldam|AH70\_04330  
is1381: clau|AEV95748.1 ldam|AH70\_09805  
is1382: clau|AEV95749.1 ldam|AH70\_08785  
is1383: clau|AEV95751.1 ldam|AH70\_06550  
is1384: clau|AEV95753.1 ldam|AH70\_07815  
is1385: clau|AEV95756.1 ldam|AH70\_03090  
is1386: clau|AEV95764.1 pent|YP\_805107  
is1387: clau|AEV95768.1 pent|YP\_805129  
is1388: clau|AEV95769.1 ldam|AH70\_09695  
is1389: clau|AEV95773.1 pent|YP\_804091  
is1390: clau|AEV95777.1 pent|YP\_804090  
is1391: clau|AEV95779.1 pent|YP\_804088  
is1392: clau|AEV95781.1 pent|YP\_805248  
is1393: clau|AEV95789.1 pent|YP\_805134

is1394: clau|AEV95790.1 pent|YP\_805133  
is1395: clau|AEV95791.1 pent|YP\_805132  
is1396: clau|AEV95793.1 pent|YP\_805130  
is1397: clau|AEV95800.1 ldam|AH70\_01765  
is1398: clau|AEV95801.1 ldam|AH70\_01760  
is1399: clau|AEV95803.1 pent|YP\_803886  
is1400: clau|AEV95805.1 pent|YP\_803847  
is1401: clau|AEV95806.1 pent|YP\_803846  
is1402: clau|AEV95809.1 pent|YP\_804419  
is1403: clau|AEV95810.1 pent|YP\_804420  
is1404: clau|AEV95811.1 ldam|AH70\_05315  
is1405: clau|AEV95812.1 ldam|AH70\_10735  
is1406: clau|AEV95814.1 ldam|AH70\_10740  
is1407: clau|AEV95815.1 ldam|AH70\_10745  
is1408: clau|AEV95817.1 pent|YP\_805285  
is1409: clau|AEV95818.1 pent|YP\_805284  
is1410: clau|AEV95819.1 ldam|AH70\_01545  
is1411: clau|AEV95825.1 ldam|AH70\_02395  
is1412: clau|AEV95834.1 pent|YP\_804545  
is1413: clau|AEV95838.1 ldam|AH70\_08790  
is1414: clau|AEV95840.1 pent|YP\_805271  
is1415: clau|AEV95847.1 pent|YP\_805143  
is1416: clau|AEV95850.1 ldam|AH70\_02390  
is1417: clau|AEV95859.1 ldam|AH70\_01265  
is1418: clau|AEV95876.1 pent|YP\_803751  
is1419: clau|AEV95877.1 pent|YP\_803662  
is1420: clau|AEV95880.1 pent|YP\_804047  
is1421: clau|AEV95882.1 ldam|AH70\_05380  
is1422: clau|AEV95886.1 ldam|AH70\_00890  
is1423: clau|AEV95914.1 ldam|AH70\_07770  
is1424: clau|AEV95917.1 ldam|AH70\_07760  
is1425: clau|AEV95918.1 ldam|AH70\_07755  
is1426: clau|AEV95919.1 ldam|AH70\_07750  
is1427: clau|AEV95920.1 pent|YP\_803740  
is1428: clau|AEV95921.1 pent|YP\_803739  
is1429: clau|AEV95922.1 pent|YP\_803738  
is1430: clau|AEV95923.1 pent|YP\_803737  
is1431: clau|AEV95924.1 pent|YP\_803736  
is1432: clau|AEV95925.1 pent|YP\_803735  
is1433: clau|AEV95926.1 ldam|AH70\_08290  
is1434: clau|AEV95927.1 ldam|AH70\_08285  
is1435: clau|AEV95928.1 ldam|AH70\_08280  
is1436: clau|AEV95933.1 pent|YP\_805200  
is1437: clau|AEV95940.1 ldam|AH70\_01315  
is1438: clau|AEV95943.1 pent|YP\_805205  
is1439: clau|AEV95948.1 pent|YP\_805088  
is1440: clau|AEV95949.1 pent|YP\_805123  
is1441: clau|AEV95950.1 pent|YP\_805124  
is1442: clau|AEV95951.1 pent|YP\_805125  
is1443: clau|AEV95952.1 ldam|AH70\_00950  
is1444: clau|AEV95956.1 ldam|AH70\_06585  
is1445: clau|AEV95957.1 pent|YP\_803597  
is1446: clau|AEV95960.1 ldam|AH70\_03085  
is1447: clau|AEV95977.1 ldam|AH70\_07070  
is1448: clau|AEV95978.1 pent|YP\_805113  
is1449: ldam|AH70\_08770 pent|YP\_805274  
is1450: clau|AEV95991.1 pent|YP\_805152  
is1451: clau|AEV95992.1 pent|YP\_805153  
is1452: clau|AEV95993.1 pent|YP\_805154  
is1453: clau|AEV95994.1 pent|YP\_805155  
is1454: clau|AEV95995.1 pent|YP\_805156  
is1455: clau|AEV95996.1 pent|YP\_805157  
is1456: clau|AEV95997.1 pent|YP\_805158  
is1457: clau|AEV95998.1 pent|YP\_805159  
is1458: clau|AEV96003.1 ldam|AH70\_05750  
is1459: clau|AEV96008.1 pent|YP\_803952  
is1460: clau|AEV96009.1 ldam|AH70\_04435  
is1461: clau|AEV96013.1 ldam|AH70\_05770  
is1462: clau|AEV96017.1 ldam|AH70\_04475  
is1463: clau|AEV96019.1 pent|YP\_803697  
is1464: clau|AEV96023.1 pent|YP\_805246  
is1465: clau|AEV96027.1 ldam|AH70\_01670  
is1466: clau|AEV96029.1 pent|YP\_805303  
is1467: clau|AEV96036.1 pent|YP\_804064  
is1468: clau|AEV96053.1 ldam|AH70\_05375  
is1469: clau|AEV96071.1 ldam|AH70\_04285  
is1470: clau|AEV96089.1 ldam|AH70\_02035  
is1471: clau|AEV96097.1 ldam|AH70\_06845  
is1472: clau|AEV96100.1 ldam|AH70\_03630  
is1473: clau|AEV96101.1 ldam|AH70\_03625  
is1474: clau|AEV96104.1 ldam|AH70\_11075  
is1475: clau|AEV96108.1 ldam|AH70\_01865  
is1476: clau|AEV96111.1 ldam|AH70\_10710

is1477: clau|AEV96120.1 ldam|AH70\_00010  
is1478: clau|AEV96141.1 ldam|AH70\_11260  
is1479: clau|AEV96150.1 ldam|AH70\_03070  
is1480: clau|AEV96152.1 ldam|AH70\_00585  
is1481: clau|AEV96153.1 ldam|AH70\_00580  
is1482: clau|AEV96177.1 ldam|AH70\_10920  
is1483: clau|AEV96180.1 ldam|AH70\_03150  
is1484: clau|AEV96187.1 ldam|AH70\_10310  
is1485: clau|AEV96191.1 ldam|AH70\_09615  
is1486: clau|AEV96193.1 ldam|AH70\_02080  
is1487: clau|AEV96195.1 ldam|AH70\_02070  
is1488: clau|AEV96196.1 ldam|AH70\_11205  
is1489: clau|AEV96197.1 ldam|AH70\_11210  
is1490: clau|AEV96202.1 ldam|AH70\_03335  
is1491: clau|AEV96204.1 ldam|AH70\_11240  
is1492: clau|AEV96205.1 ldam|AH70\_11245  
is1493: ldam|AH70\_00130 pent|YP\_803672  
is1494: ldam|AH70\_00290 pent|YP\_803610  
is1495: ldam|AH70\_00295 pent|YP\_804808  
is1496: ldam|AH70\_00300 pent|YP\_804807  
is1497: ldam|AH70\_00315 pent|YP\_803668  
is1498: ldam|AH70\_00350 pent|YP\_803825  
is1499: ldam|AH70\_00380 pent|YP\_803701  
is1500: ldam|AH70\_00385 pent|YP\_803700  
is1501: ldam|AH70\_00550 pent|YP\_805127  
is1502: ldam|AH70\_00595 pent|YP\_805279  
is1503: ldam|AH70\_00685 pent|YP\_804554  
is1504: ldam|AH70\_00920 pent|YP\_805242  
is1505: ldam|AH70\_00960 pent|YP\_803614  
is1506: ldam|AH70\_00965 pent|YP\_804368  
is1507: ldam|AH70\_00975 pent|YP\_805231  
is1508: ldam|AH70\_00990 pent|YP\_803650  
is1509: ldam|AH70\_01340 pent|YP\_803631  
is1510: ldam|AH70\_01390 pent|YP\_804099  
is1511: ldam|AH70\_01425 pent|YP\_803723  
is1512: ldam|AH70\_01430 pent|YP\_803722  
is1513: ldam|AH70\_01435 pent|YP\_803721  
is1514: ldam|AH70\_01440 pent|YP\_803720  
is1515: ldam|AH70\_01500 pent|YP\_803629  
is1516: ldam|AH70\_01565 pent|YP\_804056  
is1517: ldam|AH70\_01715 pent|YP\_803586  
is1518: ldam|AH70\_01725 pent|YP\_804943  
is1519: ldam|AH70\_01930 pent|YP\_805106  
is1520: ldam|AH70\_02405 pent|YP\_803947  
is1521: ldam|AH70\_02410 pent|YP\_803949  
is1522: ldam|AH70\_02640 pent|YP\_805003  
is1523: ldam|AH70\_02680 pent|YP\_803580  
is1524: ldam|AH70\_02760 pent|YP\_804609  
is1525: ldam|AH70\_02815 pent|YP\_804618  
is1526: ldam|AH70\_02875 pent|YP\_804467  
is1527: ldam|AH70\_03220 pent|YP\_804827  
is1528: ldam|AH70\_03510 pent|YP\_804052  
is1529: ldam|AH70\_03780 pent|YP\_805219  
is1530: ldam|AH70\_04290 pent|YP\_803866  
is1531: ldam|AH70\_04455 pent|YP\_805083  
is1532: ldam|AH70\_05295 pent|YP\_803611  
is1533: ldam|AH70\_05455 pent|YP\_803948  
is1534: ldam|AH70\_05495 pent|YP\_804478  
is1535: ldam|AH70\_05500 pent|YP\_804477  
is1536: ldam|AH70\_05745 pent|YP\_803814  
is1537: ldam|AH70\_06360 pent|YP\_804875  
is1538: ldam|AH70\_06420 pent|YP\_803872  
is1539: ldam|AH70\_06425 pent|YP\_803871  
is1540: ldam|AH70\_06430 pent|YP\_803870  
is1541: ldam|AH70\_06435 pent|YP\_803869  
is1542: ldam|AH70\_06465 pent|YP\_804354  
is1543: ldam|AH70\_06685 pent|YP\_804782  
is1544: ldam|AH70\_06850 pent|YP\_805126  
is1545: ldam|AH70\_07120 pent|YP\_805196  
is1546: ldam|AH70\_07125 pent|YP\_805197  
is1547: ldam|AH70\_07130 pent|YP\_805198  
is1548: ldam|AH70\_07135 pent|YP\_805199  
is1549: ldam|AH70\_07190 pent|YP\_803856  
is1550: ldam|AH70\_07195 pent|YP\_805261  
is1551: ldam|AH70\_07200 pent|YP\_805263  
is1552: ldam|AH70\_07295 pent|YP\_804029  
is1553: ldam|AH70\_07475 pent|YP\_804018  
is1554: ldam|AH70\_07810 pent|YP\_805040  
is1555: ldam|AH70\_07860 pent|YP\_805038  
is1556: ldam|AH70\_08050 pent|YP\_804112  
is1557: ldam|AH70\_08090 pent|YP\_803592  
is1558: ldam|AH70\_08105 pent|YP\_804845  
is1559: ldam|AH70\_08225 pent|YP\_804059

---


is1560: ldam|AH70\_08305 pent|YP\_803660  
is1561: ldam|AH70\_08335 pent|YP\_803894  
is1562: ldam|AH70\_08365 pent|YP\_804168  
is1563: ldam|AH70\_08410 pent|YP\_805192  
is1564: ldam|AH70\_08545 pent|YP\_804153  
is1565: ldam|AH70\_08775 pent|YP\_805275  
is1566: ldam|AH70\_09005 pent|YP\_804701  
is1567: ldam|AH70\_09060 pent|YP\_805247  
is1568: ldam|AH70\_09345 pent|YP\_804481  
is1569: ldam|AH70\_09350 pent|YP\_804482  
is1570: ldam|AH70\_09360 pent|YP\_804484  
is1571: ldam|AH70\_09380 pent|YP\_804489  
is1572: ldam|AH70\_09385 pent|YP\_804490  
is1573: ldam|AH70\_09400 pent|YP\_804492  
is1574: ldam|AH70\_09405 pent|YP\_804493  
is1575: ldam|AH70\_09410 pent|YP\_804494  
is1576: ldam|AH70\_09415 pent|YP\_804495  
is1577: ldam|AH70\_09420 pent|YP\_804496  
is1578: ldam|AH70\_09425 pent|YP\_804497  
is1579: ldam|AH70\_09440 pent|YP\_804500  
is1580: ldam|AH70\_09450 pent|YP\_804502  
is1581: ldam|AH70\_09560 pent|YP\_804252  
is1582: ldam|AH70\_09760 pent|YP\_805140  
is1583: ldam|AH70\_09765 pent|YP\_803741  
is1584: ldam|AH70\_09795 pent|YP\_803730  
is1585: ldam|AH70\_09930 pent|YP\_803978  
is1586: ldam|AH70\_09965 pent|YP\_803985  
is1587: ldam|AH70\_10085 pent|YP\_803752  
is1588: ldam|AH70\_10225 pent|YP\_803745  
is1589: ldam|AH70\_10235 pent|YP\_804466  
is1590: ldam|AH70\_10480 pent|YP\_804994  
is1591: ldam|AH70\_10890 pent|YP\_805054  
is1592: ldam|AH70\_11045 pent|YP\_803788  
is1593: ldam|AH70\_11050 pent|YP\_803789





## Part IV

# General Discussion and Future Perspectives





# Preamble

The goals of this thesis were to evaluate the state-of-the-art approaches for microbial diversity analyses of food ecosystems and for the description of novel LAB taxa. This Part presents a GENERAL DISCUSSION of the results obtained and provides FUTURE PERSPECTIVES.

In **Chapter 6**, the effectiveness of culture-dependent and culture-independent molecular approaches, traditionally used in microbial diversity analyses of food ecosystems, was assessed. Furthermore, the value of HT sequencing technologies, which were applied to unravel the microbial diversity of mature Belgian red-brown acidic ales (Section 5.1) and to sequence the genome of the dominant bacterial member (*i.e.*, *Pediococcus damnosus* LMG 28219) of this beer ecosystem (Section 5.2), was evaluated. Finally, perspectives on the importance of cultivation and the application of an integrative ecosystem biology approach in future microbial diversity analyses of food ecosystems were provided.

In **Chapter 7**, the value of the traditionally used polyphasic taxonomy approach, which was implemented to characterize several novel LAB species (Sections 4.1, 4.2, and 4.3), was assessed. For the genus *Carnobacterium*, this characterization included the extension of the existing MLSA scheme (Section 4.3). Furthermore, Chapter 7 includes an evaluation of the possibilities of incorporating whole-genome sequences into the description of novel LAB species.

Finally, **Chapter 8** comprises guidelines on how HT sequencing technologies should be integrated in future microbial diversity analyses of food ecosystems and outlines which role the genomic taxonomy could have in these analyses.



# 6

## Considerations for Microbial Diversity Analyses

### 6.1 Traditional Methods

A variety of culture-dependent and culture-independent molecular techniques are traditionally used in microbial diversity analyses to study the phylogenetic and functional diversity of food ecosystems. The culture-dependent approach, involving plating on a range of empirically chosen media and conditions followed by dereplication using MALDI-TOF MS and identification of representative isolates using 16S rRNA gene sequencing, is labor-intensive and reveals only a fraction of the actual diversity. This is due to the challenging recovery of microorganisms, which are either stressed or entered into a VBNC state (Fleet, 1999; Rappé & Giovannoni, 2003). As a consequence, this empirical approach is inadequate for the complete characterization of microbial communities. On the other hand, culture-independent molecular techniques (*i.e.*, DGGE, RFLP, ARISA, qPCR, FISH, clone libraries, and others) do not depend on the cultivability of microorganisms in a community, but the limitations associated with these methods are their labor-intensity, their low-throughput, and their time-consuming nature. Furthermore, the Sanger DNA-sequencing technology is inadequate for processing complex and/or multiple samples. Analysis of these samples requires the capacity to read DNA from

multiple templates in parallel, which HT sequencing technologies do effectively at low costs.

## 6.2 Methods Involving HT Sequencing

HT sequencing technologies have become indispensable tools for studying the diversity and the metabolic potential of environmental microorganisms. Because of the biases introduced during microbial diversity analyses applying HT sequencing technologies (Ercolini, 2013; Temperton & Giovannoni, 2012), it is crucial to consider those before launching an experiment and while handling the data. In this chapter, several pre- and post-sequencing considerations are discussed and related to the experimental work done in the framework of the present study.

### 6.2.1 Pre-Sequencing Considerations

#### General Biases

As is the case for most of the cultivation-independent methods, it is of utmost importance to obtain metagenomic DNA that represents the actual diversity of a sample of interest. This implies minimizing the number of contaminants and inhibitory substances, while maximizing DNA yield of all community members. The most challenging aspect in the DNA extraction process is cell lysis, as not all members of the community (*i.e.*, Gram-stain-positive bacteria, Gram-stain-negative bacteria, fungi, and/or others) are lysed with the same efficacy, which will inevitably introduce biases in the community composition observed. The DNA extraction protocol, as utilized in Section 5.1, applied a combination of freezing and thawing, enzymatic lysis, and physical breaking to release genomic DNA from the cells and was capable of capturing both Gram-stain-positive (*e.g.*, *Pediococcus*, *Lactobacillus*), Gram-stain-negative (*e.g.*, AAB), and fungal (*e.g.*, *Dekkera*) DNA. An alternative approach is to separate prokaryotic and eukaryotic cells by filtration, followed by the application of DNA extraction procedures optimized for the filtered fractions. However, this alternative approach will distort the actual community structure of the sample. As a consequence of the concerns highlighted above, prior knowledge on the microbial community composition is crucial to determine the optimal DNA extraction procedure for the sample of interest.

Next to the biases associated with metagenomic DNA extraction, applying PCR will inevitably introduce biases to estimations of the actual microbial community diversity. Issues regarding primer universality, chimera formation, PCR errors, non-target and preferential amplification, primer and target-region choice, and others, are common to all PCR-based molecular methods. These biases can be minimized by, for instance, optimization of the PCR protocol, which reduces the incidence of chimera formation by applying a limited number of amplification cycles and a long elongation time (as was applied in Section 5.1). Furthermore, a PCR polymerase with proofreading activity was used in Section 5.1 to minimize the PCR error rate. Another major issue is preferential amplification of shorter fragments when sequencing the ITS region, which shows considerable length variation across fungal taxa. Moreover, the primer and target-region choice will result in phylogenetic analyses into varying taxonomic depths, depending on the length and region of the retrieved partial 16S rRNA gene sequence (Kunin *et al.*, 2008; Engelbrektson *et al.*, 2010).

## Habitat

The biodiversity of a habitat, including both species richness (*i.e.*, total number of species) and species evenness (*i.e.*, relative abundance of each of the community members), influences the degree of difficulty of the microbial diversity analyses applying HT sequencing technologies. For metagenomic shotgun sequencing strategies involving assembly, habitats with few abundant microbial species are better targets than habitats with many species of even abundance (Kunin *et al.*, 2008). Furthermore, samples containing a large amount of eukaryotic DNA, as was the case for the Belgian red-brown acidic ale ecosystem (Section 5.1), will require extensive sequencing, elevating the actual costs and complicating downstream data analysis and storage. Because of the issues highlighted above, pre-discovery of the ecosystem of interest is recommended prior to launching metagenomic shotgun sequencing experiments. Preceding microbial diversity analysis using target enrichment strategies in conjunction with abundance estimations (*e.g.*, qPCR) can aid to properly assess the most suited approach to analyze the ecosystem of interest.

As discussed in Section 5.1, the mature Belgian red-brown acidic ale ecosystem was analyzed using 454 pyrosequencing of the partial 16S rRNA gene and ITS1 region and revealed that this ecosystem contained only one dominant bacterial and one dominant fungal member, next to several less abundant microbial members. Subsequent targeted isolation and whole-genome sequencing of the dominant community

members allows the study of the metabolism and gene repertoire of these isolates, providing valuable information on their eco-physiological roles (as was done for the dominant bacterial community member in Section 5.2). In case of cultivation problems with the community members of interest, sequencing the metagenome or the genome of single microbial cells (Kalisky & Quake, 2011) could be valuable alternatives. However, if the major objective of the study is to target the whole community diversity and function, instead of focusing on the dominant members, deep metagenome sequencing may be the preferred solution (van Hijum *et al.*, 2013).

## Platform

In earlier times, sequencing of 16S rRNA gene amplicons mostly involved 454 pyrosequencing (as was applied in Section 5.1), whereas now many studies use Illumina sequencing because of its higher throughput and resulting reduced expenses (Caporaso *et al.*, 2010). A valid concern regarding the use of the Illumina platform for sequencing 16S rRNA gene amplicons is that the single-end Illumina reads are less than half the length of the 454 pyrosequencing reads, at least at the time of writing. Nevertheless, Illumina sequencing can take advantage of its higher throughput capacity and its ability to generate paired-end (PE) reads in downstream analyses, enabling PE read assembly into a consensus sequence. Because of the conserved nature of the 16S rRNA gene (1500 bp) (as shown in Sections 4.1, 4.2, and 4.3), sequencing of the partial 16S rRNA gene using 454 pyrosequencing or Illumina sequencing does not allow to distinguish between closely related LAB species and only provides a tentative identification. Therefore, complete or nearly complete sequencing of the 16S rRNA gene would be valuable for future research, with the PacBio RS platform being promising in this respect.

A similar evolution in the choice of platform was apparent for shotgun sequencing of (meta)genomic DNA. The initially most often used 454 pyrosequencing is gradually replaced by the increasing use of Illumina sequencing (as was applied for genome sequencing in Section 5.2), providing shorter reads but a much higher throughput and hence coverage at the same price. A recent comparative study of a freshwater lake planktonic community has shown that Illumina sequencing and 454 pyrosequencing lead to similar results with respect to assemblies and the taxonomic and functional repertoires covered (Luo *et al.*, 2012). Next to 454 pyrosequencing and Illumina sequencing platforms, the PacBio RS platform could be promising, because it generates read lengths that are long enough to span multiple prokaryotic genes



and thus are able to provide reliable genetic contexts and to improve annotations. Despite the high error rate inherent to use of the PacBio RS platform, the errors are mostly random and can be reduced in an almost linear fashion by increased coverage, which contrasts with the inherent systematic errors (*i.e.*, error accumulation at the end of the sequence) of other platforms. Currently, a combination of long- and short-read technologies constitutes a particularly promising approach in future (meta)genomics that bears the potential to significantly advance the field.

## Replication

Although HT sequencing technologies bring extensive sample replication and subsequent robust statistical analyses into reach, this is rarely done in microbial ecology studies (Prosser, 2010). It is a good scientific practice to analyze replicates (*i.e.*, biological, technical, and experimental replicates) of a sample and assess whether observed differences are statistically meaningful. Replication implies that researchers should critically assess the techniques and experimental design used and determine whether they are capable of achieving the aims of the experiment (Knight *et al.*, 2012). Nevertheless, sampling replicates is almost impossible because of spatial variability occurring in many habitats (Prosser, 2010). Hence, comparing such alleged replicates may reveal little information on methodological reproducibility.

The research presented in Section 5.1 included biological replicates (*i.e.*, subsequent brews) and technical variation was taken into consideration by performing DNA extractions and PCRs in triplicate and pooling them afterwards. A drawback associated with pooling of replicates is that it destroys spatial variability and makes it impossible to calculate true variability, which is required for comparisons (Prosser, 2010). In Section 5.1, experimental replication was applied for the generous donor sample (*i.e.*, a DNA sample that was previously sequenced) and showed a high reproducibility of 454 pyrosequencing. Similarly, Luo and colleagues (2012) assessed the reproducibility of 454 pyrosequencing and Illumina sequencing by re-analyzing the same microbial community DNA sample and showed that library preparation and sequencing are highly reproducible. Nevertheless, Schloss and coworkers (2011a) measured intra- and inter-sequencing center variation and revealed that their mock community (*i.e.*, artificial community) samples clustered by sequencing center and by sequencing run, demonstrating the value of replicating samples in microbial ecology studies.

## Contextual Data

Contextual data are the data associated with microbial diversity analyses, such as habitat description, measures of environmental parameters, sampling procedure, sample storage, and others. The genomic standards consortium (GSC) (Field *et al.*, 2011) has published standards for the minimum information about a (meta)genome sequence [MIMS/MIGS (Field *et al.*, 2008; Kottmann *et al.*, 2008)] and about a marker gene sequence [MIMARKS (Yilmaz *et al.*, 2011)], as part of the minimum information about any sequence (MlXs) standards and checklist (Yilmaz *et al.*, 2011), which are supported by the international nucleotide sequence databases collaboration (INSDC). It is of utmost importance that contextual data are collected and integrated into databases (as was done for the data generated in Sections 5.1 and 5.2), because in the long run these data will allow to extract correlations between geography, time, prevailing environmental conditions, and functions from data that otherwise would never be uncovered (Yilmaz *et al.*, 2011; Delmont *et al.*, 2011; Knight *et al.*, 2012).

### 6.2.2 Post-Sequencing Considerations

Until recently, sequencing capacity has been the limiting factor in microbial diversity analyses involving HT sequencing. However, the on-going increase in sequencing capacity and resulting reduction of prices have made post-sequencing data analysis the main bottleneck. Although progress in sequencing technologies still continues at an exponential pace, data analysis cannot keep up. As a consequence, the cost of sequencing declines continuously, whereas the cost for bioinformatics data analysis increases (Sboner *et al.*, 2011).

### Use of the 16S rRNA Gene for Taxonomic Purposes

About one in thousand genes in a metagenomic shotgun sequencing dataset is a 16S rRNA gene and only a fraction of these reads has sufficient length and quality for subsequent phylogenetic analysis. Furthermore, depending on the length and region of the partial 16S rRNA gene sequence retrieved, phylogenetic analysis results into OTUs with varying taxonomic depths (Kunin *et al.*, 2008). Extracting 16S rRNA gene reads from a metagenome dataset is superior to targeted sequencing of 16S rRNA gene amplicons in terms of PCR biases, because no primer selection and amplification are involved for library preparation. Nevertheless,

when used for phylogenetic analysis, metagenomic shotgun sequencing requires extensive sequencing depth, resulting in complex data handling and higher prices compared to target enrichment strategies. Furthermore, the highly conserved nature of the 16S rRNA gene in closely related LAB species complicates identification at species levels (as discussed above). Alternatively, as sequence analysis of protein-encoding genes has proven to be superior for the accurate species level identification of a variety of LAB and AAB [(Naser *et al.*, 2005a; Cleenwerck *et al.*, 2010; De Bruyne *et al.*, 2007) and as shown in Sections 4.1, 4.2, and 4.3], mining a metagenomic shotgun dataset for housekeeping genes could aid in obtaining species-level identifications.

## Error Removal

In genome sequencing projects, highly redundant consensus assemblies compensate for sequencing errors (as was applied in Section 5.2). Similarly, assembly of sequences from metagenomic libraries can result in a good draft or even complete genomes (Tyson *et al.*, 2004) when the target species shows little intraspecies variation, but usually requires a substantial amount of sequencing. In addition, assembly of large metagenomic datasets presents challenges with respect to memory requirements. Although the assembly of metagenomes yields longer sequences, it also bears the risk of creating chimeric contigs, particularly in habitats with closely related species or highly conserved sequences occurring across species (Teeling & Glöckner, 2012). Furthermore, assembly distorts abundance information, as overlapping sequences from an abundant species will be identified as belonging to the same genome and consequently joined. This leads to a relative underrepresentation of sequences of abundant species. Hence, gene frequencies are better compared based on read representation rather than on assemblages (Teeling & Glöckner, 2012).

In contrast, target enrichment strategies based on PCR (as applied in Section 5.1) cannot take advantage of consensus assemblies to mask incorrect base calls (Huse *et al.*, 2010). As a consequence, caution must be taken when analyzing datasets that do not rely on assembled reads, because sequencing errors can lead to the artificial inflation of diversity estimates (Kunin *et al.*, 2010) and to erroneous taxonomic assignments. A feasible but imperfect alternative to building consensus sequences for target enrichment strategies is to identify and remove reads that are likely to be incorrect. Different HT sequencing platforms are prone to different types of errors, with homopolymers being frequently associated with 454 pyrosequencing (as was applied in Section 5.1).

Bioinformatics' solutions exist to remove PCR and sequencing errors in target enrichment datasets, with the latter having a higher incidence compared to the former. Schloss and coworkers (2011a) identified several features that are linked to low-quality 454 pyrosequences. For instance, sequences containing homopolymers, ambiguous bases, mismatches in their barcode or primer sequence, and short sequences are more likely to be of low quality. The parameters applied in Section 5.1 were largely similar to those proposed by Schloss and coworkers (2011a), with the exception of several parameters for which more stringent values based on the analysis of a mock community were preferred. Moreover, error accumulation at the end of the reads makes trimming of raw 454 pyrosequencing reads crucial to reduce the overall error rate, although there is still much debate on how stringent this trimming should be (Schloss *et al.*, 2011a; Quince *et al.*, 2011). For instance, the raw reads of approximately 500 bp provided by the 454 GS FLX Titanium system of Roche Diagnostics (as applied in Section 5.1) were trimmed to a length of approximately 250 bp, using the parameter settings proposed by Schloss and coworkers (2011a). Further reduction of the error rate can be achieved by single-linkage pre-clustering (Huse *et al.*, 2010), SeqNoise (Quince *et al.*, 2011) (applied in Section 5.1), or modifications thereof. The analysis of reads generated by the Illumina MiSeq (250 bp) differs in several aspects from those generated by the 454 GS FLX system, because Illumina can take advantage of the PE read assembly, which can be used as a tool to reduce the sequencing error rate. However, Kozich and coworkers (2013) showed that these PE reads should fully overlap to obtain good-quality reads.

Subsequently, the de-noised and trimmed sequences are preferably aligned against a curated reference alignment, which is available for 16S rRNA gene sequences (*e.g.*, the SILVA reference alignment was used in Section 5.1) (Schloss, 2009). Sequences generated due to aspecific amplification and therefore aligning to a wrong region or sequences that were classified as chloroplasts or mitochondria, were removed from the dataset. Furthermore, singletons (*i.e.*, OTUs containing only one sequence) are often sequencing errors and may lead to the artificial inflation of the 'rare biosphere' diversity (*i.e.*, microorganisms present in extremely low abundances) and thereby affect the conclusions (Dickie, 2010; Reeder & Knight, 2009). A lot of debate exists on whether the members of this 'rare biosphere' have ecological significance or should be removed from the dataset (Dickie, 2010; Reeder & Knight, 2009).

Although several steps can be taken prior to sequencing to reduce the rate of chimerism (Acinas *et al.*, 2005; Haas *et al.*, 2011; Thompson *et al.*, 2002) as discussed above, several bioinformatics approaches have been developed to identify and remove chimeras. For instance, Haas and coworkers (2011) developed the ChimeraSlayer algorithm, which was superior to Bellerephon (Huber *et al.*, 2004) and Pintail (Ashelford *et al.*, 2005), relying on a reference database to distinguish chimeric and non-chimeric sequences. Additionally, Quince and colleagues (2011) developed Perseus, which does not use a reference database but requires a training set of sequences similar to the sequences being characterized. Finally, Edgar and coworkers (2011) developed Uchime (as applied in Section 5.1), showing improved performance over ChimeraSlayer, but comparable to that of Perseus.

The mothur software package ([www.mothur.org](http://www.mothur.org), as applied in Section 5.1) is an open-source, platform-independent, community-supported software for describing and comparing microbial communities (Schloss, 2009). It builds upon previous tools [e.g., Amplicon Noise (Quince *et al.*, 2009), single-linkage pre-clustering (Huse *et al.*, 2010), Perseus (Quince *et al.*, 2011), Uchime (Edgar *et al.*, 2011), and others] and provides a flexible and powerful software package for analyzing sequencing data. Furthermore, they present a pipeline for handling 454 and Illumina 16S rRNA gene sequencing data (Schloss *et al.*, 2011a; Kozich *et al.*, 2013). QIIME (Caporaso *et al.*, 2010) (*i.e.*, quantitative insights into microbial ecology) is a valid alternative for the analysis of target enrichment sequencing data.

## Database Dependencies

A major concern in microbial diversity analyses involving HT sequencing is that often a significant proportion of the data cannot be assigned to a taxon due to a lack of close matches in reference databases (Qin *et al.*, 2010; Temperton & Giovannoni, 2012). The primary source of annotated nucleotide sequences is the INSDC resource (Nakamura *et al.*, 2013), which includes nucleic acid sequencing data and associated information from the DDBJ (Sugawara *et al.*, 2007), the European nucleotide archive (ENA) (Leinonen *et al.*, 2011), and GenBank (Benson *et al.*, 2013), including the SRA (Kodama *et al.*, 2012) repository. These sequence data collections constitute huge resources of sequence information but, at the same time, their contents are often compromised by identification errors, low-quality sequence data, redundancy, and incompleteness (Turenne *et al.*, 2001). Current functional assignment of genes from (meta)genomes is based on time-consuming homology searches, using BLAST tools (Altschul *et al.*, 1997) (as was applied in

Section 5.2) that depend heavily on the quality and completeness of these available databases. Furthermore, homology-based algorithms are not sensitive, particularly for genomes that lack sequenced relatives and can miss novel genes that may ecologically be the most interesting (Suenaga, 2012). A review of prokaryotic protein diversity in different shotgun metagenomic studies indicated that 30–60% of the proteins cannot be assigned to known functions using current public databases (Vieites *et al.*, 2009), indicating that there is an urgent need to develop specific and quality controlled reference databases that are able to reduce redundancies, validate sequence annotations, and increase accuracy and effectiveness of taxon assignments (Santamaria *et al.*, 2012; Walker *et al.*, 2014).

The approach applied in Section 5.1 for the annotation of the 16S rRNA gene and ITS region sequence data, was based on what has been proposed by Schloss and coworkers (2009; 2010; 2011a). Firstly, reads were binned (*i.e.*, grouping of reads into clusters) into OTUs and subsequently a consensus classification was obtained for each OTU using a trainset (*i.e.*, quality controlled reference sequences), which is available for 16S rRNA gene sequences on the mothur website to train the Naïve Bayesian Classifier (Wang *et al.*, 2007). Unfortunately, no trainset was available for fungal ITS sequences and was therefore developed in-house using reference sequences extracted from curated ITS databases (Weiss *et al.*, 2013; Robert *et al.*, 2013) (as applied in Section 5.1). Another issue complicating the taxonomic assignment of fungal ITS sequences is the lack of means to refer to fungal species, for which no Latin name is available in a standardized way. These shortcomings negatively affect the identification of fungal species based on ITS markers and result in many unclassified fungi, which was apparent in the results of Section 5.1. As a consequence, the development of curated fungal ITS databases will be crucial for future research on fungal diversity.

## 6.3 Importance of Cultivation

Although purely sequence-based descriptions give valuable information about community content and metabolic potential, they suffer from several inherent limitations (as discussed in Section 6.2). As a consequence, the cultivation of microorganisms will remain essential for future research (Allen-Vercoe, 2013). One benefit of having strains in pure culture is that their genome can be sequenced rapidly and at low-cost per base, thus adding information to reference databases (Walker *et al.*, 2014) and contributing to the interpretation of physiological as well as (meta)genomic data. Furthermore, reference genome data derived from food isolates

can be mined for interesting functionalities to apply in the food industry. Of course, a microorganism can only be applied in the food industry after it has first been grown, isolated, and tested for safety in the laboratory. In addition, it is becoming increasingly clear that microbial ecosystems contain rare phylotypes for which abundance is less important than the ability to serve a key metabolic function upon which the larger community structure may rely (Ze *et al.*, 2013). Culture-based approaches offer the route through which sampling and detailed analysis of these low-abundance species may be accomplished.

Cultivation of microbial community members *in vitro* can be approached in several ways, which are summarized in Figure 22. The first of these is an empirical approach (as discussed in Section 1.1), which traditionally involves the cultivation of microorganisms onto a set of empirically chosen growth media and conditions. The diversity yield using this approach is low because of the enormous range of microbial growth rates and conditions that unknown targets may use, which makes the approach time-consuming and inefficient (Allen-Vercoe, 2013).

Another approach to cultivate microbial communities requires previous knowledge on the ecosystem of interest and involves the use of custom-made growth media for targeted growth of microorganisms that have thus far evaded cultivation (Allen-Vercoe, 2013). In this targeted bioinformatics approach, the molecular signatures of a microbial target are studied before culture is attempted to infer culture conditions. The resulting diversity yield is intentionally low and the target microbiota is efficiently recovered. To our knowledge, this approach has not been applied to a food fermentation ecosystem. An example from research on the gut microbiota is the isolation of *Akkermansia muciniphila* from feces through its unusual ability to use mucin as the sole carbon and nitrogen source (Derrien *et al.*, 2004). An even more targeted approach would be to utilize the genome of a given uncultured species to infer its necessary growth requirements [example publication: (Livermore *et al.*, 2014)].

The third approach to culture microorganisms that thus far evaded cultivation has been developed by Lagier and coworkers (2012) and named culturomics. They started with eliminating the predominant microbial population from stool samples using cocktails of lysogenic phages or filtration methods. This was followed by growth enrichment steps with blood, rumen fluid, or fecal water, and finally 212 separate culture conditions were applied to the resulting pool of microorganisms that had been diluted to give single cells. This approach enabled cultivation of a significant number of low abundant species, which were previously missed by

HT sequencing approaches. The main bottleneck of the culturomics method is the identification step, which is normally performed through time-consuming 16S rRNA gene sequencing of isolates. However, Lagier and colleagues (2012) were able to demonstrate that MALDI-TOF MS (as was applied for dereplication and identification in Sections 4.2 and 4.1, respectively) is a fast and cheap alternative, greatly facilitating the work. Combined with automated colony picking technology and HT growth arrays, culturomics represents a powerful technique that can revolutionize microbial cultivation technology. This approach has the potential to generate a huge amount of bacterial isolates, which may be used to define interactions between bacterial species and between bacterial and eukaryotic cells that are difficult to predict unless tested biologically.




Figure 22: Schematic outlining of the general approaches to culturing microbial diversity

(Allen-Vercoe, 2013)



## 6.4 Integrative Ecosystems Biology

The present thesis focused on microbial community analyses involving HT sequencing technologies, which were based on DNA and therefore cannot distinguish between metabolically (in)active and dead microorganisms. Important functional information of the metabolically active fraction of a community can be gained from the fields of metatranscriptomics and metaproteomics. These have the inherent advantage to reveal functional activity rather than predicted functional capabilities that are derived from (meta)genomic analyses. Combining these approaches with metagenomics and exhaustive cultivation efforts (Allen-Vercoe, 2013) (Figure 22) could provide new insights into relations between members of a microbial community and will be important for designing interventions targeted at functions of the microbial communities in food rather than specific constituent species. Furthermore, metagenome data can be integrated with (meta)-metabolome data (Turnbaugh & Gordon, 2008). Such integrative ecosystems biology studies [*e.g.*, (Shi *et al.*, 2011; Teeling *et al.*, 2012)] will pose new challenges due to their increasing data complexities, in particular with respect to bioinformatics post-processing. To cope with these huge amounts of data, there will be a growing need for automated pipelines enabling bioinformaticians to perform the analysis and to interpret the outcome.



# 7

## LAB Taxonomy in a HT Sequencing Era

### 7.1 Polyphasic Taxonomy

Polyphasic taxonomy of LAB comprises the application of comparative 16S rRNA gene sequence analysis, MLSA of less conserved phylogenetic markers, DNA-DNA hybridization, and phenotypic characterization to provide conclusive evidence about the taxonomic status of an organism (as applied in Sections 4.1, 4.2, and 4.3). Furthermore, the determination of the G+C content is part of the standard description of novel taxa, whereas the analysis of the peptidoglycan structure of the cell wall is more specific for LAB taxonomy. Different aspects make this polyphasic taxonomy approach time- and money-consuming, labor-intensive, and not accessible to all laboratories.

Due to the conserved nature of the 16S rRNA gene in LAB species, MLSA of less conserved genes is often required in LAB species descriptions. One aspect that makes MLSA unsuited for HT applications is the lack of universal primers and, as a consequence, novel primer design and PCR optimization are often required to extend the existing MLSA scheme to several LAB genera (as was put into practice for the genus *Carnobacterium* in Section 4.3). In Section 4.1, amplification of the

*rpoA* and *atpA* genes proved unsuccessful for many strains of the genus *Weissella* using the primers available, and therefore sequencing was restricted to the *pheS* gene. As a consequence, novel primer design and optimization of the PCR will be mandatory to extend the existing MLSA scheme to the genus *Weissella*.

Next to the issues related to sequencing of the 16S rRNA gene and the construction of an MLSA scheme for LAB, DDH experiments also have several inherent characteristics that make it unsuited to apply in HT. These include the requirement of large quantities of high-quality DNA, the time- and money-consuming and labor-intensive nature, the inability to build databases (Gevers *et al.*, 2005), and the lack of intra- and inter-laboratory reproducibility (Stackebrandt & Ebers, 2006; Rosselló-Mora, 2005).

Furthermore, a range of labor-intensive and time-consuming tests are utilized for the determination of phenotypic characteristics of novel taxa. Phenotypic characteristics that are part of the standard description of LAB taxa commonly include a positive Gram reaction, absence of endospores, oxidase and catalase activity (typically absent), glucose fermentation products (typically lactic acid), carbohydrate fermentation patterns, ratio of D- and L-lactic acid production, hydrolysis of aesculin and arginine, reduction of nitrate, gelatine liquefaction, growth at different temperatures, pH range values and NaCl concentrations, and tolerance to oxygen (Mattarelli *et al.*, 2014). Unfortunately, the tests to determine these phenotypic characteristics are time-consuming and labor-intensive and most of these phenotypic characteristics are poorly discriminatory among species and genera and mainly have a descriptive value in LAB taxonomy. Another phenotypic characteristic recommended for novel LAB species descriptions is the peptidoglycan type, which is species-specific for many LAB species. Nevertheless, the diversity in peptidoglycan type is very limited among species, with the Lys-D-Asp type being the predominant type within the genus *Lactobacillus*, *Pediococcus*, and several other LAB genera. The determination of the peptidoglycan type is not suited for HT applications because it is labor-intensive, time- and money-consuming, and not accessible to all laboratories.

As discussed above, the polyphasic taxonomy approach to describe novel LAB taxa is not suited to perform with HT capacity, which is contradictory with the huge amount of strains that are currently awaiting description and formal naming (Vandamme & Peeters, 2014). Furthermore, this approach contrasts with the emerging culturomics approaches, which will generate a huge amount of not yet characterized strains. Therefore, the traditionally used polyphasic taxonomy approach is outdated and should urgently be reconsidered to enable fast description of novel LAB taxa.

## 7.2 Genomic Taxonomy

With the advent of HT sequencing technologies that are becoming cheaper and the development of many tools that guide bioinformaticians during the assembly and annotation of whole-genome sequences, routine sequencing and downstream analysis come into reach of many research groups. Because whole-genome sequences provide insights into the genetic nature of microbial species, they can be used as a tool for delineating bacterial species and for studying their phylogeny.

LAB genome sequences are continuously accumulating, providing a good opportunity to study the evolutionary history of LAB. For instance, sequences of the 16S rRNA gene and less conserved genes for MLSA can be extracted out of the whole-genome sequence, evading current issues related to primer universality and PCR optimizations (as discussed above). Furthermore, whole-genome sequence-based approaches are superior for studying phylogeny, because of the better resolution of whole-genome sequences for discriminating both distantly and closely related bacteria compared to single or multiple genes (Vandamme & Peeters, 2014). In addition, the G+C content can be calculated from the whole-genome sequence, making the application of HPLC approaches obviate for G+C content analyses.

While the DDH species threshold is being translated into MLSA or whole-genome based thresholds (Gevers *et al.*, 2005; Goris *et al.*, 2007; Richter & Rossello-Mora, 2009; Tindall *et al.*, 2010), the reassessment of bacterial taxonomy and the species definition will require more than a methodological translation of threshold levels (Achtman & Wagner, 2008). In 1987, the ad hoc committee on reconciliation of approaches to bacterial systematics (Wayne *et al.*, 1987) stated that taxonomy should be determined phylogenetically and that the complete genome sequence should therefore be the standard for species delineation. As DDH experiments were originally developed because whole-genome sequences were not available, the current accumulation of whole-genome sequences enables to abandon DDH experiments out of the polyphasic taxonomy approach and to establish taxonomic schemes based on evolutionary information contained in whole-genome sequences (Eisen, 2000; Wolf *et al.*, 2001). Comparing whole-genome sequences have many advantages compared to DDH experiments, such as the ability to build databases and to perform these comparisons in HT.

Furthermore, whole-genome sequences yield information on the metabolic potential of a particular strain (as was applied in Section 5.2), which could make extensive phenotypic tests obviate. Nevertheless, to date, complete genome sequences and improved insights from genome annotations have not yet resulted in reliable predictions of metabolic and chemotaxonomic features, but it is likely that this will occur at some point in the future (Oren & Garrity, 2014). However, it is not known if these properties are actually expressed in the phenotype of the organism as it grows in the laboratory or in its natural environment. Therefore, at the time of writing, analysis of whole-genome sequences cannot be considered a proxy for the phenotypic characterization contained in novel species descriptions.

### 7.3 Integrating Genomics into LAB Taxonomy

There is a strong consensus that prokaryotic classification should be based on a polyphasic taxonomy approach, integrating data from traditional physiological and chemotaxonomic tests with complementary results from genetic methods (Vandamme *et al.*, 1996; Rossello-Mora & Amann, 2001; Stackebrandt *et al.*, 2002; Schleifer, 2009). Recently, Mattarelli and coworkers (2014) have provided the recommended minimal standards for description of new taxa of the genera *Bifidobacterium*, *Lactobacillus*, and related genera. Explicit in their guidelines is an emphasis that novel LAB taxa should be characterized as comprehensive as possible based on phenotypic, genotypic, and ecological characteristics. Nevertheless, how much characterization is considered enough?

Recently, Vandamme and Peeters (2014) proposed to use a whole-genome sequence and a minimal description of phenotypic characteristics as a basic biological identity card that be considered sufficient, cost-effective, and appropriate for species descriptions, which is a workable option for future large-scale descriptions of novel LAB species. In my opinion, novel LAB species descriptions should include a draft genome sequence accompanied with contextual data (*e.g.*, elaborated description of the sampling site, growth requirements, isolation strategies, date of sampling, ...) and data with taxonomic relevance (*e.g.*, the analysis of carbohydrate fermentation patterns and of glucose fermentation products, the ratio of D- and L-lactic acid production, Gram stain, catalase and oxidase activity). Phenotypic characteristics without or with limited taxonomic relevance, (*e.g.*, cell and colony morphology) should be excluded from LAB species descriptions, preventing taxonomists to perform a plethora of time- and money-consuming physiological tests of questionable

taxonomic relevance (Sutcliffe *et al.*, 2012). Furthermore, the LAB draft genome sequences should be subjected to a range of *in silico* calculations, including ANI, G+C content, phylogenetic trees of the core gene set, the prediction of metabolic pathways of peptidoglycan synthesis, ...

These basic biological identity cards should be stored and maintained in high-quality, readily accessible, iterative, and adaptable taxonomic databases (Sutcliffe *et al.*, 2012). The accumulation of these curated data will enable the determination of thresholds for species delineation and provide valuable resources for different research areas. The re-evaluation of how novel taxa are described will undoubtedly demand online publication formats, coupled to a more effective use of online databases to allow the iterative comparison of prokaryotic characteristics. Finally, these online taxonomic databases should consider how to deal with genome sequences of strains that cannot be cultivated, and hence cannot be deposited into culture collections.





# 8

## Future Perspectives

This Chapter provides perspectives on the integration of HT sequencing technologies in future microbial diversity analyses of food ecosystems and outlines the possible role of genomic taxonomy to improve the quality of microbial diversity analyses. Figure 23 proposes a workflow to characterize the microbial ecosystem contained in a food sample of interest by merging the fields of genomic taxonomy, reference database construction and management, (meta)genomics, (meta)transcriptomics, (meta)-metabolomics, and culturomics, which have the potential to collectively provide a comprehensive, integrative analysis of the food ecosystem of interest.

If a food sample with an unknown microbial ecosystem is provided, the starting point is the extensive collection of contextual metadata (as described in Section 6.2) and the sampling of at least three biological and technical replicates, if possible (Figure 23). These replicates are subsequently pre-discovered using a target enrichment strategy, which includes amplification of the bacterial 16S rRNA gene and the fungal ITS region. Following, these amplicons are subjected to HT sequencing using the platform of interest. This pre-discovery step will provide basic knowledge on the ecosystem of interest, which is crucial for subsequent in-depth analyses.




Figure 23: Microbial diversity analyses in a HT sequencing era

This Figure is extensively discussed in Chapter 8

Following, dependent on the outcome of the pre-discovery analysis and the research question, different approaches are preferred. (i) If the goal is to obtain an in-depth analysis of the low-abundant community members of the food ecosystem of interest, culturomics is a valuable approach (as discussed in Section 6.3). Knowledge on the identity of these low-abundant community members, which was previously obtained by the pre-discovery approach, is valuable to define a set of growth media and conditions that could promote the growth of these low-abundant community members and could discourage growth of the dominant community members. Robotic plating on an extensive set of growth media and conditions, followed by MALDI-TOF MS dereplication and identification using 16S rRNA gene sequencing, have the potential to significantly increase the throughput of this culturomics approach. (ii) If the goal is to target a limited number of community members of interest, the targeted bioinformatics approach is suited to isolate the target strain(s) (as discussed in Section 6.3). Similarly to the culturomics approach, the pre-discovery

step will provide information on the most suited strategy to isolate the target strain(s) of interest. The isolates obtained by the culturomics or the targeted bioinformatics approach [(i) & (ii)] can subsequently be subjected to whole-genome sequencing. In case of the inability to cultivate the target strain, the whole-genome sequence of that particular strain can be obtained by means of single-cell genomics or metagenome sequencing to infer its necessary growth requirements. These whole-genome sequences add interesting information to reference databases, which can be mined by other researchers. Another value of whole-genome sequencing of bacterial isolates is that functionalities can be assigned to specific microbial lineages, which contrasts with the lack of connectivity (*i.e.*, which species harbors which functionality) of metagenomic datasets (Walker *et al.*, 2014). As such, culture and phylogeny will continue to have crucially important roles in food fermentation microbiota research and will be required for the development of novel starter cultures. (iii) If one wants to provide a thorough functional analysis of the whole microbial community, metagenome sequencing is the approach preferred. The previously obtained knowledge on the species richness and composition will assist in determining how deep one needs to sequence to obtain enough coverage.

As (meta)genomics cannot distinguish between metabolically (in)active and dead microorganisms, the fields of (meta)transcriptomics and (meta)proteomics can reveal functional activity rather than predicted functional capabilities, derived from (meta)genomic analyses. Furthermore, combining (meta)genomics, (meta)transcriptomics, (meta)proteomics, and (meta)metabolomics with culturomics could provide new insights into relations between members of a microbial community and will be important for designing interventions targeted at functions of the microbial communities in food rather than specific constituent species. Automated pipelines will enable bioinformaticians to cope with the huge amounts of data and to analyze and interpret the outcome of these integrative ecosystem biology studies.


Although numerous large-scale genome sequencing studies have been performed, they have tended to focus on particular habitats of high interest to the scientific community or on the relatives of specific microorganisms. As a consequence, the currently available whole-genome sequences are limited by their highly biased phylogenetic distribution, creating a major obstacle in progressing towards a genome-based classification of microorganisms. Therefore, the Genomic encyclopaedia of bacteria and archaea (GEBA) project was launched aiming at reducing this bias (Klenk & Goker, 2010). They initiated a phylogenetically driven genome sequencing effort, selecting microorganisms based on their position in a phylogenetic tree of small subunit rRNA gene sequences. Although the GEBA project is already a

first attempt to correct for this bias, taxonomists could contribute by making the availability of a whole-genome sequence a prerequisite for the description of novel bacterial taxa.

Next to reducing the biased phylogenetic distribution of the whole-genome sequences that are currently available, switching to a genomic taxonomy enables the construction, curation, and maintenance of comprehensive taxonomic databases containing valuable contextual and taxonomic data, next to whole-genome sequences. This effort could aid in gene prediction of (meta)genomic datasets, because many gene finders require reference sequences from single species that are subsequently used to build a species-specific gene prediction model (Teeling *et al.*, 2012) (as discussed above). This will also have a major influence on the annotations of (meta)transcriptomic, (meta)proteomic, and (meta)-metabolomic data, which could improve the quality of the integrative ecosystem biology studies, and thereby bring future microbial diversity analyses of food ecosystems to a higher level. Furthermore, these curated taxonomic reference databases can be mined for strains with interesting functionalities, which could be of interest to the food industry.

# Part V

## Bibliography





- Abbott, D. A., Hynes, S. H., & Ingledew, W. M. (2005). Growth rates of *Dekkera/Brettanomyces* yeasts hinder their ability to compete with *Saccharomyces cerevisiae* in batch corn mash fermentations. *Applied Microbiology and Biotechnology*, 66, 641–647.
- Achtman, M., & Wagner, M. (2008). Microbial diversity and the genetic nature of microbial species. *Nature Reviews Microbiology*, 6, 431–440.
- Acinas, S. G., Marcelino, L. A., Klepac-Ceraj, V., & Polz, M. F. (2004). Divergence and redundancy of 16S rRNA sequences in genomes with multiple *rrn* operons. *Journal of Bacteriology*, 186, 2629–2635.
- Acinas, S. G., Sarma-Rupavtarm, R., Klepac-Ceraj, V., & Polz, M. F. (2005). PCR-induced sequence artifacts and bias: insights from comparison of two 16S rRNA clone libraries constructed from the same sample. *Applied and Environmental Microbiology*, 71, 8966–8969.
- Albert, T. J., Molla, M. N., Muzny, D. M., Nazareth, L., Wheeler, D., Song, X., Richmond, T. A., Middle, C. M., Rodesch, M. J., Packard, C. J., Weinstock, G. M., & Gibbs, R. A. (2007). Direct selection of human genomic loci by microarray hybridization. *Nature Methods*, 4, 903–905.
- Alegria, A., Szczesny, P., Mayo, B., Bardowski, J., & Kowalczyk, M. (2012). Biodiversity in Oscypek, a traditional Polish cheese, determined by culture-dependent and -independent approaches. *Applied and Environmental Microbiology*, 78, 1890–1898.
- Allen-Vercoe, E. (2013). Bringing the gut microbiota into focus through microbial culture: recent progress and future perspective. *Current Opinion in Microbiology*, 16, 625–9.
- Altschul, S. F., Madden, T. L., Schaffer, A. A., Zhang, J., Zhang, Z., Miller, W., & Lipman, D. J. (1997). Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Research*, 25, 3389–3402.
- Andorra, I., Monteiro, M., Esteve-Zarzoso, B., Albergaria, H., & Mas, A. (2011). Analysis and direct quantification of *Saccharomyces cerevisiae* and *Hanseniaspora guilliermondii* populations during alcoholic fermentation by fluorescence *in situ* hybridization, flow cytometry and quantitative PCR. *Food Microbiology*, 28, 1483–1491.
- Andres-Barrao, C., Benagli, C., Chappuis, M., Ortega Perez, R., Tonolla, M., & Barja, F. (2013). Rapid identification of acetic acid bacteria using MALDI-TOF mass spectrometry fingerprinting. *Systematic and Applied Microbiology*, 36, 75–81.
- Angiuoli, S. V., Gussman, A., Klimke, W., Cochrane, G., Field, D., Garrity, G., Kodira, C. D., Kyrpides, N., Madupu, R., Markowitz, V., Tatusova, T., Thomson, N., & White, O. (2008). Toward an online repository of standard operating procedures (SOPs) for (meta)genomic annotation. *OMICS*, 12, 137–141.
- Arena, M. P., Romano, A., Capozzi, V., Beneduce, L., Ghariani, M., Grieco, F., Lucas, P., & Spano, G. (2011). Expression of *Lactobacillus brevis* IOEB 9809 tyrosine decarboxylase and agmatine deiminase genes in wine correlates with substrate availability. *Letters in Applied Microbiology*, 53, 395–402.
- Ashelford, K. E., Chuzhanova, N. A., Fry, J. C., Jones, A. J., & Weightman, A. J. (2005). At least 1 in 20 16S rRNA sequence records currently held in public repositories is estimated to contain substantial anomalies. *Applied and Environmental Microbiology*, 71, 7724–7736.

- Aziz, R., Bartels, D., Best, A., DeJongh, M., Disz, T., Edwards, R., Formsma, K., Gerdes, S., Glass, E., Kubal, M., Meyer, F., Olsen, G., Olson, R., Osterman, A., Overbeek, R., McNeil, L., Paarmann, D., Paczian, T., Parrello, B., Pusch, G., Reich, C., Stevens, R., Vassieva, O., Vonstein, V., Wilke, A., & Zagnitko, O. (2008). The RAST server: rapid annotations using subsystems technology. *BMC Genomics*, 9, 1–15.
- Babot, J. D., Hidalgo, M., Arganaraz-Martinez, E., Apella, M. C., & Perez Chaia, A. (2011). Fluorescence *in situ* hybridization for detection of classical propionibacteria with specific 16S rRNA-targeted probes and its application to enumeration in Gruyere cheese. *International Journal of Food Microbiology*, 145, 221–228.
- Begerow, D., Nilsson, H., Unterseher, M., & Maier, W. (2010). Current state and perspectives of fungal DNA barcoding and rapid identification procedures. *Applied Microbiology and Biotechnology*, 87, 99–108.
- Behr, J., Israel, L., Gänzle, M. G., & Vogel, R. F. (2007). Proteomic approach for characterization of hop-inducible proteins in *Lactobacillus brevis*. *Applied and Environmental Microbiology*, 73, 3300–3306.
- Behr, J., & Vogel, R. F. (2010). Mechanisms of hop inhibition include the transmembrane redox reaction. *Applied and Environmental Microbiology*, 76, 142–149.
- Benson, D. A., Cavanaugh, M., Clark, K., Karsch-Mizrachi, I., Lipman, D. J., Ostell, J., & Sayers, E. W. (2013). GenBank. *Nucleic Acids Research*, 41, D36–42.
- Bergsveinson, J., Pittet, V., & Ziola, B. (2012). RT-qPCR analysis of putative beer-spoilage gene expression during growth of *Lactobacillus brevis* BSO 464 and *Pediococcus clausenii* ATCC BAA-344<sup>T</sup> in beer. *Applied Microbiology and Biotechnology*, 96, 461–470.
- Bermúdez-Humarán, L. G., Aubry, C., Motta, J.-P., Deraison, C., Steidler, L., Vergnolle, N., Chatel, J.-M., & Langella, P. (2013). Engineering lactococci and lactobacilli for human health. *Current Opinion in Microbiology*, 16, 278–283.
- Bernal, A., Ear, U., & Kyrpides, N. (2001). Genomes onLine database (GOLD): a monitor of genome projects world-wide. *Nucleic Acids Research*, 29, 126–127.
- Berry, D., Mahfoudh, K. B., Wagner, M., & Loy, A. (2011). Barcoded primers used in multiplex amplicon pyrosequencing bias amplification. *Applied and Environmental Microbiology*, 77, 7846–7849.
- Björkroth, K. J., Schillinger, U., Geisen, R., Weiss, N., Hoste, B., Holzapfel, W. H., Korkeala, H. J., & Vandamme, P. (2002). Taxonomic study of *Weissella confusa* and description of *Weissella cibaria* sp. nov., detected in food and clinical samples. *International Journal of Systematic and Evolutionary Microbiology*, 52, 141–148.
- Blasco, L., Ferrer, S., & Pardo, I. (2003). Development of specific fluorescent oligonucleotide probes for *in situ* identification of wine lactic acid bacteria. *FEMS Microbiology Letters*, 225, 115–123.
- Bokulich, N. A., & Bamforth, C. W. (2013). The microbiology of malting and brewing. *Microbiology and Molecular Biology Reviews*, 77, 157–172.



- Bokulich, N. A., Bamforth, C. W., & Mills, D. A. (2012a). Brewhouse-resident microbiota are responsible for multi-stage fermentation of American coolship ale. *PLoS ONE*, *7*, e35507.
- Bokulich, N. A., Joseph, C. M., Allen, G., Benson, A. K., & Mills, D. A. (2012b). Next-generation sequencing reveals significant bacterial diversity of botrytized wine. *PLoS ONE*, *7*, e36357.
- Bokulich, N. A., & Mills, D. A. (2012c). Differentiation of mixed lactic acid bacteria communities in beverage fermentations using targeted terminal restriction fragment length polymorphism. *Food Microbiology*, *31*, 126–132.
- Bokulich, N. A., & Mills, D. A. (2012d). Next-generation approaches to the microbial ecology of food fermentations. *BMB Reports*, *45*, 377–389.
- Bolotin, A., Wincker, P., Mauger, S., Jaillon, O., Malarne, K., Weissenbach, J., Ehrlich, S. D., & Sorokin, A. (2001). The complete genome sequence of the lactic acid bacterium *Lactococcus lactis* ssp. *lactis* IL1403. *Genome Research*, *11*, 731–753.
- Bottari, B., Santarelli, M., Neviani, E., & Gatti, M. (2010). Natural whey starter for Parmigiano Reggiano: culture-independent approach. *Journal of Applied Microbiology*, *108*, 1676–1684.
- Breidt, F., Medina, E., Wafa, D., Pérez-Díaz, I., Franco, W., Huang, H.-Y., Johanningsmeier, S. D., & Kim, J. H. (2013). Characterization of cucumber fermentation spoilage bacteria by enrichment culture and 16S rDNA cloning. *Journal of Food Science*, *78*, M470–M476.
- Caporaso, J. G., Kuczynski, J., Stombaugh, J., Bittinger, K., Bushman, F. D., Costello, E. K., Fierer, N., Pena, A. G., Goodrich, J. K., Gordon, J. I., Huttley, G. A., Kelley, S. T., Knights, D., Koenig, J. E., Ley, R. E., Lozupone, C. A., McDonald, D., Muegge, B. D., Pirrung, M., Reeder, J., Sevinsky, J. R., Turnbaugh, P. J., Walters, W. A., Widmann, J., Yatsunenko, T., Zaneveld, J., & Knight, R. (2010). QIIME allows analysis of high-throughput community sequencing data. *Nature Methods*, *7*, 335–336.
- Carver, T. J., Rutherford, K. M., Berriman, M., Rajandream, M. A., Barrell, B. G., & Parkhill, J. (2005). ACT: the artemis comparison tool. *Bioinformatics (Oxford, England)*, *21*, 3422–3423.
- Case, R. J., Boucher, Y., Dahllöf, I., Holmström, C., Doolittle, W. F., & Kjelleberg, S. (2007). Use of 16S rRNA and *rpoB* genes as molecular markers for microbial ecology studies. *Applied and Environmental Microbiology*, *73*, 278–288.
- Chambel, L., Chelo, I. M., Ze-Ze, L., Pedro, L. G., Santos, M. A., & Tenreiro, R. (2006). *Leuconostoc pseudoficulneum* sp. nov., isolated from a ripe fig. *International Journal of Systematic and Evolutionary Microbiology*, *56*, 1375–1381.
- Chen, T., Yu, W.-H., Izard, J., Baranova, O. V., Lakshmanan, A., & Dewhirst, F. E. (2010). The human oral microbiome database: a web accessible resource for investigating oral microbe taxonomic and genomic information. *Database*, (p. baq013).
- Cho, J. C., & Tiedje, J. M. (2001). Bacterial species determination from DNA–DNA hybridization by using genome fragments and DNA microarrays. *Applied and Environmental Microbiology*, *67*, 3677–3682.

- Cho, R. J., Mindrinos, M., Richards, D. R., Sapolsky, R. J., Anderson, M., Drenkard, E., Dewdney, J., Reuber, T. L., Stammers, M., Federspiel, N., Theologis, A., Yang, W.-H., Hubbell, E., Au, M., Chung, E. Y., Lashkari, D., Lemieux, B., Dean, C., Lipshutz, R. J., Ausubel, F. M., Davis, R. W., & Oefner, P. J. (1999). Genome-wide mapping with biallelic markers in *Arabidopsis thaliana*. *Nature Genetics*, 23, 203–207.
- Claesson, M. J., van Sinderen, D., & O'Toole, P. W. (2008). *Lactobacillus* phylogenomics – towards a reclassification of the genus. *International Journal of Systematic and Evolutionary Microbiology*, 58, 2945–2954.
- Clark, C. G., Kruczkiewicz, P., Guan, C., McCorrister, S. J., Chong, P., Wylie, J., van Caesele, P., Tabor, H. A., Snarr, P., Gilmour, M. W., Taboada, E. N., & Westmacott, G. R. (2013). Evaluation of MALDI-TOF mass spectroscopy methods for determination of *Escherichia coli* pathotypes. *Journal of Microbiological Methods*, 94, 180–191.
- Cleenwerck, I., De Vos, P., & De Vuyst, L. (2010). Phylogeny and differentiation of species of the genus *Gluconacetobacter* and related taxa based on multilocus sequence analyses of housekeeping genes and reclassification of *Acetobacter xylinus* subsp. *sucrofermentans* as *Gluconacetobacter sucrofermentans*. *International Journal of Systematic and Evolutionary Microbiology*, 60, 2277–2283.
- Cocolin, L., Alessandria, V., Dolci, P., Gorra, R., & Rantsiou, K. (2013). Culture independent methods to assess the diversity and dynamics of microbiota during food fermentation. *International Journal of Food Microbiology*, 167, 29 – 43.
- Coenye, T., Falsen, E., Vancanneyt, M., Hoste, B., Govan, J. R. W., Kersters, K., & Vandamme, P. (1999). Classification of *Alcaligenes faecalis*-like isolates from the environment and human clinical samples as *Ralstonia gilardii* sp. nov. *International Journal of Systematic Bacteriology*, 49, 405–413.
- Coenye, T., Gevers, D., Van de Peer, Y., Vandamme, P., & Swings, J. (2005). Towards a prokaryotic genomic taxonomy. *FEMS Microbiology Reviews*, 29, 147–167.
- Coenye, T., & Vandamme, P. (2003). Extracting phylogenetic information from whole-genome sequencing projects: the lactic acid bacteria as a test case. *Microbiology*, 149, 3507–3517.
- Collins, M. D., Samelis, J., Metaxopoulos, J., & Wallbanks, S. (1993). Taxonomic studies on some *Leuconostoc*-like organisms from fermented sausages: description of a new genus *Weissella* for the *Leuconostoc paramesenteroides* group of species. *Journal of Applied Microbiology*, 75, 595–603.
- Coorevits, A., De Jonghe, V., Vandroemme, J., Reekmans, R., Heyrman, J., Messens, W., De Vos, P., & Heyndrickx, M. (2008). Comparative analysis of the diversity of aerobic spore-forming bacteria in raw milk from organic and conventional dairy farms. *Systematic and Applied Microbiology*, 31, 126–140.
- de Crecy-Lagard, V., El Yacoubi, B., de la Garza, R. D., Noiriél, A., & Hanson, A. D. (2007). Comparative genomics of bacterial and plant folate synthesis and salvage: predictions and validations. *BMC Genomics*, 8, 245.

- Crous, P. W., Gams, W., Stalpers, J. A., Robert, V., & Stegehuis, G. (2004). MycoBank: an online initiative to launch mycology into the 21st century. *Studies in Mycology*, 50, 19–22.
- Darling, A. C. E., Mau, B., Blattner, F. R., & Perna, N. T. (2004). MAUVE: multiple alignment of conserved genomic sequence with rearrangements. *Genome Research*, 14, 1394–1403.
- Darmon, E., & Leach, D. R. F. (2014). Bacterial genome instability. *Microbiology and Molecular Biology Reviews*, 78, 1–39.
- De Bruyne, K., Camu, N., De Vuyst, L., & Vandamme, P. (2010). *Weissella fabaria* sp. nov., from a Ghanaian cocoa fermentation. *International Journal of Systematic and Evolutionary Microbiology*, 60, 1999–2005.
- De Bruyne, K., Camu, N., Lefebvre, K., De Vuyst, L., & Vandamme, P. (2008a). *Weissella ghanensis* sp. nov., isolated from a Ghanaian cocoa fermentation. *International Journal of Systematic and Evolutionary Microbiology*, 58, 2721–2725.
- De Bruyne, K., Franz, C. M., Vancanneyt, M., Schillinger, U., Mozzi, F., de Valdez, G. F., De Vuyst, L., & Vandamme, P. (2008b). *Pediococcus argentinus* sp. nov. from Argentinean fermented wheat flour and identification of *Pediococcus* species by *pheS*, *rpoA* and *atpA* sequence analysis. *International Journal of Systematic and Evolutionary Microbiology*, 58, 2909–2916.
- De Bruyne, K., Schillinger, U., Caroline, L., Boehringer, B., Cleenwerck, I., Vancanneyt, M., De Vuyst, L., Franz, C. M. A. P., & Vandamme, P. (2007). *Leuconostoc holzapfelii* sp. nov., isolated from Ethiopian coffee fermentation and assessment of sequence analysis of housekeeping genes for delineation of *Leuconostoc* species. *International Journal of Systematic and Evolutionary Microbiology*, 57, 2952–2959.
- De Bruyne, K., Slabbinck, B., Waegeman, W., Vauterin, P., De Baets, B., & Vandamme, P. (2011). Bacterial species identification from MALDI-TOF mass spectra through data analysis and machine learning. *Systematic and Applied Microbiology*, 34, 20–29.
- De Clerck, E., & De Vos, P. (2002). Study of the bacterial load in a gelatine production process focussed on *Bacillus* and related endosporeforming genera. *Systematic and Applied Microbiology*, 25, 611–617.
- De Vuyst, L., Lefeber, T., Papalexandratou, Z., & Camu, N. (2010). The functional role of lactic acid bacteria in cocoa bean fermentation. In *Biotechnology of lactic acid bacteria* (pp. 301–325). Wiley-Blackwell.
- Delcher, A. L., Bratke, K. A., Powers, E. C., & Salzberg, S. L. (2007). Identifying bacterial genes and endosymbiont DNA with Glimmer. *Bioinformatics*, 23, 673–679.
- Delcher, A. L., Harmon, D., Kasif, S., White, O., & Salzberg, S. L. (1999). Improved microbial gene identification with GLIMMER. *Nucleic Acids Research*, 27, 4636–4641.
- Delmont, T. O., Malandain, C., Prestat, E., Larose, C., Monier, J.-M., Simonet, P., & Vogel, T. M. (2011). Metagenomic mining for microbiologists. *ISME Journal*, 5, 1837–1843.
- Deloger, M., El Karoui, M., & Petit, M. A. (2009). A genomic distance based on MUM indicates discontinuity between most bacterial species and genera. *Journal of Bacteriology*, 191, 91–99.

- DeLong, E. F., & Karl, D. M. (2005). Genomic perspectives in microbial oceanography. *Nature*, 437, 336–342.
- Derrien, M., Vaughan, E. E., Plugge, C. M., & de Vos, W. M. (2004). *Akkermansia muciniphila* gen. nov., sp. nov., a human intestinal mucin-degrading bacterium. *International Journal of Systematic and Evolutionary Microbiology*, 54, 1469–1476.
- Di Cagno, R., Coda, R., De Angelis, M., & Gobbetti, M. (2013). Exploitation of vegetables and fruits through lactic acid fermentation. *Food Microbiology*, 33, 1–10.
- Di Mattia, E., Grego, S., & Cacciari, I. (2002). Eco-physiological characterization of soil bacterial populations in different states of growth. *Microbial Ecology*, 43, 34–43.
- Dickie, I. A. (2010). Insidious effects of sequencing errors on perceived diversity in molecular surveys. *New Phytologist*, 188, 916–918.
- Dieckmann, R., Graeber, I., Kaesler, I., Szewzyk, U., & von Dohren, H. (2005). Rapid screening and dereplication of bacterial isolates from marine sponges of the sula ridge by intact-cell-MALDI-TOF mass spectrometry (ICM-MS). *Applied Microbiology and Biotechnology*, 67, 539–548.
- Dijkshoorn, L., Ursing, B. M., & Ursing, J. B. (2000). Strain, clone and species: comments on three basic concepts of bacteriology. *Journal of Medical Microbiology*, 49, 397–401.
- Doolittle, W. F., & Papke, R. T. (2006). Genomics and the bacterial species problem. *Genome Biology*, 7, 116.
- Duskova, M., Sedo, O., Ksicova, K., Zdrahal, Z., & Karpiskova, R. (2012). Identification of lactobacilli isolated from food by genotypic methods and MALDI-TOF MS. *International Journal of Food Microbiology*, 159, 107–114.
- Dykhuizen, D. (2005). Species numbers in bacteria. *Proceedings of the California Academy of Sciences*, 56, 62–71.
- Eddy, S. R. (1998). Profile hidden Markov models. *Bioinformatics*, 14, 755–763.
- Edgar, R. C., Haas, B. J., Clemente, J. C., Quince, C., & Knight, R. (2011). UCHIME improves sensitivity and speed of chimera detection. *Bioinformatics*, 27, 2194–2200.
- Ehrmann, M. A., Freiding, S., & Vogel, R. F. (2009). *Leuconostoc palmae* sp. nov., a novel lactic acid bacterium isolated from palm wine. *International Journal of Systematic and Evolutionary Microbiology*, 59, 943–947.
- Eisen, J. A. (2000). Assessing evolutionary relationships among microbes from whole-genome analysis. *Current Opinion in Microbiology*, 3, 475–480.
- Endo, A., Irisawa, T., Futagawa-Endo, Y., Sonomoto, K., Itoh, K., Takano, K., Okada, S., & Dicks, L. M. T. (2010). *Fructobacillus tropaeoli* sp. nov., a novel fructophilic lactic acid bacterium isolated from a flower. *International Journal of Systematic and Evolutionary Microbiology*, (pp. ijs.0.023838–0).

- Endo, A., & Okada, S. (2008). Reclassification of the genus *Leuconostoc* and proposals of *Fructobacillus fructosus* gen. nov., comb. nov., *Fructobacillus durionis* comb. nov., *Fructobacillus ficulneus* comb. nov. and *Fructobacillus pseudoficulneus* comb. nov. *International Journal of Systematic and Evolutionary Microbiology*, 58, 2195–2205.
- Engelbrektson, A., Kunin, V., Wrighton, K. C., Zvenigorodsky, N., Chen, F., Ochman, H., & Hugenholtz, P. (2010). Experimental factors affecting PCR-based estimates of microbial species richness and evenness. *ISME Journal*, 4, 642–647.
- Ercolini, D. (2013). High-throughput sequencing and metagenomics: Moving forward in the culture-independent analysis of food microbial ecology. *Applied and Environmental Microbiology*, 79, 3148–3155.
- Ezaki, T., Hashimoto, Y., & Yabuuchi, E. (1989). Fluorometric deoxyribonucleic acid–deoxyribonucleic acid hybridization in microdilution wells as an alternative to membrane filter hybridization in which radioisotopes are used to determine genetic relatedness among bacterial strains. *International Journal of Systematic Bacteriology*, 39, 224–229.
- Fernandez, M., del Rio, B., Linares, D. M., Martin, M. C., & Alvarez, M. A. (2006). Real-time polymerase chain reaction for quantitative detection of histamine-producing bacteria: use in cheese production. *Journal of Dairy Science*, 89, 3763–3769.
- Field, D., Amaral-Zettler, L., Cochrane, G., Cole, J. R., Dawyndt, P., Garrity, G. M., Gilbert, J., Glockner, F. O., Hirschman, L., Karsch-Mizrachi, I., Klenk, H. P., Knight, R., Kottmann, R., Kyrpides, N., Meyer, F., San Gil, I., Sansone, S. A., Schriml, L. M., Sterk, P., Tatusova, T., Ussery, D. W., White, O., & Wooley, J. (2011). The genomic standards consortium. *PLoS Biology*, 9, e1001088.
- Field, D., Garrity, G., Gray, T., Morrison, N., Selengut, J., Sterk, P., Tatusova, T., Thomson, N., Allen, M. J., Angiuoli, S. V., Ashburner, M., Axelrod, N., Baldauf, S., Ballard, S., Boore, J., Cochrane, G., Cole, J., Dawyndt, P., De Vos, P., DePamphilis, C., Edwards, R., Faruque, N., Feldman, R., Gilbert, J., Gilna, P., Glockner, F. O., Goldstein, P., Guralnick, R., Haft, D., Hancock, D., Hermjakob, H., Hertz-Fowler, C., Hugenholtz, P., Joint, I., Kagan, L., Kane, M., Kennedy, J., Kowalchuk, G., Kottmann, R., Kolker, E., Kravitz, S., Kyrpides, N., Leebens-Mack, J., Lewis, S. E., Li, K., Lister, A. L., Lord, P., Maltsev, N., Markowitz, V., Martiny, J., Methe, B., Mizrachi, I., Moxon, R., Nelson, K., Parkhill, J., Proctor, L., White, O., Sansone, S. A., Spiers, A., Stevens, R., Swift, P., Taylor, C., Tatenos, Y., Tett, A., Turner, S., Ussery, D., Vaughan, B., Ward, N., Whetzel, T., San Gil, I., Wilson, G., & Wipat, A. (2008). The minimum information about a genome sequence (MIGS) specification. *Nature Biotechnology*, 26, 541–547.
- Fleet, G. H. (1999). Microorganisms in food ecosystems. *International Journal of Food Microbiology*, 50, 101–117.
- Fleet, G. H. (2007). Yeasts in foods and beverages: impact on product quality and safety. *Current Opinion in Biotechnology*, 18, 170–175.
- Fleischmann, R. D., Adams, M. D., White, O., Clayton, R. A., Kirkness, E. F., Kerlavage, A. R., Bult, C. J., Tomb, J. F., Dougherty, B. A., Merrick, J. M., & al., E. (1995). Whole-genome random sequencing and assembly of *Haemophilus influenzae* Rd. *Science*, 269, 496–512.

- Franzmann, P. D., Höpfl, P., Weiss, N., & Tindall, B. J. (1991). Psychrotrophic, lactic acid-producing bacteria from anoxic waters in Ace Lake, Antarctica; *Carnobacterium funditum* sp. nov. and *Carnobacterium alterfunditum* sp. nov. *Archives of Microbiology*, 156, 255–262.
- Fraser, C., Hanage, W. P., & Spratt, B. G. (2007). Recombination and the nature of bacterial speciation. *Science*, 315, 476–480.
- Freiwald, A., & Sauer, S. (2009). Phylogenetic classification and identification of bacteria by mass spectrometry. *Nature Protocols*, 4, 732–742.
- Garcia-Armisen, T., Papalexandratou, Z., Hendryckx, H., Camu, N., Vrancken, G., Vuyst, L., & Cornelis, P. (2010). Diversity of the total bacterial community associated with Ghanaian and Brazilian cocoa bean fermentation samples as revealed by a 16 S rRNA gene clone library. *Applied Microbiology and Biotechnology*, 87, 2281–2292.
- García-Gimeno, R. M., & Zurera-Cosano, G. (1997). Determination of ready-to-eat vegetable salad shelf-life. *International Journal of Food Microbiology*, 36, 31–38.
- Gardes, M., & Bruns, T. D. (1993). ITS primers with enhanced specificity for *basidiomycetes* – application to the identification of mycorrhizae and rusts. *Molecular Ecology*, 2, 113–118.
- Garrity, G. M., Bell, J. A., & Lilburn, T. G. (2004). *Taxonomic outline of the prokaryotes. Bergey's manual of systematic bacteriology*. New York: Springer.
- Gevers, D., Cohan, F. M., Lawrence, J. G., Spratt, B. G., Coenye, T., Feil, E. J., Stackebrandt, E., Van de Peer, Y., Vandamme, P., Thompson, F. L., & Swings, J. (2005). Opinion: Re-evaluating prokaryotic species. *Nature Reviews Microbiology*, 3, 733–739.
- Gevers, D., Huys, G., & Swings, J. (2001). Applicability of rep-PCR fingerprinting for identification of *Lactobacillus* species. *FEMS Microbiology Letters*, 205, 31–36.
- Ghyselinck, J., Van Hoorde, K., Hoste, B., Heylen, K., & De Vos, P. (2011). Evaluation of MALDI-TOF MS as a tool for high-throughput dereplication. *Journal of Microbiological Methods*, 86, 327–336.
- Gkatzionis, K., Yunita, D., Linforth, R. S. T., Dickinson, M., & Dodd, C. E. R. (2014). Diversity and activities of yeasts from different parts of a Stilton cheese. *International Journal of Food Microbiology*, 177, 109–116.
- Goll, J., Rusch, D. B., Tanenbaum, D. M., Thiagarajan, M., Li, K., Methé, B. A., & Yooseph, S. (2010). METAREP: JCVI metagenomics reports—an open source tool for high-performance comparative metagenomics. *Bioinformatics*, 26, 2631–2632.
- Goris, J., Konstantinidis, K., Klappenbach, J. A., Coenye, T., Vandamme, P., & Tiedje, J. M. (2007). DNA-DNA hybridization values and their relationship to whole-genome sequence similarities. *International Journal of Systematic and Evolutionary Microbiology*, 57, 81–91.
- Goris, J., Suzuki, K.-i., Vos, P. D., Nakase, T., & Kersters, K. (1998). Evaluation of a microplate DNA – DNA hybridization method compared with the initial renaturation method. *Canadian Journal of Microbiology*, 44, 1148–1153.

- Gouy, M., Guindon, S. A., & Gascuel, O. (2010). SeaView Version 4: A Multiplatform Graphical User Interface for Sequence Alignment and Phylogenetic Tree Building. *Molecular Biology and Evolution*, 27, 221–224.
- Haas, B. J., Gevers, D., Earl, A. M., Feldgarden, M., Ward, D. V., Giannoukos, G., Ciulla, D., Tabbaa, D., Highlander, S. K., Sodergren, E., Methé, B., DeSantis, T. Z., The Human Microbiome, C., Petrosino, J. F., Knight, R., & Birren, B. W. (2011). Chimeric 16S rRNA sequence formation and detection in Sanger and 454-pyrosequenced PCR amplicons. *Genome Research*, 21, 494–504.
- Hall, B. (2011). *Phylogenetic trees made easy: A how-to manual*. (4th ed.). Sinauer Associates, Inc. Publishers.
- Hardenbol, P., Baner, J., Jain, M., Nilsson, M., Namsaraev, E. A., Karlin-Neumann, G. A., Fakhrai-Rad, H., Ronaghi, M., Willis, T. D., Landegren, U., & Davis, R. W. (2003). Multiplexed genotyping with sequence-tagged molecular inversion probes. *Nature Biotechnology*, 21, 673–678.
- Hartmann, M., Howes, C. G., Abarenkov, K., Mohn, W. W., & Nilsson, R. H. (2010). V-Xtractor: an open-source, high-throughput software tool to identify and extract hypervariable regions of small subunit (16S/18S) ribosomal RNA gene sequences. *Journal of Microbiological Methods*, 83, 250–253.
- Hayashi, N., Ito, M., Horiike, S., & Taguchi, H. (2001). Molecular cloning of a putative divalent-cation transporter gene as a new genetic marker for the identification of *Lactobacillus brevis* strains capable of growing in beer. *Applied Microbiology and Biotechnology*, 55, 596–603.
- van Hijum, S. A., Vaughan, E. E., & Vogel, R. F. (2013). Application of state-of-art sequencing technologies to indigenous food fermentations. *Current Opinion in Biotechnology*, 24, 178 – 186.
- Holdeman, L. V., & Moore, W. E. C. (1977). *Anaerobic laboratory manual*. (4th ed.). Blacksburg, Virginia, USA: Virginia Polytechnic Institute and State University, Anaerobe Laboratory.
- Holzapfel, W., Franz, C. A. P., Ludwig, W., Back, W., & Dicks, L. T. (2006). The genera *Pediococcus* and *Tetragenococcus*. In M. Dworkin, S. Falkow, E. Rosenberg, K.-H. Schleifer, & E. Stackebrandt (Eds.), *The Prokaryotes* chapter 8. (pp. 229–266). Springer US.
- Huber, T., Faulkner, G., & Hugenholtz, P. (2004). Bellerophon: a program to detect chimeric sequences in multiple sequence alignments. *Bioinformatics*, 20, 2317–2319.
- Humblot, C., & Guyot, J. P. (2009). Pyrosequencing of tagged 16S rRNA gene amplicons for rapid deciphering of the microbiomes of fermented foods such as pearl millet slurries. *Applied and Environmental Microbiology*, 75, 4354–4361.
- Huse, S., Huber, J., Morrison, H., Sogin, M., & Welch, D. (2007). Accuracy and quality of massively parallel DNA pyrosequencing. *Genome Biology*, 8, R143.
- Huse, S., Welch, D. M., Morrison, H. G., & Sogin, M. L. (2010). Ironing out the wrinkles in the rare biosphere through improved OTU clustering. *Environmental Microbiology*, 12, 1889–1898.
- Huson, D., & Mitra, S. (2012). Introduction to the analysis of environmental sequences: metagenomics with MEGAN. In M. Anisimova (Ed.), *Evolutionary genomics* chapter 17. (pp. 415–429). Humana Press volume 856.

- Ibarburu, I., Aznar, R., Elizaquivel, P., Garcia-Quintans, N., Lopez, P., Munduate, A., Irastorza, A., & Duenas, M. T. (2010). A real-time PCR assay for detection and quantification of 2-branched (1,3)-beta-D-glucan producing lactic acid bacteria in cider. *International Journal of Food Microbiology*, 143, 26–31.
- Iijima, K., Suzuki, K., Asano, S., Kuriyama, H., & Kitagawa, Y. (2007). Isolation and identification of potential beer-spoilage *Pediococcus inopinatus* and beer-spoilage *Lactobacillus backi* strains carrying the *horA* and *horC* gene clusters. *Journal of the Institute of Brewing*, 113, 96–101.
- Iijima, K., Suzuki, K., Ozaki, K., & Yamashita, H. (2006). HorC confers beer-spoilage ability on hop-sensitive *Lactobacillus brevis* ABBC45cc. *Journal of Applied Microbiology*, 100, 1282–1288.
- Illegghems, K., De Vuyst, L., Papalexandratou, Z., & Weckx, S. (2012). Phylogenetic analysis of a spontaneous cocoa bean fermentation metagenome reveals new insights into its bacterial and fungal community diversity. *PLoS ONE*, 7, e38040.
- Illegghems, K., De Vuyst, L., & Weckx, S. (2013). Complete genome sequence and comparative analysis of *Acetobacter pasteurianus* 386B, a strain well-adapted to the cocoa bean fermentation ecosystem. *BMC Genomics*, 14, 526.
- Irlinger, F. (2008). Safety assessment of dairy microorganisms: coagulase-negative staphylococci. *International Journal of Food Microbiology*, 126, 302–310.
- Janssens, M., Myter, N., De Vuyst, L., & Leroy, F. (2012). Species diversity and metabolic impact of the microbiota are low in spontaneously acidified Belgian sausages with an added starter culture of *Staphylococcus carnosus*. *Food Microbiology*, 29, 167–177.
- Jung, J. Y., Lee, S. H., Kim, J. M., Park, M. S., Bae, J.-W., Hahn, Y., Madsen, E. L., & Jeon, C. O. (2011). Metagenomic analysis of kimchi, a traditional Korean fermented food. *Applied and Environmental Microbiology*, 77, 2264–2274.
- Jung, M. J., Nam, Y. D., Roh, S. W., & Bae, J. W. (2012). Unexpected convergence of fungal and bacterial communities during fermentation of traditional Korean alcoholic beverages inoculated with various natural starters. *Food Microbiology*, 30, 112–123.
- Kalisky, T., & Quake, S. R. (2011). Single-cell genomics. *Nature Methods*, 8, 311–314.
- Kim, J., Chun, J., & Han, H. U. (2000). *Leuconostoc kimchii* sp. nov., a new species from kimchi. *International Journal of Systematic and Evolutionary Microbiology*, 50, 1915–1919.
- Kim, M., & Chun, J. (2005). Bacterial community structure in kimchi, a Korean fermented vegetable food, as revealed by 16S rRNA gene analysis. *International Journal of Food Microbiology*, 103, 91–96.
- Kim, M.-S., Roh, S. W., Nam, Y.-D., Yoon, J.-H., & Bae, J.-W. (2009). *Carnobacterium jeotgali* sp. nov., isolated from a Korean traditional fermented food. *International Journal of Systematic and Evolutionary Microbiology*, 59, 3168–3171.
- Kim, Y. S., Kim, M. C., Kwon, S. W., Kim, S. J., Park, I. C., Ka, J. O., & Weon, H. Y. (2011). Analyses of bacterial communities in meju, a Korean traditional fermented soybean bricks, by cultivation-based and pyrosequencing methods. *The Journal of Microbiology*, 49, 340–348.



- Kimura, M. (1980). A simple method for estimating evolutionary rates of base substitutions through comparative studies of nucleotide sequences. *Journal of Molecular Evolution*, 16, 111–120.
- Kiyohara, M., Koyanagi, T., Matsui, H., Yamamoto, K., Take, H., Katsuyama, Y., Tsuji, A., Miyamae, H., Kondo, T., Nakamura, S., Katayama, T., & Kumagai, H. (2012). Changes in microbiota population during fermentation of narezushi as revealed by pyrosequencing analysis. *Bioscience, Biotechnology, and Biochemistry*, 76, 48–52.
- Klappenbach, J. A., Dunbar, J. M., & Schmidt, T. M. (2000). rRNA operon copy number reflects ecological strategies of bacteria. *Applied and Environmental Microbiology*, 66, 1328–1333.
- Klenk, H. P., & Goker, M. (2010). En route to a genome-based classification of Archaea and Bacteria? *Systematic and Applied Microbiology*, 33, 175–182.
- Knight, R., Jansson, J., Field, D., Fierer, N., Desai, N., Fuhrman, J., Hugenholtz, P., van der Lelie, D., Meyer, F., Stevens, R., Bailey, M., Gordon, J., Kowalchuk, G., & Gilbert, J. (2012). Unlocking the potential of metagenomics through replicated experimental design. *Nature biotechnology*, 30, 513–520. doi:10.1038/nbt.2235.
- Kodama, Y., Shumway, M., Leinonen, R., & International Nucleotide Sequence Database, C. (2012). The sequence read archive: explosive growth of sequencing data. *Nucleic Acids Research*, 40, D54–6.
- Konstantinidis, K., Ramette, A., & Tiedje, J. M. (2006). The bacterial species definition in the genomic era. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 361, 1929–1940.
- Konstantinidis, K., & Stackebrandt, E. (2013). Defining taxonomic ranks. In E. Rosenberg, E. DeLong, S. Lory, E. Stackebrandt, & F. Thompson (Eds.), *The prokaryotes* chapter 4. (pp. 229–254). Springer Berlin Heidelberg.
- Konstantinidis, K., & Tiedje, J. M. (2005). Genomic insights that advance the species definition for prokaryotes. *Proceedings of the National Academy of Sciences of the United States of America*, 102, 2567–2572.
- Kottmann, R., Gray, T., Murphy, S., Kagan, L., Kravitz, S., Lombardot, T., Field, D., Glockner, F. O., & Genomic Standards, C. (2008). A standard MGS/MIMS compliant XML schema: toward the development of the genomic contextual data markup language (GCDML). *OMICS*, 12, 115–121.
- Koyanagi, T., Kiyohara, M., Matsui, H., Yamamoto, K., Kondo, T., Katayama, T., & Kumagai, H. (2011). Pyrosequencing survey of the microbial diversity of 'narezushi', an archetype of modern Japanese sushi. *Letters in Applied Microbiology*, 53, 635–640.
- Kozich, J. J., Westcott, S. L., Baxter, N. T., Highlander, S. K., & Schloss, P. D. (2013). Development of a dual-index sequencing strategy and curation pipeline for analyzing amplicon sequence data on the MiSeq Illumina sequencing platform. *Applied and Environmental Microbiology*, .
- Kumara, H. M. C. S., De Cort, S., & Verachtert, H. (1993). Localization and characterization of  $\alpha$ -glucosidase activity in *Brettanomyces lambicus*. *Applied and Environmental Microbiology*, 59, 2352–2358.
- Kumara, H. M. C. S., & Verachtert, H. (1991). Identification of lambic superattenuating micro-organisms by the use of selective antibiotics. *Journal of the Institute of Brewing*, 97, 181–185.

- Kunin, V., Copeland, A., Lapidus, A., Mavromatis, K., & Hugenholtz, P. (2008). A bioinformatician's guide to metagenomics. *Microbiology and Molecular Biology Reviews*, 72, 557–578.
- Kunin, V., Engelbrektson, A., Ochman, H., & Hugenholtz, P. (2010). Wrinkles in the rare biosphere: pyrosequencing errors can lead to artificial inflation of diversity estimates. *Environmental Microbiology*, 12, 118–123.
- Kurtzman, C. P., & Robnett, C. J. (1998). Identification and phylogeny of ascomycetous yeasts from analysis of nuclear large subunit (26S) ribosomal DNA partial sequences. *Antonie van Leeuwenhoek*, 73, 331–371.
- Lacerda, I. C. A., Gomes, F. C. O., Borelli, B. M., Faria Jr, C. L. L., Franco, G. R., Mourão, M. M., Morais, P. B., & Rosa, C. A. (2011). Identification of the bacterial community responsible for traditional fermentation during sour cassava starch, cachaça and minas cheese production using culture-independent 16s rRNA gene sequence analysis. *Brazilian Journal of Microbiology*, 42, 650–657.
- Ladero, V., Coton, M., Fernandez, M., Buron, N., Martin, M. C., Guichard, H., Coton, E., & Alvarez, M. A. (2011). Biogenic amines content in Spanish and French natural ciders: application of qPCR for quantitative detection of biogenic amine-producers. *Food Microbiology*, 28, 554–561.
- Lagier, J. C., Armougom, F., Million, M., Hugon, P., Pagnier, I., Robert, C., Bittar, F., Fournous, G., Gimenez, G., Maraninchi, M., Trape, J. F., Koonin, E. V., La Scola, B., & Raoult, D. (2012). Microbial culturomics: paradigm shift in the human gut microbiome study. *Clinical Microbiology and Infection*, 18, 1185–1193.
- Lapage, S., Sneath, P., Lessel, E., Skerman, V., Seeliger, H., & Clark, W. (1992). *International code of nomenclature of bacteria: Bacteriological code (1990 revision)*. *Bacterial Code*. Washington DC: American Society for Microbiology.
- Laureys, D., & De Vuyst, L. (2014). Microbial species diversity, community dynamics, and metabolite kinetics of water kefir fermentation. *Applied and Environmental Microbiology*, 80, 2564–2572.
- Lee, J. S., Lee, K. C., Ahn, J.-S., Mheen, T.-I., Pyun, Y.-R., & Park, Y.-H. (2002). *Weissella koreensis* sp. nov., isolated from kimchi. *International Journal of Systematic and Evolutionary Microbiology*, 52, 1257–1261.
- Lee, S. H., Park, M. S., Jung, J. Y., & Jeon, C. O. (2012). *Leuconostoc miyukkimchii* sp. nov., isolated from brown algae (*Undaria pinnatifida*) kimchi. *International Journal of Systematic and Evolutionary Microbiology*, 62, 1098–1103.
- Lefébure, T., & Stanhope, M. (2007). Evolution of the core and pan-genome of *Streptococcus*: positive selection, recombination, and genome composition. *Genome Biology*, 8, 1–17.
- Leinonen, R., Akhtar, R., Birney, E., Bower, L., Cerdano-Tarraga, A., Cheng, Y., Cleland, I., Faruque, N., Goodgame, N., Gibson, R., Hoad, G., Jang, M., Pakseresht, N., Plaister, S., Radhakrishnan, R., Reddy, K., Sobhany, S., Ten Hoopen, P., Vaughan, R., Zalunin, V., & Cochrane, G. (2011). The European nucleotide archive. *Nucleic Acids Research*, 39, D28–31.
- Leroy, F., & De Vuyst, L. (2004). Lactic acid bacteria as functional starter cultures for the food fermentation industry. *Trends in Food Science and Technology*, 15, 67–78.

- Leroy, F., Verluuyten, J., & De Vuyst, L. (2006). Functional meat starter cultures for improved sausage fermentation. *International Journal of Food Microbiology*, 106, 270–285.
- Li, X. R., Ma, E. B., Yan, L. Z., Meng, H., Du, X. W., Zhang, S. W., & Quan, Z. X. (2011). Bacterial and fungal diversity in the traditional Chinese liquor fermentation process. *International Journal of Food Microbiology*, 146, 31–37.
- Livermore, J. A., Emrich, S. J., Tan, J., & Jones, S. E. (2014). Freshwater bacterial lifestyles inferred from comparative genomics. *Environmental Microbiology*, 16, 746–758.
- Lodolo, E. J., Kock, J. L. F., Axcell, B. C., & Brooks, M. (2008). The yeast *Saccharomyces cerevisiae*—the main character in beer brewing. *FEMS Yeast Research*, 8, 1018–1036.
- Loman, N. J., Constantinidou, C., Chan, J. Z. M., Halachev, M., Sergeant, M., Penn, C. W., Robinson, E. R., & Pallen, M. J. (2012). High-throughput bacterial genome sequencing: an embarrassment of choice, a world of opportunity. *Nature Reviews Microbiology*, 10, 599–606.
- Lozupone, C. A., Hamady, M., Kelley, S. T., & Knight, R. (2007). Quantitative and qualitative  $\beta$  diversity measures lead to different insights into factors that structure microbial communities. *Applied and Environmental Microbiology*, 73, 1576–1585.
- Lucas, P. M., Claisse, O., & Lonvaud-Funel, A. (2008). High frequency of histamine-producing bacteria in the enological environment and instability of the histidine decarboxylase production phenotype. *Applied and Environmental Microbiology*, 74, 811–817.
- Ludwig, W., Strunk, O., Westram, R., Richter, L., Meier, H., Yadhukumar, Buchner, A., Lai, T., Steppi, S., & Others, . (2004). ARB: a software environment for sequence data. *Nucleic Acids Research*, 32, 1363–1371.
- Lukashin, A. V., & Borodovsky, M. (1998). GeneMark.HMM: new solutions for gene finding. *Nucleic Acids Research*, 26, 1107–1115.
- Luo, C., Tsementzi, D., Kyrpides, N., Read, T., & Konstantinidis, K. T. (2012). Direct comparisons of Illumina vs. Roche 454 sequencing technologies on the same microbial community DNA sample. *PLoS ONE*, 7, e30087.
- Lyhs, U., Koort, J. M. K., Lundström, H. S., & Björkroth, K. J. (2004). *Leuconostoc gelidum* and *Leuconostoc gasicomitatum* strains dominated the lactic acid bacterium population associated with strong slime formation in an acetic-acid herring preserve. *International Journal of Food Microbiology*, 90, 207–218.
- Magnusson, J., Jonsson, H., Schnürer, J., & Roos, S. (2002). *Weissella soli* sp. nov., a lactic acid bacterium isolated from soil. *International Journal of Systematic and Evolutionary Microbiology*, 52, 831–834.
- Mahenthiralingam, E., Campbell, M. E., Henry, D. A., & Speert, D. P. (1996). Epidemiology of *Burkholderia cepacia* infection in patients with cystic fibrosis: analysis by randomly amplified polymorphic DNA fingerprinting. *Journal of Clinical Microbiology*, 34, 2914–2920.
- Maidak, B. L., Olsen, G. J., Larsen, N., Overbeek, R., McCaughey, M. J., & Woese, C. R. (1997). The RDP (ribosomal database project). *Nucleic Acids Research*, 25, 109–111.

- Makarova, K., Slesarev, A., Wolf, Y., Sorokin, A., Mirkin, B., Koonin, E., Pavlov, A., Pavlova, N., Karamychev, V., Polouchine, N., Shakhova, V., Grigoriev, I., Lou, Y., Rohksar, D., Lucas, S., Huang, K., Goodstein, D. M., Hawkins, T., Plengvidhya, V., Welker, D., Hughes, J., Goh, Y., Benson, A., Baldwin, K., Lee, J. H., Diaz-Muniz, I., Dosti, B., Smeianov, V., Wechter, W., Barabote, R., Lorca, G., Altermann, E., Barrangou, R., Ganesan, B., Xie, Y., Rawsthorne, H., Tamir, D., Parker, C., Breidt, F., Broadbent, J., Hutkins, R., O'Sullivan, D., Steele, J., Unlu, G., Saier, M., Klaenhammer, T., Richardson, P., Kozyavkin, S., Weimer, B., & Mills, D. (2006). Comparative genomics of the lactic acid bacteria. *Proceedings of the National Academy of Sciences of the United States of America*, 103, 15611–15616.
- Mamanova, L., Coffey, A. J., Scott, C. E., Kozarewa, I., Turner, E. H., Kumar, A., Howard, E., Shendure, J., & Turner, D. J. (2010). Target-enrichment strategies for next-generation sequencing. *Nature Methods*, 7, 111–118.
- Marcobal, A., Underwood, M. A., & Mills, D. A. (2008). Rapid determination of the bacterial composition of commercial probiotic products by terminal restriction fragment length polymorphism analysis. *Journal of Pediatric Gastroenterology and Nutrition*, 46, 608–611.
- Markowitz, V. M., Chen, I. M. A., Chu, K., Szeto, E., Palaniappan, K., Pillay, M., Ratner, A., Huang, J., Pagani, I., Tringe, S., Huntemann, M., Billis, K., Varghese, N., Tennesen, K., Mavromatis, K., Pati, A., Ivanova, N. N., & Kyrpides, N. C. (2014). IMG/M 4 version of the integrated metagenome comparative analysis system. *Nucleic Acids Research*, 42, D568–D573.
- Martens, H., Iserentant, D., & Verachtert, H. (1997). Microbiological aspects of a mixed yeast-bacterial fermentation in the production of a special Belgian acidic ale. *Journal of the Institute of Brewing*, 103, 85–91.
- Martín-Platero, A. M., Maqueda, M., Valdivia, E., Purswani, J., & Martínez-Bueno, M. (2009). Polyphasic study of microbial communities of two Spanish farmhouse goats' milk cheeses from Sierra de Aracena. *Food Microbiology*, 26, 294–304.
- Mattarelli, P., Holzapfel, W., Franz, C. M. A. P., Endo, A., Felis, G. E., Hammes, W., Pot, B., Dicks, L., & Dellaglio, F. (2014). Recommended minimal standards for description of new taxa of the genera *Bifidobacterium*, *Lactobacillus* and related genera. *International Journal of Systematic and Evolutionary Microbiology*, 64, 1434–1451.
- Mavromatis, K., Chu, K., Ivanova, N., Hooper, S. D., Markowitz, V. M., & Kyrpides, N. C. (2009). Gene context analysis in the integrated microbial genomes (IMG) data management system. *PLoS ONE*, 4, e7979.
- Mendes-Ferreira, A., Barbosa, C., Jimenez-Marti, E., Del Olmo, M. L., & Mendes-Faia, A. (2010). The wine yeast strain-dependent expression of genes implicated in sulfide production in response to nitrogen availability. *Journal of Microbiology and Biotechnology*, 20, 1314–1321.
- Mesbah, M., & Whitman, W. B. (1989). Measurement of deoxyguanosine/thymidine ratios in complex mixtures by high-performance liquid chromatography for determination of the mole percentage guanine + cytosine of DNA. *Journal of Chromatography*, 479, 297–306.
- Metzker, M. L. (2005). Emerging technologies in DNA sequencing. *Genome Research*, 15, 1767–1776.

- Meuzelaar, L. S., Lancaster, O., Pasche, J. P., Kopal, G., & Brookes, A. J. (2007). MegaPlex PCR: a strategy for multiplex amplification. *Nature Methods*, 4, 835–837.
- Meyer, F., Paarmann, D., D'Souza, M., Olson, R., Glass, E. M., Kubal, M., Paczian, T., Rodriguez, A., Stevens, R., Wilke, A., Wilkening, J., & Edwards, R. A. (2008). The metagenomics RAST server – a public resource for the automatic phylogenetic and functional analysis of metagenomes. *BMC Bioinformatics*, 9, 386.
- Midha, S., Ranjan, M., Sharma, V., Kumari, A., Singh, P. K., Korpole, S., & Patil, P. B. (2012). Genome sequence of *Pediococcus pentosaceus* strain IE-3. *Journal of Bacteriology*, 194, 4468.
- Moreno Switt, A. I., Andrus, A. D., Ranieri, M. L., Orsi, R. H., Ivy, R., den Bakker, H. C., Martin, N. H., Wiedmann, M., & Boor, K. J. (2014). Genomic comparison of sporeforming bacilli isolated from milk. *BMC Genomics*, 15, 26.
- Mounier, J., Monnet, C., Jacques, N., Antoinette, A., & Irlinger, F. (2009). Assessment of the microbial diversity at the surface of Livarot cheese using culture-dependent and independent approaches. *International Journal of Food Microbiology*, 133, 31–37.
- Muyzer, G., de Waal, E. C., & Uitterlinden, A. G. (1993). Profiling of complex microbial populations by denaturing gradient gel electrophoresis analysis of polymerase chain reaction-amplified genes coding for 16S rRNA. *Applied and Environmental Microbiology*, 59, 695–700.
- Nakamura, Y., Cochrane, G., Karsch-Mizrachi, I., & International Nucleotide Sequence Database, C. (2013). The international nucleotide sequence database collaboration. *Nucleic Acids Research*, 41, D21–4.
- Nam, Y. D., Lee, S. Y., & Lim, S. I. (2012a). Microbial community analysis of Korean soybean pastes by next-generation sequencing. *International Journal of Food Microbiology*, 155, 36–42.
- Nam, Y. D., Park, S. L., & Lim, S. I. (2012b). Microbial composition of the Korean traditional food "kochujang" analyzed by a massive sequencing technique. *Journal of Food Science*, 77, M250–6.
- Naser, S., Dawyndt, P., Hoste, B., Gevers, D., Vandemeulebroecke, K., Cleenwerck, I., Vancanneyt, M., & Swings, J. (2007). Identification of lactobacilli by *pheS* and *rpoA* gene sequence analyses. *International Journal of Systematic and Evolutionary Microbiology*, 57, 2777–2789.
- Naser, S., Thompson, F. L., Hoste, B., Gevers, D., Dawyndt, P., Vancanneyt, M., & Swings, J. (2005a). Application of multilocus sequence analysis (MLSA) for rapid identification of *Enterococcus* species based on *rpoA* and *pheS* genes. *Microbiology*, 151, 2141–2150.
- Naser, S., Thompson, F. L., Hoste, B., Gevers, D., Vandemeulebroecke, K., Cleenwerck, I., Thompson, C. C., Vancanneyt, M., & Swings, J. (2005b). Phylogeny and identification of enterococci by *atpA* gene sequence analysis. *Journal of Clinical Microbiology*, 43, 2224–2230.
- Nesbo, C. L., Dlutek, M., & Doolittle, W. F. (2006). Recombination in *Thermotoga*: implications for species concepts and biogeography. *Genetics*, 172, 759–769.
- Nguyen, D. T. L., Hoorde, K. V., Cnockaert, M., Brandt, E. D., Bruyne, K. D., Le, B. T., & Vandamme, P. (2013). A culture-dependent and -independent approach for the identification of lactic acid bacteria associated with the production of nem chua, a vietnamese fermented meat product. *Food Research International*, 50, 232 – 240.

- Nilsson, R. H., Veldre, V., Hartmann, M., Unterseher, M., Amend, A., Bergsten, J., Kristiansson, E., Ryberg, M., Jumpponen, A., & Abarenkov, K. (2010). An open source software package for automated extraction of ITS1 and ITS2 from fungal ITS sequences for use in high-throughput community assays and molecular ecology. *Fungal Ecology*, 3, 284–287.
- Oren, A., & Garrity, G. (2014). Then and now: a systematic review of the systematics of prokaryotes in the last 80 years. *Antonie van Leeuwenhoek*, 106, 43–56.
- Padonou, S. W., Schillinger, U., Nielsen, D. S., Franz, C. M. A. P., Hansen, M., Hounhouigan, J. D., Nago, M. C., & Jakobsen, M. (2010). *Weissella beninensis* sp. nov., a motile lactic acid bacterium from submerged cassava fermentations, and emended description of the genus *Weissella*. *International Journal of Food Microbiology*, 60, 2193–2198.
- Panelli, S., Brambati, E., Bonacina, C., & Feligini, M. (2013). Detection of *Clostridium tyrobutyricum* in milk to prevent late blowing in cheese by automated ribosomal intergenic spacer analysis. *Journal of Food Science*, 78, M1569–M1574.
- Pangallo, D., Šaková, N., KoreÁlová, J., Puškárová, A., Kraková, L., Valík, L., & Kuchta, T. (2014). Microbial diversity and dynamics during the production of May bryndza cheese. *International Journal of Food Microbiology*, 170, 38–43.
- Papalexandratou, Z., Camu, N., Falony, G., & De Vuyst, L. (2011a). Comparison of the bacterial species diversity of spontaneous cocoa bean fermentations carried out at selected farms in Ivory Coast and Brazil. *Food Microbiology*, 28, 964–973.
- Papalexandratou, Z., & De Vuyst, L. (2011b). Assessment of the yeast species composition of cocoa bean fermentations in different cocoa-producing regions using denaturing gradient gel electrophoresis. *FEMS Yeast Research*, 11, 564–574.
- Papalexandratou, Z., Falony, G., Romanens, E., Jimenez, J. C., Amores, F., Daniel, H.-M., & De Vuyst, L. (2011c). Species diversity, community dynamics, and metabolite kinetics of the microbiota associated with traditional Ecuadorian spontaneous cocoa bean fermentations. *Applied and Environmental Microbiology*, 77, 7698–7714.
- Papalexandratou, Z., Lefeber, T., Bahrim, B., Lee, O. S., Daniel, H.-M., & De Vuyst, L. (2013). *Hanseniaspora opuntiae*, *Saccharomyces cerevisiae*, *Lactobacillus fermentum*, and *Acetobacter pasteurianus* predominate during well-performed Malaysian cocoa bean box fermentations, underlining the importance of these microbial species for a successful cocoa . *Food Microbiology*, 35, 73–85.
- Papalexandratou, Z., Vrancken, G., De Bruyne, K., Vandamme, P., & De Vuyst, L. (2011d). Spontaneous organic cocoa bean box fermentations in Brazil are characterized by a restricted species diversity of lactic acid bacteria and acetic acid bacteria. *Food Microbiology*, 28, 1326–1338.
- Parameswaran, P., Jalili, R., Tao, L., Shokralla, S., Gharizadeh, B., Ronaghi, M., & Fire, A. Z. (2007). A pyrosequencing-tailored nucleotide barcode design unveils opportunities for large-scale sample multiplexing. *Nucleic Acids Research*, 35, e130.
- Park, E. J., Chun, J., Cha, C. J., Park, W. S., Jeon, C. O., & Bae, J. W. (2012). Bacterial community analysis during fermentation of ten representative kinds of kimchi with barcoded pyrosequencing. *Food Microbiology*, 30, 197–204.

- Peeters, K., Hodgson, D., Convey, P., & Willems, A. (2011). Culturable diversity of heterotrophic bacteria in Forlidas Pond (Pensacola Mountains) and Lundström Lake (Shackleton Range), Antarctica. *Microbial Ecology*, 62, 399–413.
- Petrosino, J. F., Highlander, S., Luna, R. A., Gibbs, R. A., & Versalovic, J. (2009). Metagenomic pyrosequencing and microbial identification. *Clinical Chemistry*, 55, 856–866.
- Pikuta, E. V., Marsic, D., Bej, A., Tang, J., Krader, P., & Hoover, R. B. (2005). *Carnobacterium pleistocenium* sp. nov., a novel psychrotolerant, facultative anaerobe isolated from permafrost of the Fox Tunnel in Alaska. *International Journal of Systematic and Evolutionary Microbiology*, 55, 473–478.
- Pires, E., Teixeira, J., Brányik, T., & Vicente, A. (2014). Yeast: the soul of beer's aroma—a review of flavour-active esters and higher alcohols produced by the brewing yeast. *Applied Microbiology and Biotechnology*, 98, 1937–1949.
- Pitcher, D. G., Saunders, N. A., & Owen, R. J. (1989). Rapid extraction of bacterial genomic DNA with guanidium thiocyanate. *Letters in Applied Microbiology*, 8, 151–156.
- Pittet, V., Abegunde, T., Marfleet, T., Haakensen, M., Morrow, K., Jayaprakash, T., Schroeder, K., Trost, B., Byrns, S., Bergsveinson, J., Kusalik, A., & Ziola, B. (2012). Complete genome sequence of the beer spoilage organism *Pediococcus claussenii* ATCC BAA-344<sup>T</sup>. *Journal of Bacteriology*, 194, 1271–1272.
- Pittet, V., Morrow, K., & Ziola, B. (2011). Ethanol tolerance of lactic acid bacteria, including relevance of the exopolysaccharide gene *gtf*. *Journal of the American Society of Brewing Chemists*, 69, 57–61.
- Pittet, V., Phister, T. G., & Ziola, B. (2013). Transcriptome sequence and plasmid copy number analysis of the brewery isolate *Pediococcus claussenii* ATCC BAA-344<sup>T</sup> during growth in beer. *PLoS ONE*, 8, e73627.
- Pothakos, V., Snauwaert, C., De Vos, P., Huys, G., & Devlieghere, F. (2014). Psychrotrophic members of *Leuconostoc gasicomitatum*, *Leuconostoc gelidum* and *Lactococcus piscium* dominate at the end of shelf-life in packaged and chilled-stored food products in Belgium. *Food Microbiology*, 39, 61–67.
- Prosser, J. I. (2010). Replicate or lie. *Environmental Microbiology*, 12, 1806–1810.
- Pruesse, E., Quast, C., Knittel, K., Fuchs, B. M., Ludwig, W., Peplies, J., & Glöckner, F. O. (2007). SILVA: a comprehensive online resource for quality checked and aligned ribosomal RNA sequence data compatible with ARB. *Nucleic Acids Research*, 35, 7188–7196.
- Qin, J., Li, R., Raes, J., Arumugam, M., Burgdorf, K. S., Manichanh, C., Nielsen, T., Pons, N., Levenez, F., Yamada, T., Mende, D. R., Li, J., Xu, J., Li, S., Li, D., Cao, J., Wang, B., Liang, H., Zheng, H., Xie, Y., Tap, J., Lepage, P., Bertalan, M., Batto, J.-M., Hansen, T., Le Paslier, D., Linneberg, A., Nielsen, H. B., Pelletier, E., Renault, P., Sicheritz-Ponten, T., Turner, K., Zhu, H., Yu, C., Li, S., Jian, M., Zhou, Y., Li, Y., Zhang, X., Li, S., Qin, N., Yang, H., Wang, J., Brunak, S., Dore, J., Guarner, F., Kristiansen, K., Pedersen, O., Parkhill, J., Weissenbach, J., Bork, P., Ehrlich, S. D., & Wang, J. (2010). A human gut microbial gene catalogue established by metagenomic sequencing. *Nature*, 464, 59–65.

- de Queiroz, K. (2005). Ernst Mayr and the modern concept of species. *Proceedings of the National Academy of Sciences of the United States of America*, 102 Suppl, 6600–6607.
- Quince, C., Lanzen, A., Curtis, T. P., Davenport, R. J., Hall, N., Head, I. M., Read, L. F., & Sloan, W. T. (2009). Accurate determination of microbial diversity from 454 pyrosequencing data. *Nature Methods*, 6, 639–641.
- Quince, C., Lanzen, A., Davenport, R. J., & Turnbaugh, P. J. (2011). Removing noise from pyrosequenced amplicons. *BMC Bioinformatics*, 12, 38.
- Rademaker, J. L. W., Herbet, H., Starrenburg, M. J. C., Naser, S. M., Gevers, D., Kelly, W. J., Hugenholtz, J., Swings, J., & van Hylckama Vlieg, J. E. T. (2007). Diversity analysis of dairy and nondairy *Lactococcus lactis* isolates, using a novel multilocus sequence analysis scheme and (GTG)<sub>5</sub>-PCR fingerprinting. *Applied and Environmental Microbiology*, 73, 7128–7137.
- Rademaker, J. L. W., Hoolwerf, J. D., Wagendorp, A. A., & te Giffel, M. C. (2006). Assessment of microbial population dynamics during yoghurt and hard cheese fermentation and ripening by DNA population fingerprinting. *International Dairy Journal*, 16, 457–466.
- Rahkila, R., De Bruyne, K., Johansson, P., Vandamme, P., & Björkroth, J. (2014). Reclassification of *Leuconostoc gasicomitatum* as *Leuconostoc gelidum* subsp. *gasicomitatum* comb. nov., description of *Leuconostoc gelidum* subsp. *aenigmaticum* subsp. nov., designation of *Leuconostoc gelidum* subsp. *gelidum* subsp. nov. and emended description. *International Journal of Systematic and Evolutionary Microbiology*, 64, 1290–1295.
- Rappé, M. S., & Giovannoni, S. J. (2003). The uncultured microbial majority. *Annual Review of Microbiology*, 57, 369–394.
- Raspor, P., & Goranovic, D. (2008). Biotechnological applications of acetic acid bacteria. *Critical Reviews in Biotechnology*, 28, 101–124.
- Ravyts, F., De Vuyst, L., & Leroy, F. (2012). Bacterial diversity and functionalities in food fermentations. *Engineering in Life Sciences*, 12, 356–367.
- Ravyts, F., Vrancken, G., D'Hondt, K., Vasilopoulos, C., De Vuyst, L., & Leroy, F. (2009). Kinetics of growth and 3-methyl-1-butanol production by meat-borne, coagulase-negative staphylococci in view of sausage fermentation. *International Journal of Food Microbiology*, 134, 89–95.
- Reddy, V. S., Shlykov, M. A., Castillo, R., Sun, E. I., & Saier, M. H. (2012). The major facilitator superfamily (MFS) revisited. *FEBS Journal*, 279, 2022–2035.
- Reeder, J., & Knight, R. (2009). The 'rare biosphere': a reality check. *Nature Methods*, 6, 636–637.
- Richardson, E. J., & Watson, M. (2012). The automatic annotation of bacterial genomes. *Briefings in Bioinformatics*, .
- Richter, M., & Rossello-Mora, R. (2009). Shifting the genomic gold standard for the prokaryotic species definition. *Proceedings of the National Academy of Sciences of the United States of America*, 106, 19126–19131.



- Robert, V., Vu, D., Amor, A. B., van de Wiele, N., Brouwer, C., Jabas, B., Szoke, S., Dridi, A., Triki, M., Ben Daoud, S., Chouchen, O., Vaas, L., de Cock, A., Stalpers, J. A., Stalpers, D., Verkley, G. J., Groenewald, M., Dos Santos, F. B., Stegehuis, G., Li, W., Wu, L., Zhang, R., Ma, J., Zhou, M., Gorjon, S. P., Eurwilaichitr, L., Ingsriswang, S., Hansen, K., Schoch, C., Robbertse, B., Irinyi, L., Meyer, W., Cardinali, G., Hawksworth, D. L., Taylor, J. W., & Crous, P. W. (2013). MycoBank gearing up for new horizons. *IMA Fungus*, 4, 371–379.
- Robinson, C. J., Bohannon, B. J., & Young, V. B. (2010). From structure to function: the ecology of host-associated microbial communities. *Microbiology and Molecular Biology Reviews*, 74, 453–476.
- Roh, S. W., Kim, K. H., Nam, Y. D., Chang, H. W., Park, E. J., & Bae, J. W. (2010). Investigation of archaeal and bacterial diversity in fermented seafood using barcoded pyrosequencing. *ISME Journal*, 4, 1–16.
- Rosselló-Mora, R. (2005). Updating prokaryotic taxonomy. *Journal of Bacteriology*, 187, 6255–6257.
- Rosselló-Mora, R. (2006). DNA-DNA reassociation methods applied to microbial taxonomy and their critical evaluation. In E. Stackebrandt (Ed.), *Molecular identification, systematics, and population structure of prokaryotes* chapter 2. (pp. 23–50). Springer, Berlin, Germany.
- Rossello-Mora, R., & Amann, R. (2001). The species concept for prokaryotes. *FEMS Microbiology Reviews*, 25, 39–67.
- Säde, E. (2011). *Leuconostoc spoilage of refrigerated, packaged foods*. Academic dissertation University of Helsinki.
- Sakamoto, K., & Konings, W. N. (2003). Beer spoilage bacteria and hop resistance. *International Journal of Food Microbiology*, 89, 105–124.
- Sakamoto, K., Margolles, A., van Veen, H. W., & Konings, W. N. (2001). Hop resistance in the beer spoilage bacterium *Lactobacillus brevis* is mediated by the ATP-binding cassette multidrug transporter HorA. *Journal of Bacteriology*, 183, 5371–5375.
- Sakamoto, N., Tanaka, S., Sonomoto, K., & Nakayama, J. (2011). 16S rRNA pyrosequencing-based investigation of the bacterial community in nukadoko, a pickling bed of fermented rice bran. *International Journal of Food Microbiology*, 144, 352–359.
- Sanchez, J. I., Rossetti, L., Martinez, B., Rodriguez, A., & Giraffa, G. (2006). Application of reverse transcriptase PCR-based T-RFLP to perform semi-quantitative analysis of metabolically active bacteria in dairy fermentations. *Journal of Microbiological Methods*, 65, 268–277.
- Sanger, F., Nicklen, S., & Coulson, A. R. (1992). DNA sequencing with chain-terminating inhibitors. *Biotechnology*, 24, 104–108.
- Santamaria, M., Fosso, B., Consiglio, A., De Caro, G., Grillo, G., Licciulli, F., Liuni, S., Marzano, M., Alonso-Aleman, D., Valiente, G., & Pesole, G. (2012). Reference databases for taxonomic assignment in metagenomics. *Briefings in Bioinformatics*, 13, 682–695.
- Sauer, S., & Kliem, M. (2010). Mass spectrometry tools for the classification and identification of bacteria. *Nature Reviews Microbiology*, 8, 74–82.

- Sboner, A., Mu, X. J., Greenbaum, D., Auerbach, R. K., & Gerstein, M. B. (2011). The real cost of sequencing: higher than you think! *Genome Biology*, 12, 125.
- Schleifer, K. H. (2009). Classification of bacteria and archaea: Past, present and future. *Systematic and Applied Microbiology*, 32, 533 – 542.
- Schleifer, K. H., & Bergan, T. (1985). Analysis of the chemical composition and primary structure of murein. In *Methods in microbiology* (pp. 123–156). Academic Press volume Volume 18.
- Schloss, P. D. (2009). A high-throughput DNA sequence aligner for microbial ecology studies. *PLoS ONE*, 4, e8230.
- Schloss, P. D. (2010). The effects of alignment quality, distance calculation method, sequence filtering, and region on the analysis of 16S rRNA gene-based studies. *PLoS Computational Biology*, 6, e1000844.
- Schloss, P. D., Gevers, D., & Westcott, S. L. (2011a). Reducing the effects of PCR amplification and sequencing artifacts on 16S rRNA-based studies. *PLoS ONE*, 6, e27310.
- Schumann, P. (2011). Peptidoglycan structure. In F. O. A. Rainey (Ed.), *Taxonomy of Prokaryotes* (pp. 101–129). Methods in Microbiology volume 38. (38th ed.).
- Seifert, K. A. (2009). Progress towards DNA barcoding of fungi. *Molecular Ecology Resources*, 9, 83–89.
- Seshadri, R., Kravitz, S. A., Smarr, L., Gilna, P., & Frazier, M. (2007). CAMERA: a community resource for metagenomics. *PLoS Biology*, 5, e75.
- Shanta Kumara, H. M. C., & Verachtert, H. (1991). Identification of lambic superattenuating micro-organisms by the use of selective antibiotics. *Journal of the Institute of Brewing*, 97, 181–185.
- Shi, Y., Tyson, G. W., Eppley, J. M., & DeLong, E. F. (2011). Integrated metatranscriptomic and metagenomic analyses of stratified microbial assemblages in the open ocean. *ISME Journal*, 5, 999–1013.
- Shimwell, J. L. (1937). On the relation between the staining properties of bacteria and their reaction towards hop antiseptic. *Journal of the Institute of Brewing*, 43, 111–118.
- Simon, F., Johan, B., Fredrik, D., Angela, C., Hanlee, J., Katrina, W., & Ronald, W. D. (2007). Multiplex amplification of all coding sequences within 10 cancer genes by Gene-Collector. *Nucleic Acids Research*, 35.
- Simpson, W. J. (1993). Ionophoric action of trans-isohumulone on *Lactobacillus brevis*. *Journal of General Microbiology*, 139, 1041–1045.
- Sleator, R. (2011). Phylogenetics. *Archives of Microbiology*, 193, 235–239.
- Smid, W. S., Hugenholtz, J., de Vos, W. M., & J., E. (2006). Safe use of genetically modified lactic acid bacteria in food. Bridging the gap between consumers, green groups, and industry. *Electronic Journal of Biotechnology*, 9, 1–27.

- Smith, M. T., Yamazaki, M., & Poot, G. A. (1990). *Dekkera*, *Brettanomyces* and *Eeniella*: Electrophoretic comparison of enzymes and DNA-DNA homology. *Yeast*, 6, 299–310.
- Snauwaert, I., Hoste, B., De Bruyne, K., Peeters, K., De Vuyst, L., Willems, A., & Vandamme, P. (2013a). *Carnobacterium iners* sp. nov., a psychrophilic, lactic acid-producing bacterium from the littoral zone of an Antarctic pond. *International Journal of Systematic and Evolutionary Microbiology*, 63, 1370–1375.
- Sogin, M. L., Morrison, H. G., Huber, J. A., Mark Welch, D., Huse, S. M., Neal, P. R., Arrieta, J. M., & Herndl, G. J. (2006). Microbial diversity in the deep sea and the underexplored "rare biosphere". *Proceedings of the National Academy of Sciences of the United States of America*, 103, 12115–12120.
- Spaepen, M., Van Oevelen, D., & Verachtert, H. (1978). Fatty acids and esters produced during the spontaneous fermentation of lambic and gueuze. *Journal of the Institute of Brewing*, 84, 278–282.
- Spaepen, M., Van Oevelen, D., & Verachtert, H. (1979). Higher fatty acid (HFA) and HFA-ester content of spontaneously fermented Belgian beers and evaluation of their analytical determination. *Brauwissenschaft*, 32, S1–S6.
- Spaepen, M., & Verachtert, H. (1982). Esterase activity in the genus *Brettanomyces*. *Journal of the Institute of Brewing*, 88, 11–17.
- Spitaels, F., Li, L., Wieme, A., Balzarini, T., Cleenwerck, I., Van Landschoot, A., De Vuyst, L., & Vandamme, P. (2014a). *Acetobacter lambici* sp. nov., isolated from fermenting lambic beer. *International Journal of Systematic and Evolutionary Microbiology*, 64, 1083–1089.
- Spitaels, F., Wieme, A., Janssens, M., Aerts, M., Daniel, H.-M., Van Landschoot, A., De Vuyst, L., & Vandamme, P. (2014b). The microbial diversity of traditional spontaneously fermented lambic beer. *PLoS ONE*, 9, e95384.
- Stackebrandt, E., & Ebers, J. (2006). Taxonomic parameters revisited: tarnished gold standards. *Microbiology Today*, 33, 152.
- Stackebrandt, E., Frederiksen, W., Garrity, G. M., Grimont, P. A. D., Kämpfer, P., Maiden, M. C. J., Nesme, X., Rossell-Mora, R., Swings, J., Träijper, H. G., Vauterin, L., Ward, A. C., & Whitman, W. B. (2002). Report of the ad hoc committee for the re-evaluation of the species definition in bacteriology. *International Journal of Systematic and Evolutionary Microbiology*, 52, 1043–7.
- Stackebrandt, E., & Goebel, B. M. (1994). Taxonomic note: a place for dna-dna reassociation and 16s rRNA sequence analysis in the present species definition in bacteriology. *International Journal of Systematic Bacteriology*, 44, 846–849.
- Staley, J. T. (2006). The bacterial species dilemma and the genomic-phylogenetic species concept. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 361, 1899–1909.
- Stiles, M. E., & Holzapfel, W. H. (1997). Lactic acid bacteria of foods and their current taxonomy. *International Journal of Food Microbiology*, 36, 1–29.
- Suárez, R., Suárez-Lepe, J., Morata, A., & Calderón, F. (2007). The production of ethylphenols in wine by yeasts of the genera *Brettanomyces* and *Dekkera*: A review. *Food Chemistry*, 102, 10–21.

- Suenaga, H. (2012). Targeted metagenomics: a high-resolution metagenomics approach for specific gene clusters in complex microbial communities. *Environmental Microbiology*, 14, 13–22.
- Sugawara, H., Abe, T., Gojobori, T., & Tateno, Y. (2007). DDBJ working on evaluation and classification of bacterial genes in INSDC. *Nucleic Acids Research*, 35, D13–5.
- Sutcliffe, I. C., Trujillo, M. E., & Goodfellow, M. (2012). A call to arms for systematists: revitalising the purpose and practises underpinning the description of novel microbial taxa. *Antonie van Leeuwenhoek*, 101, 13–20.
- Suzuki, K., Ozaki, K., & Yamashita, H. (2004). Comparative analysis of conserved genetic markers and adjacent DNA regions identified in beer-spoilage lactic acid bacteria. *Letters in Applied Microbiology*, 39, 240–245.
- Sybesma, W., Hugenholtz, J., de Vos, W. M., & Smid, E. J. (2006). Safe use of genetically modified lactic acid bacteria in food: Bridging the gap between consumers, green groups, and industry. *Electronic Journal of Biotechnology*, 9.
- Tamura, K. (1992). Estimation of the number of nucleotide substitutions when there are strong transition-transversion and g+c-content biases. *Molecular Biology and Evolution*, 9, 678–687.
- Tamura, K., Peterson, D., Peterson, N., Stecher, G., Nei, M., & Kumar, S. (2011). MEGA5: Molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Molecular Biology and Evolution*, 28, 2731–2739.
- Tanasupawat, S., Shida, O., Okada, S., & Komagata, K. (2000). *Lactobacillus acidipiscis* sp. nov. and *Weissella thailandensis* sp. nov., isolated from fermented fish in Thailand. *International Journal of Systematic and Evolutionary Microbiology*, 50, 1479–1485.
- Teeling, H., Fuchs, B. M., Becher, D., Klockow, C., Gardebrecht, A., Bennke, C. M., Kassabgy, M., Huang, S., Mann, A. J., Waldmann, J., Weber, M., Klindworth, A., Otto, A., Lange, J., Bernhardt, J., Reinsch, C., Hecker, M., Peplies, J., Bockelmann, F. D., Callies, U., Gerdt, G., Wichels, A., Wiltshire, K. H., Glöckner, F. O., Schweder, T., & Amann, R. (2012). Substrate-controlled succession of marine bacterioplankton populations induced by a phytoplankton bloom. *Science*, 336, 608–611.
- Teeling, H., & Glöckner, F. O. (2012). Current opportunities and challenges in microbial metagenome analysis – a bioinformatic perspective. *Briefings in Bioinformatics*, 13, 728–742.
- Temperton, B., & Giovannoni, S. J. (2012). Metagenomics: microbial diversity through a scratched lens. *Current Opinion in Microbiology*, 15, 605 – 612.
- Teuber, M., & Schmalreck, A. F. (1973). Membrane leakage in *Bacillus subtilis* 168 induced by the hop constituents lupulone, humulone, isohumulone and humulinic acid. *Archiv für Mikrobiologie*, 94, 159–171.
- Thompson, J. R., Marcelino, L. A., & Polz, M. F. (2002). Heteroduplexes in mixed-template amplifications: formation, consequence and elimination by 'reconditioning PCR'. *Nucleic Acids Research*, 30, 2083–2088.

- Tilden, E. B., & Svec, M. (1952). Further studies of a differential culture technique for estimations of acidogenic bacteria in saliva. II. Species of lactobacilli isolated from saliva and their distribution in a group of children. *Journal of Dental Research*, 31, 831–838.
- Timke, M., Wang-Lieu, N. Q., Altendorf, K., & Lipski, A. (2008). Identity, beer spoiling and biofilm forming potential of yeasts from beer bottling plant associated biofilms. *Antonie van Leeuwenhoek*, 93, 151–161.
- Tindall, B. J., Rossello-Mora, R., Busse, H. J., Ludwig, W., & Kampfer, P. (2010). Notes on the characterization of prokaryote strains for taxonomic purposes. *International Journal of Systematic and Evolutionary Microbiology*, 60, 249–266.
- Tiukova, I. A., Petterson, M. E., Tellgren-Roth, C., Bunikis, I., Eberhard, T., Pettersson, O. V., & Passoth, V. (2013). Transcriptome of the alternative ethanol production strain *Dekkera bruxellensis* CBS 11270 in sugar limited, low oxygen cultivation. *PLoS ONE*, 8, e58455.
- Torriani, S., Gatto, V., Sembeni, S., Tofalo, R., Suzzi, G., Belletti, N., Gardini, F., & Bover-Cid, S. (2008). Rapid detection and quantification of tyrosine decarboxylase gene (*tdc*) and its expression in gram-positive bacteria associated with fermented foods using PCR-based methods. *Journal of Food Protection*, 71, 93–101.
- Turenne, C. Y., Tschetter, L., Wolfe, J., & Kabani, A. (2001). Necessity of quality-controlled 16S rRNA gene sequence databases: identifying nontuberculous *Mycobacterium* species. *Journal of Clinical Microbiology*, 39, 3637–3648.
- Turnbaugh, P. J., & Gordon, J. I. (2008). An invitation to the marriage of metagenomics and metabolomics. *Cell*, 134, 708–713.
- Tyson, G. W., Chapman, J., Hugenholtz, P., Allen, E. E., Ram, R. J., Richardson, P. M., Solovyev, V. V., Rubin, E. M., Rokhsar, D. S., & Banfield, J. F. (2004). Community structure and metabolism through reconstruction of microbial genomes from the environment. *Nature*, 428, 37–43.
- van der Aa Kuhle, A., & Jespersen, L. (1998). Detection and identification of wild yeasts in lager breweries. *International Journal of Food Microbiology*, 43, 205–213.
- Van Oevelen, D., L'Escaille, F., & Verachtert, H. (1976). Synthesis of aroma components during the spontaneous fermentation of lambic and gueuze. *Journal of the Institute of Brewing*, 82, 322–326.
- Van Oevelen, D., Spaepen, M., Timmermans, P., & Verachtert, H. (1977). Microbiological aspects of spontaneous wort fermentation in the production of lambic and gueuze. *Journal of the Institute of Brewing*, 83, 356–360.
- Van Oevelen, D., & Verachtert, H. (1979). Slime production by brewery strains of *Pediococcus cerevisiae*. *Journal of the American Society of Brewing Chemists*, 37, 34–37.
- Vancanneyt, M., Mengaud, J., Cleenwerck, I., Vanhonacker, K., Hoste, B., Dawyndt, P., Degivry, M. C., Ringuet, D., Janssens, D., & Swings, J. (2004). Reclassification of *Lactobacillus kefirgranum* Takizawa et al. 1994 as *Lactobacillus kefiranofaciens* subsp. *kefirgranum* subsp. nov. and emended description of *L. kefiranofaciens* Fujisawa et al. 1988. *International Journal of Systematic and Evolutionary Microbiology*, 54, 551–556.

- Vandamme, P., & Peeters, C. (2014). Time to revisit polyphasic taxonomy. *Antonie Van Leeuwenhoek*, .
- Vandamme, P., Pot, B., Gillis, M., de Vos, P., Kersters, K., & Swings, J. (1996). Polyphasic taxonomy, a consensus approach to bacterial systematics. *Microbiological Reviews*, 60, 407–438.
- Varley, K. E., & Mitra, R. D. (2008). Nested patch PCR enables highly multiplexed mutation discovery in candidate genes. *Genome Research*, 18, 1844–1850.
- Vela, A. I., Fernández, A., Bernaldo de Quirós, Y., Herráez, P., Domínguez, L., & Fernández-Garayzábal, J. F. (2011). *Weissella ceti* sp. nov., isolated from beaked whales (*Mesoplodon bidens*). *International Journal of Systematic and Evolutionary Microbiology*, 61, 2758–2762.
- Verachtert, H., & Derdelinckx, G. (2005). Acidic beers: enjoyable reminiscences of the past. *Cerevisia, Belgian Journal of Brewing and Biotechnology*, 30, 38–47.
- Verachtert, H., & Iserentant, D. (1995). Properties of Belgian acid beers and their microflora. Part I. The production of gueuze and related refreshing acid beers. *Cerevisia, Belgian Journal of Brewing and Biotechnology*, 20, 37–41.
- Verstrepen, K. J., Derdelinckx, G., Dufour, J. P., Winderickx, J., Thevelein, J. M., Pretorius, I. S., & Delvaux, F. R. (2003). Flavor-active esters: adding fruitiness to beer. *Journal of Bioscience and Bioengineering*, 96, 110–118.
- Vieites, J. M., Guazzaroni, M. E., Beloqui, A., Golyshin, P. N., & Ferrer, M. (2009). Metagenomics approaches in systems microbiology. *FEMS Microbiology Reviews*, 33, 236–255.
- Vihavainen, E. J., Murros, A. E., & Björkroth, K. J. (2008). *Leuconostoc* spoilage of vacuum-packaged vegetable sausages. *Journal of Food Protection*, 71, 2312–2315.
- Walker, A. W., Duncan, S. H., Louis, P., & Flint, H. J. (2014). Phylogeny, culturing, and metagenomics of the human gut microbiota. *Trends in Microbiology*, 22, 267–274.
- Walling, E., Gindreau, E., & Lonvaud-Funel, A. (2005). A putative glucan synthase gene *dps* detected in exopolysaccharide-producing *Pediococcus damnosus* and *Oenococcus oeni* strains isolated from wine and cider. *International Journal of Food Microbiology*, 98, 53–62.
- Wang, D. G., Fan, J.-B., Siao, C.-J., Berno, A., Young, P., Sapolsky, R., Ghandour, G., Perkins, N., Winchester, E., Spencer, J., Kruglyak, L., Stein, L., Hsie, L., Topaloglou, T., Hubbell, E., Robinson, E., Mittmann, M., Morris, M. S., Shen, N., Kilburn, D., Rioux, J., Nusbaum, C., Rozen, S., Hudson, T. J., Lipshutz, R., Chee, M., & Lander, E. S. (1998). Large-scale identification, mapping, and genotyping of single-nucleotide polymorphisms in the human genome. *Science*, 280, 1077–1082.
- Wang, Q., Garrity, G. M., Tiedje, J. M., & Cole, J. R. (2007). Naïve bayesian classifier for rapid assignment of rRNA Sequences into the new bacterial taxonomy. *Applied and Environmental Microbiology*, 73, 5261–5267.
- Wang, X., Eijkemans, M. J. C., Wallinga, J., Biesbroek, G., TrzciÅŹski, K., Sanders, E. A. M., & Bogaert, D. (2012). Multivariate approach for studying interactions between environmental variables and microbial communities. *PLoS ONE*, 7, e50267.


- Wattam, A. R., Abraham, D., Dalay, O., Disz, T. L., Driscoll, T., Gabbard, J. L., Gillespie, J. J., Gough, R., Hix, D., Kenyon, R., Machi, D., Mao, C., Nordberg, E. K., Olson, R., Overbeek, R., Pusch, G. D., Shukla, M., Schulman, J., Stevens, R. L., Sullivan, D. E., Vonstein, V., Warren, A., Will, R., Wilson, M. J. C., Yoo, H. S., Zhang, C., Zhang, Y., & Sobral, B. W. (2014). PATRIC, the bacterial bioinformatics database and analysis resource. *Nucleic Acids Research*, 42, D581–D591.
- Wayne, L. G., Brenner, D. J., Colwell, R. R., Grimont, P. A. D., Kandler, O., Krichevsky, M. I., Moore, L. H., Moore, W. E. C., Murray, R. G. E., Stackebrandt, E., Starr, M. P., & Truper, H. G. (1987). Report of the ad hoc committee on reconciliation of approaches to bacterial systematics. *International Journal of Systematic Bacteriology*, 37, 463–464.
- Weiss, S., Samson, F., Navarro, D., & Casaregola, S. (2013). YeastIP: a database for identification and phylogeny of *Saccharomycotina* yeasts. *FEMS Yeast Research*, 13, 117–125.
- White, T., Bruns, T., Lee, S., & Taylor, J. (1990). Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In M. Innis, D. Gelfand, J. Shinsky, & T. White (Eds.), *PCR protocols: a guide to methods and applications* (pp. 315–322). Academic Press.
- Wieme, A., Cleenwerck, I., Van Landschoot, A., & Vandamme, P. (2012). *Pediococcus lolii* DSM 19927<sup>T</sup> and JCM 15055<sup>T</sup> are strains of *Pediococcus acidilactici*. *International Journal of Systematic and Evolutionary Microbiology*, 62, 3105–3108.
- Wieme, A., Spitaels, F., Aerts, M., De Bruyne, K., Van Landschoot, A., & Vandamme, P. (2014). Effects of growth medium on matrix-assisted laser desorption-ionization time of flight mass spectra: a case study of acetic acid bacteria. *Applied and Environmental Microbiology*, 80, 1528–1538.
- Wolf, Y. I., Rogozin, I. B., Grishin, N. V., Tatusov, R. L., & Koonin, E. V. (2001). Genome trees constructed using five different approaches suggest new major bacterial clades. *BMC Evolutionary Biology*, 1, 8.
- Xufre, A., Albergaria, H., Inacio, J., Spencer-Martins, I., & Girio, F. (2006). Application of fluorescence *in situ* hybridisation (FISH) to the analysis of yeast population dynamics in winery and laboratory grape must fermentations. *International Journal of Food Microbiology*, 108, 376–384.
- Yasuhara, T., Yuuki, T., & Kagami, N. (2001). Novel quantitative method for detection of *Pectinatus* using rRNA targeted fluorescent probes. *Journal of the American Society of Brewing Chemists*, 59, 121–177.
- Yilmaz, P., Gilbert, J. A., Knight, R., Amaral-Zettler, L., Karsch-Mizrachi, I., Cochrane, G., Nakamura, Y., Sansone, S. A., Glockner, F. O., & Field, D. (2011). The genomic standards consortium: bringing standards to life for microbial ecology. *ISME Journal*, 5, 1565–1567.
- Yilmaz, P., Parfrey, L. W., Yarza, P., Gerken, J., Pruesse, E., Quast, C., Schweer, T., Peplies, J., Ludwig, W., & Glöckner, F. O. (2013). The SILVA and "All-species Living Tree Project (LTP)" taxonomic frameworks. *Nucleic Acids Research*, .
- Yoon, B., Jang, S., & Chang, H.-I. (2011). Sequence analysis of the *Lactobacillus* temperate phage Sha1. *Archives of Virology*, 156, 1681–1684.
- You, Z., Yongjie, L., Karlene, H. L., Jonathan, J. D., & David, S. W. (2011). PHAST: a fast phage search tool. *Nucleic Acids Research*, 39, W347–W352.

- Zdobnov, E. M., & Apweiler, R. (2001). InterProScan - an integration platform for the signature-recognition methods in InterPro. *Bioinformatics*, 17, 847–848.
- Ze, X., Le Mougou, F., Duncan, S. H., Louis, P., & Flint, H. J. (2013). Some are more equal than others: the role of 'keystone' species in the degradation of recalcitrant substrates. *Gut microbes*, 4, 236–40.
- Zerbino, D. R., & Birney, E. (2008). Velvet: Algorithms for de novo short read assembly using de Bruijn graphs. *Genome Research*, 18, 821–829.
- Zhang, Z. G., Ye, Z. Q., Yu, L., & Shi, P. (2011). Phylogenomic reconstruction of lactic acid bacteria: an update. *BMC Evolutionary Biology*, 11.



## Part VI

### Curriculum Vitae







## Isabel Snauwaert

Gulden-Peerdenstraat 45 • 8310 Assebroek • Belgium

Phone: +32 (0) 471 87 12 54 • E-mail: [isabelsnauwaert@hotmail.com](mailto:isabelsnauwaert@hotmail.com)

Nationality: Belgian • Date of birth: the 29<sup>th</sup> of April 1987

Married with Bram Gilté

### Objective

---

I am an **ambitious** and **result-oriented team player** with a background in Biotechnology and experience in Genomics, Microbiology, and Bioinformatics. My objective is to work in an **international** and **challenging scientific environment** where my data handling, organizational, and communication skills can be effectively employed.

### Experience

---

I am currently finishing my PhD project under the supervision of Prof. Dr. Peter Vandamme at the Laboratory of Microbiology, Ghent University and Prof. dr. ir. Luc De Vuyst at the Research Group of Industrial Microbiology and Food Biotechnology, Vrije Universiteit Brussel.

I carried out research on different aspects of microbiology, including microbial diversity analysis, comparative genomics, metabolite target analysis, taxonomy, and microbial ecology. I am familiar with the following methods: next-generation sequencing technologies and data handling, comparative genomics, mass spectrometry, chromatography, DNA extraction, cultivation, phylogenetic analysis using different software packages, PCR-based methods, and other. Details on my experience can be found below.

### Education

---

Master of Biochemistry and Biotechnology (*magna cum laude*), Ghent University

- Specialization: Microbial biotechnology and biomedical biotechnology

Bachelor of Biochemistry and Biotechnology (*cum laude*), Ghent University

Secondary school degree, Sint-Andreaslyceum Sint-Kruis, Bruges

- Specialization in science and mathematics

### Skills

---

#### Language skills

- Dutch: mother tongue; English: proficient user; French: basic user; Portuguese: basic user

#### Social and organizational skills

- Team work through experience in various disciplines including the youth movement (Chiro: member for 14 years of which 6 years as a leader), scientific popularization ('Dag van de wetenschap': 2012 and 2013), and other.

#### Computer skills

- Excellent command of Microsoft Windows and Office (Excel, Word, PowerPoint and Outlook); mothur software package (command line interface); Internet applications
- Command of R software package
- Basics of Unix, Latex, High-Performance computing, and Python

#### Other skills

- Sports: triathlon (Olympic distance), running, bike racing, swimming, 4 years safeguard at the beach, 2 years safeguard in an indoor swimming pool; hobby brewer

## Experience: details

---

### September 2010- now

Analysis of the microbial diversity and metabolite composition of Belgian acidic ales

- Knowledge on DNA extraction, PCR optimizations, 454 pyrosequencing, data analysis using the mothur software package, and statistical analysis using the R software package. Contributed to the first use of this technology and its resulting data analysis in the laboratory.
- Collaboration with the laboratory of Prof. Dr. Ir. Luc De Vuyst (Research Group of Industrial Microbiology and Food Biotechnology, Department of Bioengineering Sciences, Vrije Universiteit Brussel) and Prof. Dr. Anita Van Landschoot (Laboratory of Biochemistry and Brewing, Faculty of Bioscience Engineering, Ghent University)
- Example publication (not yet submitted)
  - Snauwaert I, Roels S, Van Nieuwerburg F, Van Landschoot A, De Vuyst L, Vandamme P (2014) Microbial diversity and metabolite composition of Belgian red-brown acidic ales. In preparation.

Comparative genome analysis of a *Pediococcus damnosus* strain isolated from beer

- Knowledge on Illumina sequencing, data analysis using different open source resources (RAST, IMG-RE, PGAAP, MAUVE, ACT) and CLC genomics workbench, interpretation of the annotations found in the genome sequences, and other.
- Example publication (not yet submitted)
  - Snauwaert I, Stragier P, De Vuyst L, Vandamme P (2014) Comparative genome analysis of *Pediococcus damnosus* LMG 28219, a strain well-adapted to the brewery environment. In preparation.

Taxonomic study

- Knowledge on mass spectrometry, multi locus sequence analysis, phylogeny, general cultivation, and other. Contributed to the setting up of a mass spectral database containing protein profiles of lactic acid bacteria.
- Example publications
  - Snauwaert I, Hoste B, De Bruyne K, Peeters K, De Vuyst L, et al. (2013) *Carnobacterium iners* sp. nov., a psychrophilic, lactic acid-producing bacterium from the littoral zone of an Antarctic pond. International Journal of Systematic and Evolutionary Microbiology 63: 1370-1375.
  - Snauwaert I, Papalexandratou Z, De Vuyst L, Vandamme P (2013) Characterization of strains of *Weissella fabalis* sp. nov. and *Fructobacillus tropaeoli* from spontaneous cocoa bean fermentations. International Journal of Systematic and Evolutionary Microbiology 63: 1709-1716.

#### Collaboration to other projects

- Spitaels F, Van Kerrebroeck S, Wieme A, Snauwaert I, Aerts M, Van Landschoot A, De Vuyst L, Vandamme P (2014). Microbiota and metabolites of aged bottled gueuze beers converge to the same composition. In preparation.
- Lyhs U, Snauwaert I, Pihlajaviita S, De Vuyst L, Vandamme P (2014) *Leuconostoc rapi* sp. nov., isolated from sous-vide cooked rutabaga. In preparation.
- Huch M, De Bruyne K, Cleenwerck I, Bub A, Cho G, Watzl B, Snauwaert I, Franz C, Vandamme P (2013) *Streptococcus rubneri* sp. nov., isolated from the human throat. International Journal of Systematic and Evolutionary Microbiology 63:4026-4032
- Nguyen D, Cnockaert M, Van Hoorde K, De Brandt E, Snauwaert I, Snauwaert C, De Vuyst L, Thanh Le B, Vandamme P (2012) *Lactobacillus porcinae* sp. nov. isolated from traditional Vietnamese nem chua. International Journal of Systematic and Evolutionary Microbiology 0.044123-0

#### Guidance of students

- Mieke Vandewalle and Annelies Torfs (2<sup>nd</sup> Master in Biochemistry and Biotechnology); Elisa Dobbelaere (Master Pharmaceutical sciences); Serge Tambue (1<sup>st</sup> Master Biochemistry and Biotechnology); several practical courses.

#### Workshops followed in the scope of the PhD

- Mothur workshop 2012 and R workshop 2013, Detroit area (Michigan, USA) organized by Dr. Patrick Schloss; COST action workshops on 'Microbial ecology & the earth system: collaboration for insight and success with the new generation of sequencing tools' in Liverpool and 'Workshop on Linking microbial ecology & evolution' in Berlin.

#### Courses followed in the scope of the PhD (doctoral program Ghent University)

- Getting started with high-performance computing (included basics of Unix and Python); Introduction to Statistics; Brewing technology
- Advanced academic English: writing skills; Communication skills: Conflict handling and Meeting skills; Project management; Academic posters

#### Posters on symposia

- Mass spectrometry in Food and Feed (9 June 2011), Merelbeke, Belgium
- 10th symposium on lactic acid bacteria (28 August -1 September 2011), Egmond aan zee, The Netherlands
- ISME 14 congress (21-24 of august 2012), Copenhagen, Denmark

#### January 2010- June 2010

#### 2<sup>nd</sup> Master thesis

- 'Gut microbiota of Cystic fibrosis patients and their healthy siblings' under the supervision of Prof. Dr. Peter Vandamme (Laboratory of Microbiology, Ghent University)

